

## OKLAHOMA DEPARTMENT OF CORRECTIONS

### SEVERITY CLASSIFICATION OF COMMON CHRONIC ILLNESS

Chronic Illness	Disease Severity		
	Mild	Moderate	Severe
<b>Diabetes</b>	<ul style="list-style-type: none"> <li>HGBA1C &lt; 7.0</li> <li>No evidence of diabetic complications</li> <li>No hospitalization within year</li> </ul>	<ul style="list-style-type: none"> <li>HGBA1C &lt; 7.0 – 10.0</li> <li>No disabling diabetic complications</li> <li>No hospitalization within year</li> </ul>	<ul style="list-style-type: none"> <li>HGBA1C &gt; 10.0</li> <li>Disabling diabetic complications</li> <li>Hospitalized within last 6 months</li> </ul>
<b>Asthma</b>	<ul style="list-style-type: none"> <li>Peak flow 80% of personal best</li> <li>Rescue inhaler use &lt; 6 times/week</li> <li>No hospitalization within year</li> </ul>	<ul style="list-style-type: none"> <li>Peak flow 60% - 80% of personal best</li> <li>Daily rescue inhaler use</li> <li>No hospitalization within year</li> </ul>	<ul style="list-style-type: none"> <li>Peak flow &lt; 60% of personal best</li> <li>Rescue inhaler use several times daily</li> <li>Hospitalized within last 6 months</li> </ul>
<b>Hypertension</b>	<ul style="list-style-type: none"> <li>Systolic BP 120 – 139 to</li> <li>Diastolic BP 80 – 89</li> </ul>	<ul style="list-style-type: none"> <li>Systolic BP 140 – 159 to</li> <li>Diastolic BP 90 – 99</li> </ul>	<ul style="list-style-type: none"> <li>Systolic BP &gt; 160</li> <li>Diastolic BP &gt; 100</li> </ul>
	<b>The above readings are on three separate occasions in spite of adequate doses of antihypertensive medications.</b>		
<b>Seizure Disorder</b>	<ul style="list-style-type: none"> <li>History of seizures</li> <li>No current seizure medications</li> <li>No seizure in the past 12 months</li> <li>No hospitalization within year</li> </ul>	<ul style="list-style-type: none"> <li>Currently on seizure medications</li> <li>No seizure in the past 6 months</li> <li>No hospitalization within year</li> </ul>	<ul style="list-style-type: none"> <li>Seizure activity in the past 6 months</li> <li>History of status epilepticus in the past 12 months</li> <li>Hospitalized within last 6 months</li> </ul>
<b>Coronary Artery Disease</b>	<ul style="list-style-type: none"> <li>No chest pain with unusual activities</li> <li>With or without medications</li> <li>No hospitalization within year</li> </ul>	<ul style="list-style-type: none"> <li>Chest pain with exertion</li> <li>Relieved with medications</li> <li>No hospitalization within year</li> <li>Abnormal cardiac imaging</li> </ul>	<ul style="list-style-type: none"> <li>Chest pain with minimal activity</li> <li>MI or bypass surgery within past 6 months</li> <li>Hospitalized within last 6 months</li> </ul>
<b>HIV Infection</b>	<ul style="list-style-type: none"> <li>Viral load undetectable</li> <li>CD4 &gt; 350</li> <li>No history of opportunistic infections</li> <li>No hospitalization within year</li> </ul>	<ul style="list-style-type: none"> <li>Viral load &lt; 200,000</li> <li>CD4 50 – 350</li> <li>History of opportunistic infections currently well-controlled</li> <li>No hospitalization within year</li> </ul>	<ul style="list-style-type: none"> <li>Viral load &gt; 750,000</li> <li>CD4 &lt; 50</li> <li>Uncontrolled opportunistic infections</li> <li>Hospitalized within last 6 months</li> </ul>
<b>Chronic Obstructive Pulmonary Disease</b>	<ul style="list-style-type: none"> <li>No dyspnea with daily activities</li> <li>No hospitalization within year</li> </ul>	<ul style="list-style-type: none"> <li>Dyspnea with daily activities</li> <li>Requires routine steroids</li> <li>No hospitalization within year</li> </ul>	<ul style="list-style-type: none"> <li>Oxygen dependent</li> <li>Hospitalized within last 6 months</li> </ul>
<b>Congestive Heart Failure</b>	<ul style="list-style-type: none"> <li>No symptoms with daily activities</li> <li>No hospitalization within year</li> </ul>	<ul style="list-style-type: none"> <li>Dyspnea with daily activities</li> <li>No hospitalization within year</li> </ul>	<ul style="list-style-type: none"> <li>Dyspnea with minimal activity or at rest</li> <li>Hospitalized within last 6 months</li> </ul>
<b>Hepatitis B</b>	<ul style="list-style-type: none"> <li>APRI: &lt; 0.4-0.69</li> <li>FIB-4: &lt; 0.57 – 1.45</li> <li>Metavir 0, 1</li> <li>Fibrosure: F0-F1</li> <li>No stigmata of liver disease</li> </ul>	<ul style="list-style-type: none"> <li>APRI :0.7 – 1.06</li> <li>FIB-4: 1.46-2.32</li> <li>Metavir 2</li> <li>Fibrosure: F2</li> <li>No stigmata of liver disease</li> </ul>	<ul style="list-style-type: none"> <li>APRI: &gt; 1.06</li> <li>FIB-4: &gt; 2.32</li> <li>Metavir 3, 4</li> <li>Fibrosure: F3-F4</li> <li>Stigmata of liver disease</li> </ul>
<b>Hepatitis C</b>	<ul style="list-style-type: none"> <li>APRI: &lt; 0.4-0.69</li> <li>FIB-4: &lt; 0.57 – 1.45</li> <li>Metavir 0, 1</li> <li>Fibrosure: F0-F1</li> <li>No stigmata of liver disease</li> </ul>	<ul style="list-style-type: none"> <li>APRI :0.7 – 1.06</li> <li>FIB-4: 1.46-2.32</li> <li>Metavir 2</li> <li>Fibrosure: F2</li> <li>No stigmata of liver disease</li> </ul>	<ul style="list-style-type: none"> <li>APRI: &gt; 1.06</li> <li>FIB-4: &gt; 2.32</li> <li>Metavir 3, 4</li> <li>Fibrosure: F3-F4</li> <li>Stigmata of liver disease</li> </ul>
<b>Cirrhosis</b>	<ul style="list-style-type: none"> <li>Metavir 4</li> <li>Fibrosure: F4</li> <li>Child Pugh score 5 -6 (Class A)</li> </ul>	<ul style="list-style-type: none"> <li>Metavir 4</li> <li>Fibrosure: F4</li> <li>Child Pugh score 7 -8 (Class B)</li> </ul>	<ul style="list-style-type: none"> <li>Metavir 4</li> <li>Fibrosure: F4</li> <li>Child Pugh score 9 -15 (Class C)</li> </ul>