

LINCOLN COUNTY

DRUG OVERDOSE DEATHS¹ (2018-2022)

NUMBER OF UNINTENTIONAL DRUG OVERDOSE DEATHS BY TYPE OF SUBSTANCE*

32

total number of unintentional drug overdose deaths

22

Methamphetamine

8

Other Opioids (excluding fentanyl)

6

Fentanyl

<5

Benzodiazepines

0

Alcohol



0

Cocaine

* Categories are not mutually exclusive. Deaths may involve more than one substance.

From 2018-2022, **Lincoln County** had the **40th highest** drug overdose death rate in the state.

Adults aged 45-54 years had the highest rate of death.




53% of deaths occurred at a home or apartment.

NON-HISPANIC WHITE Lincoln County residents had the **highest** rate of death.

- Males were **2.6 times more likely** to die than females.
- Drug overdose **death rates increased 7%** from 17.3 per 100,000 (2013-2017) to 18.5 per 100,000 (2018-2022).
- **59%** of people who died had a history of substance use.
- **19%** of people who died had a history of mental health problems.

UNINTENTIONAL DRUG OVERDOSE DEATH RATES¹ BY COUNTY OF RESIDENCE²

► OKLAHOMA, 2018-2022



¹ Some rates are based on small numbers and may be unstable.

² County of residence was unknown for 35 decedents.

Data Source: Fatal Unintentional Poisoning Surveillance System, OSDH Injury Prevention Service

DRUG OVERDOSE HOSPITALIZATIONS[†] (2020-2022)

Lincoln County
had the 22nd
highest nonfatal
drug overdose
hospitalization
rate in the state.



**TEENS AND
ADULTS AGED
15-24**
had the highest
hospitalization
rate.



There were
105 Lincoln
County
Residents

hospitalized for a nonfatal drug overdose from 2020-2022.

- **46%** of hospitalizations were unintentional and **50%** due to self harm.
- Females were **2.3 times more likely** to be hospitalized than males.
- Drug overdose **hospitalization rates decreased 5%** from 107 per 100,000 (2017-2019) to 101.9 per 100,000 (2020-2022).
- Non-Hispanic White Lincoln County residents had the **highest rate** of hospitalizations.


The most common substances involved in hospitalizations were:

- ▶ Non-opioid pain medications
- ▶ Opioids
- ▶ Antidepressants
- ▶ Stimulants
- ▶ Antipsychotics

[†] Includes all intents of nonfatal drug overdose – unintentional, self harm, undetermined, and assault.

NONFATAL DRUG OVERDOSE HOSPITALIZATION RATES³ BY COUNTY OF RESIDENCE⁴

► OKLAHOMA, 2020-2022



³ Some rates are based on small numbers and may be unstable.

⁴ County of residence was unknown for 15 patients.

Data Source: OSDH Center for Health Statistics, Oklahoma Hospital Discharge Data
Compiled by: OSDH Injury Prevention Service

This publication was supported by Cooperative Agreement 6NU17CE0010188-01 funded by the Centers for Disease Control and Prevention. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention or the Department of Health and Human Services. This publication was issued by the Oklahoma State Department of Health (OSDH), an equal opportunity employer and provider. A digital file has been deposited with the Publications Clearinghouse of the Oklahoma Department of Libraries in compliance with section 3-114 of Title 65 of the Oklahoma Statutes and is available for download at documents.ok.gov. | oklahoma.gov/health | August 2024



OKLAHOMA
State Department
of Health