

Tobacco Use Patterns and Burden among Oklahomans 2025 Update (2011 – 2023 Trend Data)



Tobacco Use Patterns and Burden among Oklahomans 2025 Update



Prepared by

Nasir Mushtaq, PhD

Shirley James, PhD

Laura A. Beebe, PhD

Department of Biostatistics and Epidemiology

Hudson College of Public Health

University of Oklahoma Health Sciences Center

Prepared for

Oklahoma State Department of Health

Chronic Disease Prevention Service

March 27, 2025

Table of Contents

	Page
Key Findings	4
Background	5
Methods	6
Cigarette smoking	10
Current cigarette smoking	10
Patterns of cigarette smoking: Someday and Everyday Smoking	22
Former cigarette smokers	44
Smokeless tobacco use	56
Electronic Nicotine (E-cigarette or EC) product use	68
Quit attempts	80
Secondhand smoke exposure at home	92
Appendix A: Tables	104

Key Findings

- This updated report includes data on tobacco use, quit behaviors and secondhand smoke exposure among Oklahomans from 2011 – 2023.
- In Oklahoma, the prevalence of cigarette smoking overall was 15.8% in 2023. Since 2011 there has been a statistically significant average annual decline in smoking prevalence of 3.8%. Everyday smoking among Oklahomans has declined by an average of 4% each year since 2011.
- Education and income continue to demonstrate strong inverse associations with current smoking – those with more education and higher incomes have significantly lower smoking prevalence rates.
- The prevalence of current smoking continues to be substantially higher for individuals with Frequent Mental Distress (FMD) compared to those without FMD. Similar trends were observed for individuals with 14 or more days of poor physical health.
- The prevalence of current smoking was consistently higher among those who reported one or more disabilities. In 2023, more than 22% of Oklahoma adults with at least one disability were current smokers.
- In 2023, the prevalence of smokeless tobacco use was 6.3%, somewhat higher than the 5.7% observed in 2022. Males living in rural Oklahoma were twice as likely to use the product (13.8%) compared to males living in urban Oklahoma (6.9%).
- In 2023, electronic nicotine products (E-cigarettes or ECs) were used by 8.5% of Oklahomans in the past month. The prevalence of EC use was highest among those 18-24 years (26.7%) compared to those 55+ years (average of 2%). EC use was also more than twice as high for Oklahomans with Frequent Mental Distress (FMD, 17.1%) compared to those without (6.9%).
- In 2023, 51.5% of Oklahoma smokers reported a quit attempt in the last year. This proportion has significantly changed over the past 12 years by an average annual decrease of 1.6%.
- Younger Oklahoma smokers were more likely to report a quit attempt in the last year compared to older Oklahomans. In 2023, 60.6% of those 18-24 years reported a quit attempt compared to 43.0% of those 65+ years.
- The prevalence of quit attempts in the past year was higher for those with poor physical health (65.0%) compared to those with better physical health (48.4%) as well as for those with FMD (61.8%) compared to those without (47.2%).
- Similar to 2021, 87.2% of Oklahomans have a ban on smoking in the home.

Background

- Although the prevalence of smoking has been decreasing over time, certain population subgroups continue to smoke at much higher rates than the general population.
- Differences in tobacco-related behaviors have been demonstrated by sex, age, race/ethnicity, education, income, area of residence, physical and mental health status, disability status and health insurance coverage. This report explores differences in patterns of cigarette smoking, smokeless tobacco use, e-cigarette use, past quit attempts, and rules about smoking inside the home in these population subgroups and over time.
- Identifying and eliminating tobacco-related differences among groups of Oklahomans is one of the four goals for comprehensive tobacco control programs. Identifying differences in these groups of Oklahomans and examining trends over time is essential for Oklahoma to continue to make progress in reducing the tobacco-related disease burden in the state.

Methods

- This report evaluates trends in tobacco-related behaviors among Oklahomans from 2011 to 2023. The report examines variations in tobacco use prevalence, cigarette smoking patterns, former smoking rates, past quit attempts, and rules about secondhand smoke (SHS).
- Data used in this report were obtained from Oklahoma Behavioral Risk Factor Surveillance System (BRFSS). Information was obtained from 2011 to 2023 BRFSS surveys.
- Cigarette smoking was defined as having smoked at least 100 cigarettes during one's lifetime and smoking cigarettes everyday or somedays during the past 30 days.
- Cigarette smoking pattern was separated into two groups, everyday smokers and someday smokers.
- Smokeless tobacco (ST) use was defined as current use of chewing tobacco, snuff, or snus. ST use prevalence was low among women (2.0%); therefore, subgroup analysis of ST use was restricted to male ST users only.

Methods

- From 2017 to 2023, electronic nicotine product (E-cigarette or EC) use was a state-added question, and it differed slightly from data previously collected. Therefore, only the 2017 to 2023 state-added questions were used to create tables and figures related to EC use. Current EC use in this report reflects everyday or someday use in the past 30 days.
- A quit attempt in the past 12 months was defined as stopping smoking for one day or longer because the smoker was trying to quit.
- Secondhand smoke (SHS) exposure was assessed by self-reported smoking restrictions at home. This question was not asked in 2022. Home smoking ban indicated that smoking was never allowed inside one's home
- Racial/ethnic groups were defined as non-Hispanic (NH) White, NH-Black, NH-American Indian/Alaska Native, NH-Other, and Hispanic. NH-Other group included NH-Asian, NH-Hawaiian or Pacific Islander, NH-Other, and NH-Multiracial.

Methods

- Age was grouped into six categories: 18-24, 25-34, 35-44, 45-54, 55-64, and 65 years or older.
- Income was categorized as Less than \$15,000, \$15,000 to \$24,999, \$25,000 to \$34,999, \$35,000 to \$49,999, and \$50,000 or more.
- Education had four levels: less than high school, high school, some college, and college graduate or more.
- Area of residence, Urban/Rural, was defined based on geographic region: BRFSS Regions 3 and 4 (Tulsa and Central which includes Oklahoma and Cleveland counties) were considered Urban; whereas regions 1, 2, 5, and 6 (Northwest, Northeast, Southwest, and Southeast) were categorized as Rural.
- Poor physical health was defined as having 14 or more days of poor physical health in the past 30 days. Having 14 or more days of poor mental health in the past 30 days was referred to as Frequent Mental Distress (FMD)
- Disability status was defined as having one or more of the six disability types including, cognitive disability, hearing disability, mobility disability, vision disability, self-care disability, and independent living disability. Disability status data were available from 2016 to 2023.

Methods

- Because of small sample sizes for some subgroups, three-year rolling averages were calculated for age, race/ethnicity, education, and income. The rolling-averages provided more reliable estimates to express and compare prevalence proportions. This approach is consistent with CDC guidance on using BRFSS data.
- Weighted analysis was conducted using the SAS Proc Surveyfreq procedure. Weighted estimates obtained were representative of the Oklahoma population and permitted across year comparisons.
- 95% confidence intervals (CIs) were also calculated for the weighted estimates (%).
- Joinpoint regression modelling was applied to examine continuous changes in the estimates over the period of 2011-2023. To describe linear trends for this period, the estimated annual percent change (APC) was computed. Year was used as a regressor used, and p-values were obtained for significant APC. For regression function, ordinary least squares (OLS) method was used, and assumptions of constant variance and uncorrelated errors were made.
- Statistically significant values ($p < 0.05$) are noted with an asterisk.


RESULTS: CIGARETTE SMOKING

CURRENT CIGARETTE SMOKING

Prevalence of Current Cigarette Smoking among Adults

Although the adult smoking prevalence remains higher than the national average, Oklahoma has experienced a significant decline in smoking over time. The smoking prevalence in Oklahoma declined an average of 3.8% each year between 2011 and 2023, and this change was statistically significant.* In 2023, the percentage of Oklahoma adults who reported smoking (15.8%) did not change significantly from the 2022 value of 15.6%.


**Figure 1.1 Prevalence (%) of current cigarette smoking among Oklahoma adults
(Oklahoma BRFSS 2011 – 2023)**



Prevalence of Current Cigarette Smoking among Adults by Sex

The prevalence of current cigarette smoking among Oklahoma adults has consistently remained higher for males compared to females, although the differences each year were not always statistically significant. In 2023, the prevalence of smoking was 16.4% for males and 15.3% for females. Smoking prevalence has been on a downward trend for both males and females, with a statistically significant average annual percent decrease of 4.0% for males and 3.5% for females.


Figure 1.2 Prevalence (%) of current cigarette smoking among Oklahoma adults by sex (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Cigarette Smoking among Adults by Age

The 2021-23 prevalence of smoking was highest among those 35-44 years old (22.0%) followed by the 55-64-year age group (21.5%). Since 2011, current smoking has not changed substantially for the 55+ age groups. Those ages 18-24 have the lowest smoking prevalence rates (10%) and have experienced the most significant decline in prevalence rates since 2011 (APC -10.5%). Prevalence rates since 2011 have also changed significantly for those 25-34 years (-6.0%), 35-44 years (-2.0%), and those 45-54 years (-3.4%).


Figure 1.3 Three-year rolling prevalence of current cigarette smoking among Oklahoma adults by age (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Cigarette Smoking among Adults by Race/Ethnicity

The prevalence of smoking in Oklahoma varies by race/ethnicity. Non-Hispanic American Indians/Alaska Natives (NH-AI/ANs) have historically had the highest smoking prevalence rates and did again in 2021-23 (23.0%). The lowest prevalence of smoking occurred in those of Hispanic ethnicity (11.4%). All racial/ethnic groups have experienced statistically significant declines in smoking prevalence since 2011.


Figure 1.4 Three-year rolling prevalence of current cigarette smoking among Oklahoma adults by race/ethnicity (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Cigarette Smoking among Adults by Education

A strong association between education and current smoking has been consistent during the past 13 years. The prevalence of smoking among Oklahomans with less than a high school diploma (29.4%) remains much higher than those with a high school diploma (18.9%), some college (16.1%) or a college degree (6.0%). Since 2011, all declines in smoking prevalence by education were statistically significant.


Figure 1.5 Three-year rolling prevalence of current cigarette smoking among Oklahoma adults by education (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Cigarette Smoking among Adults by Income

Similar to the gradient observed with education, Oklahomans with low income have higher smoking prevalence compared to groups with higher income. In 2021-23, the prevalence of current smoking was 32.3% among individuals earning less than \$15,000 per year compared to 10.4% among those earning \$50,000 or more. Current smoking in all income categories, except for the \$25,000 to less than \$35,000 category, have declined significantly over the last 13 years.


Figure 1.6 Three-year rolling prevalence of current cigarette smoking among Oklahoma adults by Income (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Cigarette Smoking among Adults by Area of Residence

In 2011, the prevalence of smoking among Oklahomans who reported a rural area of residence was almost the same as those who reported an urban area of residence (26.4% and 25.7%, respectively). Since then, Oklahomans living in rural areas have reported a higher smoking prevalence compared to those in urban areas. In 2023, the prevalence of smoking in rural areas was 18.0% (up from 17.3% in 2022) compared to 13.0% in urban areas. Both groups had significant average annual declines during the 13-year period.


Figure 1.7 Prevalence (%) of current cigarette smoking among Oklahoma adults by area of residence (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Cigarette Smoking among Adults by Physical Health

The prevalence of smoking among individuals with poor physical health has been consistently higher than among those with better physical health. Of individuals who reported more than 14 poor physical health days during the past month, 26.6% also reported current cigarette smoking, compared to 13.7% of those with fewer than 14 poor physical health days. The downward trend was statistically significant for both groups but greater for those with fewer than 14 poor physical health days (-4.4% vs. -2.1%).


Figure 1.8 Prevalence (%) of current cigarette smoking among Oklahoma adults by physical health status (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Cigarette Smoking among Adults by Mental Health

Similar trends in smoking prevalence were observed for Oklahomans with poor mental health that were seen in those with poor physical health. In 2023, 25.9% of individuals reporting Frequent Mental Distress (FMD), or more than 14 days of poor mental health in the past month, were current smokers compared to 13.6% of those without FMD. Both groups experienced statistically significant average annual percent declines (4.8% and 4.0% respectively) in current smoking since 2011.


Figure 1.9 Prevalence (%) of current cigarette smoking among Oklahoma adults who had Frequent Mental Distress (FMD) (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Cigarette Smoking among Adults by Health Insurance Coverage

There is a consistent difference in smoking prevalence by health insurance status. Oklahomans who report having no health insurance coverage have a smoking prevalence that is significantly higher than those who report having insurance, which includes private insurance, Medicare or Medicaid, or Indian Health Service. In 2023, smoking prevalence was 21.8% for those without insurance, compared to 15.3% for those with insurance. A statistically significant average annual percent decrease was observed in both those without insurance (-5.3%) and in those with insurance (-2.7%) over the last 13 years.


Figure 1.10 Prevalence (%) of current cigarette smoking among Oklahoma adults by health insurance coverage (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Cigarette Smoking among Adults by Disability

The prevalence of current smoking is consistently higher among those who have one or more disabilities compared to those without a disability. In 2023, 22.6% of those with one or more disabilities were current smokers compared to 11.3% of those without a disability. A statistically significant average annual percent decrease was observed in both those with a disability (-3.6%) and in those without a disability (-4.9%) over the last 13 years.

Figure 1.11 Prevalence (%) of current cigarette smoking among Oklahoma adults by disability (Oklahoma BRFSS 2016 – 2023)




RESULTS: CIGARETTE SMOKING

PATTERNS OF CIGARETTE SMOKING

Prevalence of Current Everyday and Someday Smoking among Adults

The majority of current smokers in Oklahoma report smoking everyday. Throughout the past 13 years, the prevalence of someday smoking has remained consistently lower than everyday smoking. The prevalence of everyday smoking has decreased from 19.9% in 2011 to 11.1% in 2023, with a statistically significant average annual percentage change (-4.0%). During the 13-year period, the prevalence of someday smoking significantly decreased by an average annual change of -3.0%.


**Figure 2.1 Prevalence (%) of everyday and someday smoking among Oklahoma adults
(Oklahoma BRFSS 2011 – 2023)**



Prevalence of Current Everyday Smoking among Adults by Sex

Except for 2021, the proportion of males who were current everyday smokers remained higher than for females. In 2023, the prevalence of current everyday smoking among males was 11.3%, whereas in females it was 10.9%. The average annual decline in prevalence of current everyday smoking among both males (-4.3%) and females (-3.7%) was statistically significant during the past 13 years.


Figure 2.2 Prevalence (%) of everyday smoking among Oklahoma adults by sex (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Someday Smoking among Adults by Sex

The proportion of someday smoking increased for both males and females in 2023. Despite the 2023 increase, the prevalence of someday smoking significantly declined by average annual percent change of 3.1% among females and among males, 3.0%.


Figure 2.3 Prevalence (%) of someday smoking among Oklahoma adults by sex (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Everyday Smoking among Adults by Age

Everyday smoking has declined significantly among young and middle-aged adults, with the most significant decline among those ages 18-24 (APC = -11.5%). During 2021-2023, individuals in the 18–24-year age group and those in the 65 years or older age group were less likely to report everyday smoking compared to other age groups.

Figure 2.4 Three-year rolling prevalence of everyday smoking among Oklahoma adults by age (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Someday Smoking among Adults by Age

The patterns of someday smoking with age were inconsistent. Three-year rolling prevalence of someday smoking was highest among those ages 25-34 years (6.0%) and lowest in the oldest age group (2.8%) in 2021-2023. During the 13-year period, someday smoking had the highest significant decline among the youngest age group (APC = -8.8%), while it significantly increased among the oldest age group (APC = 1.9%).


Figure 2.5 Three-year rolling prevalence of someday smoking among Oklahoma adults by age (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Everyday Smoking among Adults by Race/Ethnicity

The prevalence of everyday smoking was highest among NH-AI/ANs (15.5%) followed by NH-Whites (12.2%) in 2021-2023. The Hispanic population had the lowest prevalence of current everyday smoking of any race/ethnic group (6.2%). The prevalence of everyday smoking significantly declined among all race/ethnic groups by an average annual percent change ranging from -3.2% to -6.0%.


Figure 2.6 Three-year rolling prevalence of everyday smoking among Oklahoma adults by race/ethnicity (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Someday Smoking among Adults by Race/Ethnicity

The prevalence of someday smoking was much lower than everyday smoking for each race/ethnicity group. During 2011-2023, the prevalence of someday smoking significantly declined among all groups, with highest declines in Hispanics (APC -6.0), NH-Blacks (APC -4.5%), AI/AN (-4.4%), and NH-Whites (-3.2%). The prevalence of someday smoking in 2021-2023 increased to 5.2% from 4.1% in Hispanics.


Figure 2.7 Three-year rolling prevalence of someday smoking among Oklahoma adults by race/ethnicity (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Everyday Smoking among Adults by Education

The education gradient observed for all current smokers is also apparent when looking at current everyday smokers. As education level increases, the prevalence of current everyday smoking decreases. The average annual percent decrease in prevalence of everyday smoking was significant across each level of education. During 2021-2023, the prevalence of current everyday smoking was 23.0% among those with less than a high school education and 3.9% among those with a college degree.


Figure 2.8 Three-year rolling prevalence of everyday smoking among Oklahoma adults by education (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Someday Smoking among Adults by Education

An association between education and current someday smoking also exists, although the differences are not as clear as those observed with everyday smoking. In the 2021-23 time-period the prevalence of someday smoking was highest in those with less than HS education (6.4%) and lowest for those with a college education (2.0%). All education groups had a significant decline in the prevalence of someday smoking over the 13-year period with less than high school (-4.5%) and college degree (-4.7%) showing the largest declines.


Figure 2.9 Three-year rolling prevalence of someday smoking among Oklahoma adults by education (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Everyday Smoking among Adults by Income

A consistent gradient occurs with everyday smoking and income level. Everyday smoking is lowest among those in the highest income group, \$50,000 or more (7.0%) and highest in those in the lowest income group, less than \$15,000 (25.8%). The prevalence of everyday smoking among all income groups except \$25,000 to <\$35,000 significantly declined by an average annual change of at least 2% during the 13-year period.


Figure 2.10 Three-year rolling prevalence of everyday smoking among Oklahoma adults by income (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Someday Smoking among Adults by Income

The prevalence of current someday smoking by income varied during the past 13 years. In 2021-2023, the prevalence of current someday smoking was highest (7.4%) for those earning less than \$15,000 and lowest (3.4%) for those earning \$50,000 or more. There was significant average annual change in someday smoking in income groups less than \$15,000 (-3.9%), \$15,000 to < \$25,000 (-1.9%), and the \$25,000 to < \$35,000 group (-4.1%). There was also a non-significant change in the \$50,000 or more income group (-1.8%).


Figure 2.11 Three-year rolling prevalence of someday smoking among Oklahoma adults by income (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Everyday Smoking among Adults by Area of Residence

A difference in the prevalence of current everyday smoking has persisted between rural and urban populations in Oklahoma. In 2023, 13.4% of Oklahomans reporting rural residence currently smoke everyday compared to 8.2% of urban Oklahomans. Both urban and rural Oklahomans had significant declines in the average annual prevalence of everyday smoking over time (4.9% and 3.4%, respectively).


Figure 2.12 Prevalence (%) of everyday smoking among Oklahoma adults by area of residence (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Someday Smoking among Adults by Area of Residence

The prevalence of current someday smoking among adults in Oklahoma did not differ significantly by area of residence during the past 12 years. In 2023, the prevalence of someday smoking among rural residents was almost the same as those living in urban areas of the state (4.7% vs. 4.8%). Both groups changed significantly over the 13-year time period (-2.6 in rural and -3.6 in urban Oklahomans).


Figure 2.13 Prevalence (%) of someday smoking among Oklahoma adults by area of residence (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Everyday Smoking among Adults by Physical Health

During the 2021-23 time period, the prevalence of everyday smoking remained higher among Oklahoma adults reporting poor physical health (18.6%) compared to those without poor physical health (9.6%). The 13-year average change in everyday smoking declined significantly in both groups; -2.1% in those with poor physical health and -4.7% in those not reporting poor physical health.


Figure 2.14 Prevalence (%) of everyday smoking among Oklahoma adults by poor physical health (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Someday Smoking among Adults by Physical Health

Oklahomans reporting poor physical health were also more likely to be current someday smokers compared to those without poor physical health. The prevalence of someday smoking among those with poor physical health rose from 5.9% in 2022 to 8.1% in 2023. The prevalence of someday smoking among Oklahomans without poor physical health also rose, but only slightly, from 3.7% in 2022 to 4.0% in 2023. The average annual percentage of someday smoking over time (2011-2023) changed significantly in those without poor health, decreasing by 3.6%.


Figure 2.15 Prevalence (%) of someday smoking among Oklahoma adults by poor physical health (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Everyday Smoking among Adults by Mental Health

Similar to the patterns with poor physical health, Oklahomans who reported frequent mental distress (FMD) were consistently more likely to report current everyday smoking compared to those without FMD. In 2023, the prevalence of current everyday smoking among individuals with FMD (17.9%) was almost twice that of individuals without FMD (9.6%). Both groups experienced statistically significant declines in the APC (-4.8% for those with FMD and -4.4% for those without FMD).


**Figure 2.16 Prevalence (%) of everyday smoking among Oklahoma adults by Frequent Mental Distress (FMD)
(Oklahoma BRFSS 2011 – 2023)**



Prevalence of Current Someday Smoking among Adults by Mental Health

The prevalence of someday smoking was also higher among Oklahomans reporting Frequent Mental Distress (FMD). In 2023, the percentage of people with FMD who reported smoking on some days rose from 5.1% in 2022 to 8.1%. In comparison, the percentage of people without FMD who reported smoking some days rose only slightly from the 2022 value of 3.7% to the 2023 value of 4.0%. Both groups had statistically significant declines in the APC (-4.6% for those with FMD and -3.0% for those without FMD).


Figure 2.17 Prevalence (%) of someday smoking among Oklahoma adults by Frequent Mental Distress (FMD)
(Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Everyday Smoking among Adults by Health Insurance Coverage

In 2021, there was an increase in the prevalence of current everyday smoking among individuals who reported having no health insurance coverage (23.4%), which sharply declined in 2022 (16.2%) and 2023 (12.7%). That prevalence is now close to that of everyday smoking in those with health insurance coverage (11.0%). The declines in the APC for both groups were statistically significant (-5.9% among uninsured and -2.9% among those with insurance).


Figure 2.18 Prevalence (%) of everyday smoking among Oklahoma adults by health insurance coverage (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Someday Smoking among Adults by Health Insurance Coverage

From 2020 through 2021 individuals with no health insurance coverage were unexpectedly less likely to report current someday smoking compared to those with coverage. However, during 2022 a reverse of that trend occurred and the prevalence of someday smoking was 4.8% in those without insurance and 4.0% for those with insurance. That trend continued in 2023 when the prevalence of someday smoking for those without insurance jumped to 9.1% compared to only a small increase in those with insurance (4.3%). The 13-year period APC for someday smoking significantly changed for those without health insurance by an average annual change of -4.9% compared to -2.1% in those with insurance.


Figure 2.19 Prevalence (%) of someday smoking among Oklahoma adults by health insurance coverage (Oklahoma BRFSS 2011 – 2023)



Prevalence of Current Everyday Smoking among Adults by Disability

The prevalence of current everyday smoking was consistently higher among those who reported one or more disabilities compared to those with no disability; these differences were significant each year. In 2023, the prevalence of current everyday smoking was 16.5% among those with disability and 7.6% among those without disability. There was a significant average annual change of -3.1% in everyday smoking prevalence among those with one or more disabilities, and -4.0% in those no disabilities.


Figure 2.20 Prevalence (%) of everyday smoking among Oklahoma adults by disability
(Oklahoma BRFSS 2016 – 2023)



Prevalence of Current Someday Smoking among Adults by Disability

Adults in Oklahoma with one or more disabilities were more likely to report current someday smoking compared to those with no disability, although the differences were not always significant. The prevalence of current someday smoking was 6.1% among those with one or more disabilities and 3.7% among those with no disability in 2023. The prevalence of someday smoking among those with one or more disabilities demonstrated a significant annual decline (-4.6%).

Figure 2.21 Prevalence (%) of someday smoking among Oklahoma adults by disability
(Oklahoma BRFSS 2016 – 2023)




RESULTS: CIGARETTE SMOKING

FORMER CIGARETTE SMOKERS

Prevalence of Former Smoking among Adults

Former smokers are defined as individuals who have smoked at least 100 cigarettes in their entire lifetime but do not currently smoke. The prevalence of former smoking among adults in Oklahoma has remained around 25% from 2011 to 2021, however, it increased to 27.2% in 2022 and then decreased to 26.1% in 2023.


**Figure 3.1 Prevalence (%) of former smoking among Oklahoma adults
(Oklahoma BRFSS 2011–2023)**



Prevalence of Former Smoking among Adults by Sex

The prevalence of former smoking was consistently higher among males from 2011 to 2023. In 2023, 29.9% of males were former smokers compared to 22.5% of females. Females had a significant increase in the average annual prevalence of former smoking over time by 1.0%.


Figure 3.2 Prevalence (%) of former smoking among Oklahoma adults by sex (Oklahoma BRFSS 2011 – 2023)



Prevalence of Former Smoking among Adults by Age

The prevalence of former smoking in Oklahoma increases with increasing age. More than one third (38.0%) of Oklahomans aged 65 or older report being former smokers compared to 8.3% for those aged 18 to 24 years. The prevalence of former smoking among those aged 35-44 years (APC = 3.2%) significantly increased. However, among those 55 or older, the prevalence of former smoking significantly decreased (APC = -1.0%) over the 13-year period.


Figure 3.3 Three-year rolling prevalence of former smoking among Oklahoma adults by age
(Oklahoma BRFSS 2011 – 2023)



Prevalence of Former Smoking among Adults by Race/Ethnicity

The prevalence of former smoking varies by race/ethnicity. In 2023, NH-Whites had the highest proportion of former smokers (28.8%). Hispanics had the lowest proportion of former smokers (16.8%). NH-AI/AN and NH-Other had significant annual increases in the prevalence of former smoking over time (APC 1.8% and 2.5%, respectively).


Figure 3.4 Three-year rolling prevalence of former smoking among Oklahoma adults by race/ethnicity (Oklahoma BRFSS 2011 – 2023)



Prevalence of Former Smoking among Adults by Education

The proportion of former smokers in Oklahoma did not differ significantly by education. During 2021-2023, the proportion of former smokers remained between 22.6% (college graduates) and 28.7% (some college). Over time, there was significant average annual percent decline (-0.7%) in the proportion of former smokers among college graduates, whereas the APC among those with high school level education (+0.9%), and those with some college education (+1.1%) increased.


Figure 3.5 Three-year rolling prevalence of former smoking among Oklahoma adults by education (Oklahoma BRFSS 2011 – 2023)



Prevalence of Former Smoking among Adults by Income

The proportion of former smokers varies by income and over time. In 2021-2023, those earning \$50,000 or more and those earning between \$25,000 to less than \$35,000 had the highest prevalence of former smoking (27.9%) followed by those earning \$15,000 to less than \$25,000 (26.4%) and those earning \$35,000 to less than \$50,000 (26.2%). Over time, there was significant average annual percent increase of 1.7% in the proportion of former smokers among those earning less than \$15,000, and a 0.7% increase in those earning \$25,000 to < \$35,000 whereas, among those earning \$35,000 to < \$50,000 the APC decreased by 0.7%.


Figure 3.6 Three-year rolling prevalence of former smoking among Oklahoma adults by income (Oklahoma BRFSS 2011 – 2023)



Prevalence of Former Smoking among Adults by Area of Residence

In 2023, the proportion of former smokers among adults in Oklahoma was higher for those living in rural areas compared to those in urban areas of the state (27.1% vs. 24.8%). There were no significant changes in the proportion of former smokers by area of residence during the past 13 years.


Figure 3.7 Prevalence (%) of former smoking among Oklahoma adults by area of residence (Oklahoma BRFSS 2011 – 2023)



Prevalence of Former Smoking among Adults with Poor Physical Health

Oklahomans who report having poor physical health had a higher prevalence of former smoking compared to those without poor physical health. About one third (32.6%) of those with poor physical health were former smokers compared to 24.7% of those without poor health. Over the past 13 years, the prevalence of former smoking significantly increased by an average annual change of 0.6% among those without poor physical health.


Figure 3.8 Prevalence (%) of former smoking among Oklahoma adults by physical health status (Oklahoma BRFSS 2011 – 2023)



Prevalence of Former Smoking among Adults with Poor Mental Health Status

Although there was a difference in the proportion of former smokers by physical health status, there were no significant differences in the proportion of former smokers by mental health status. During the past 13 years, the proportion of former smokers has remained similar for both those with and without FMD, however in 2023, the gap widened minimally and those with FMD had an increase in prevalence of former smoking (29.3%), while those without FMD had a decrease in the prevalence of former smoking (25.5%). Those with frequent mental distress had a significant increase in the prevalence of former smoking by average annual percent change of 2.1%.


Figure 3.9 Prevalence (%) of former smoking among Oklahoma who reported Frequent Mental Distress (FMD) (Oklahoma BRFSS 2011 – 2023)



Prevalence of Former Smoking among Adults by Health Insurance Coverage

The prevalence of adult former smoking in Oklahoma was higher for those who are insured compared to those who are uninsured, and these differences were consistent over time. About one quarter (27.5%) of the insured were former smokers, compared to 15.8% of the uninsured. Neither change in APC was significant over the past 13 years.


Figure 3.10 Prevalence (%) of former smoking among Oklahoma adults by health insurance coverage (Oklahoma BRFSS 2011 – 2023)



Prevalence of Former Smoking among Adults by Disability

In 2023 the prevalence of former smoking was significantly higher for those who reported one or more disability (30.0%) compared to those not reporting a disability (23.4%). These differences were consistent over time. There were no significant changes in the prevalence of former smoking during the past seven years for those with or without a disability.

Figure 3.11 Prevalence (%) of former smoking among Oklahoma adults by disability (Oklahoma BRFSS 2016 – 2023)




RESULTS: SMOKELESS TOBACCO USE

Prevalence of Smokeless Tobacco Use among Adults

In addition to a high prevalence of cigarette smoking, Oklahoma also faces challenges with a high prevalence of smokeless tobacco use. Among all adults, the prevalence of smokeless tobacco was 6.3% in 2023, up from 5.7% in 2022. There was not a significant change in prevalence over the past 13 years.


**Figure 4.1 Prevalence (%) of smokeless tobacco use among Oklahoma adults
(Oklahoma BRFSS 2011 – 2023)**



Prevalence of Smokeless Tobacco Use among Adults by Sex

The majority of individuals who use smokeless tobacco are males. In Oklahoma, the prevalence of smokeless tobacco use among males decreased from 13.9% in 2019 to 10.5% in 2022, with a slight increase to 10.8% in 2023. These changes are not statistically significant. Among females, the prevalence of smokeless tobacco use rose to 2.0% in 2023. Over the past 13 years, the annual percentage change (APC) for smokeless tobacco use among females has shown a significant increase of 6.2%.

Figure 4.2 Prevalence (%) of smokeless tobacco use among Oklahoma adults by sex (Oklahoma BRFSS 2011 – 2023)



Prevalence of Smokeless Tobacco Use among Male Adults by Age

Among males, the prevalence of smokeless tobacco (ST) use varies by age. The prevalence of ST use fluctuated throughout the past 13 years. Males aged 55-64 years had a 6.5% annual percent increase in ST use; however, there was a significant decline in the prevalence of ST use among males 25-34 and 35-44 years old over the past 13 years (APC -1.0% and -2.3% respectively). There was a sharp significant decline in prevalence among males 18-24 years old between 2011-2023 (APC -6.8%).


Figure 4.3 Three-year rolling prevalence of smokeless tobacco use among Oklahoma male adults by age (Oklahoma BRFSS 2011 – 2023)



Prevalence of Smokeless Tobacco Use among Male Adults by Race/Ethnicity

The prevalence of smokeless tobacco use is consistently higher among NH-American Indian males compared to all other race/ethnicity groups. In 2021-2023, 14.7% of NH-American Indian males used smokeless tobacco. Only 3.7% of NH-Blacks used smokeless tobacco, a difference that was statistically significant. The prevalence of smokeless tobacco use significantly decreased among NH-Other by average annual percent change of 3.8%.


Figure 4.4 Three-year rolling prevalence of smokeless tobacco use among Oklahoma male adults by race/ethnicity (Oklahoma BRFSS 2011 – 2023)



Prevalence of Smokeless Tobacco Use among Male Adults by Education

The prevalence of smokeless tobacco use among males was consistently lower for those with a college degree compared to those with less education. In 2021-23, the prevalence of smokeless tobacco was highest among those with high school education followed by those with some college education (15.0% and 11.1%, respectively). The prevalence of smokeless tobacco use was lowest in those with a college education, followed by those with less than a high school education (6.8% and 8.4%, respectively). Over the last 13 years, the APC was not statistically significant for any of the education categories.


Figure 4.5 Three-year rolling prevalence of smokeless tobacco use among Oklahoma male adults by education (Oklahoma BRFSS 2011 – 2023)



Prevalence of Smokeless Tobacco Use among Male Adults by Income

The prevalence of smokeless tobacco use among males by income fluctuated throughout the past 13 years, and there were no consistent differences between income groups. There was only a statistically significant in smokeless tobacco use for the less than \$15,000 income group (APC -3.9%).


Figure 4.6 Three-year rolling prevalence of smokeless tobacco use among Oklahoma male adults by income (Oklahoma BRFSS 2011 – 2023)



Prevalence of Smokeless Tobacco Use among Male Adults by Area of Residence

Among Oklahoma males, the prevalence of smokeless tobacco use among those in rural areas of residence was approximately twice as high as those in urban areas. In 2023, the prevalence of smokeless tobacco use among males in urban areas was 6.9%; however, 13.8% of males in rural areas reported using smokeless tobacco.


Figure 4.7 Prevalence (%) of smokeless tobacco use among Oklahoma male adults by area of residence (Oklahoma BRFSS 2011 – 2023)



Prevalence of Smokeless Tobacco Use among Male Adults with Poor Physical Health Status

Unlike cigarette smoking, the prevalence of smokeless tobacco use in Oklahoman did not differ significantly by physical health status among men. There were no significant changes in smokeless tobacco use prevalence across the past 13 years for either group.


Figure 4.8 Prevalence (%) of smokeless tobacco use among Oklahoma male adults by poor physical health status (Oklahoma BRFSS 2011 – 2023)



Prevalence of Smokeless Tobacco Use among Male Adults by Mental Health Status

The prevalence of smokeless tobacco use in 2022-2023 rose from 9.9% to 12.5% for Oklahoma males reporting Frequent Mental Distress (FMD) compared to those who had no FMD (10.6% to 10.4% in 2022-2023). There was a significant decrease in smokeless tobacco use during the past 13 years in the group of Oklahoma males reporting frequent mental distress (-3.6%).


Figure 4.9 Prevalence (%) of smokeless tobacco use among Oklahoma male adults by Frequent Mental Distress (FMD) (Oklahoma BRFSS 2011 – 2023)



Prevalence of Smokeless Tobacco Use among Male Adults by Health Insurance Coverage

There were no significant differences in the prevalence of smokeless tobacco use among Oklahoma males with or without health insurance coverage. The prevalence fluctuated for those uninsured, with an overall decrease in smokeless tobacco use among males across the 13 years (APC -2.7%).


Figure 4.10 Prevalence (%) of smokeless tobacco use among Oklahoma male adults by health insurance coverage (Oklahoma BRFSS 2011 – 2023)



Prevalence of Smokeless Tobacco Use among Male Adults by Disability

The prevalence of smokeless tobacco use among Oklahoma men did not differ significantly by disability status. In 2023, the prevalence of smokeless tobacco use was higher among men with a disability (10.8%) than men without a disability (10.6%). There were no significant changes in smokeless tobacco use during the past 12 years in either group.

Figure 4.11 Prevalence (%) of smokeless tobacco use among Oklahoma male adults by disability (Oklahoma BRFSS 2016 – 2023)



RESULTS: USE OF ELECTRONIC NICOTINE (E-CIGARETTE OR EC) PRODUCTS*

*Based on state-added question “Have you used an electronic cigarette or other electronic vaping device in the last 30 days?”

Prevalence of EC use among Adults

In 2023, 8.5% of Oklahomans reported using an EC in the last 30 days compared to 9.3% in 2022. Prevalence of EC use has remained consistent since 2020. There was not a significant change during the past seven years in the prevalence of EC use.


Figure 5.1 Prevalence of EC use* among Oklahoma adults (Oklahoma BRFSS 2017 – 2023)



Prevalence of EC use among Adults by Sex

Similar to prior years, prevalence of EC use is higher among males (10.1%) compared to females (6.9%) in 2023. Over the past seven years, there was no significant change in the proportion of EC use among males and females.


Figure 5.2 Prevalence of EC use* among adults by sex (Oklahoma BRFSS 2017 – 2023)



Prevalence of EC use among Adults by Age

EC use declined with increasing age; 18-24 years old had the highest three-year rolling percentage (26.7%) whereas 65 years or older had the lowest (1.2%) during 2021-23. In younger age groups, less than 35 years old, EC use significantly increased from 2017-19 to 2020-23 (6.8% in those 18-24 years and 5.7% in those 25-34 years). EC use significantly decreased from 2017-19 to 2020-23 in those 45-54 years (-4.7%), those 55-64 years (-14.0%), and in those 65+ years (-17.4%).


Figure 5.3 Three-year rolling prevalence of EC use* among Oklahoma adults by age
(Oklahoma BRFSS 2017 – 2023)



Prevalence of EC use among Adults by Race/Ethnicity

In 2020-23, NHA/ANs reported the highest levels of EC use at 11.8% followed by NH Other at 11.6% (which dropped from the 2019-22 value of 14.4%). Over time, the prevalence of EC use in the last 30 days increased among Hispanics by an average annual percent change of 3.0%.


Figure 5.4 Three-year rolling prevalence of EC use* among Oklahoma adults by race/ethnicity (Oklahoma BRFSS 2017 – 2023)



Prevalence of EC use among Adults by Education

College graduates had the lowest level of EC in 2021-23 (3.7%) and those with high school education had the highest level of EC use (11.6%). There was not a significant change during the past seven years in the prevalence of EC use in any of the education groups.


Figure 5.5 Three-year rolling prevalence of EC use* among Oklahoma adults by education (Oklahoma BRFSS 2017 – 2023)



Prevalence of EC use among Adults by Income

There were no significant differences in EC use by income level. In 2021-2023, the highest prevalence of EC use (12.0%) was reported among those earning less than \$15,000 per year, whereas it was the lowest among those earning \$50,000 or more (7.0%). There was a significant change during the past seven years in the prevalence of EC use among those earning \$25,000 to <\$35,000 (-6.9%).


Figure 5.6 Three-year rolling prevalence of EC use* among Oklahoma adults by income
(Oklahoma BRFSS 2017 – 2023)



Prevalence of EC use among Adults by Area of Residence

The prevalence of EC use among those reporting urban residence did not significantly change from the 2022 value of 9.2% to the 2023 value of 8.4%. Similarly, it did not significantly change in those with rural residence, moving from 9.5% in 2022 to 8.6% in 2023. There were no significant changes in EC use during the past seven years for either group.


**Figure 5.7 Prevalence of EC use* among adults by area of residence
(Oklahoma BRFSS 2017 – 2023)**



Prevalence of EC use among Adults with Poor Physical Health

In 2023, the prevalence of EC use among those without poor physical health was 7.9% and those with poor health it was 11.2%. There was no significant change the prevalence of EC use in both groups over the past seven years.


Figure 5.8 Prevalence of EC use* among adults by Frequent Physical Distress (FPD)
(Oklahoma BRFSS 2017 – 2023)



Prevalence of EC use among Adults with Poor Mental Health Status

Oklahomans reporting Frequent Mental Distress (FMD) have consistently higher prevalence of EC use than those without FMD. In 2023, 17.1% of those with FMD reported EC use, whereas 6.9% of those without FMD reported EC use. There was no significant change the prevalence of EC use in both groups over the past seven years.


Figure 5.8 Prevalence of EC use* among adults by Frequent Mental Distress (FMD)
(Oklahoma BRFSS 2017 – 2023)



Prevalence of EC use among adults by Insurance Coverage

The prevalence of EC use was higher among uninsured compared to those with health insurance (12.2% vs. 8.1%). There were no statistically significant changes in EC use during the past seven years for either group.


**Figure 5.10 Prevalence of EC use* among adults by insurance coverage
(Oklahoma BRFSS 2017 – 2023)**



Prevalence of EC use among adults by Disability

Those who reported one or more disabilities had consistently higher prevalence of EC use than those with no disability. In 2023, 10.9% of those with one or more disabilities and 6.8% of those without disability reported EC use. There was no significant change the prevalence of EC use in both groups over the past seven years.

Figure 5.11 Prevalence of EC use* among adults by disability (Oklahoma BRFSS 2017 – 2022)




RESULTS: QUIT ATTEMPTS

Proportion of Adult Smokers with a Quit Attempt in the Last Year

Many adult current smokers want to quit. In Oklahoma, the proportion of current smokers who report a quit attempt in the last year declined between 2019-2022. In 2023 this value rebounded, with 51.5% of adult current smokers reporting a quit attempt in the last year. From 2011 to 2023 the percentage who reported a quit attempt in the last year was between 47.0% and 61.5%, and the APC declined significantly (-1.6%).


Figure 6.1 Percent of smokers with a quit attempt in the last year (Oklahoma BRFSS 2011 – 2023)



Proportion of Adult Smokers with a Quit Attempt in the Last Year by Sex

The proportion of adult smokers who reported a quit attempt in the was higher for females compared to males for all year except 2020. Between 51.7% and 64.6% of females reported a past quit attempt each of the past 13 years. Among males, the proportion reporting a quit attempt ranged from 43.0% to 58.8%. There was a significant average annual decline in the proportion of female smokers (-1.7%) and male smokers (-1.5%) reporting a quit attempt in the last year.


Figure 6.2 Percent of smokers with a quit attempt in the last year by gender (Oklahoma BRFSS 2011 – 2023)



Proportion of Adult Smokers with a Quit Attempt in the Last Year by Age

The proportion of smokers with a quit attempt in the last year tends to decrease with increasing age. The proportion of Oklahoma adult smokers aged 18-24 reporting a quit attempt ranged from 60.6% to 77.3% from 2011 to 2023, while the proportion of smokers aged 55 and older remained below 50.2% each year. There was significant average annual percent decline of 0.9% or more in the proportion of smokers reporting a quit attempt during the past 13 years for those aged 35 years or older and -1.6% in those 18-24 years.


Figure 6.3 Three-year rolling prevalence of smokers with a quit attempt in the last year by age (Oklahoma BRFSS 2011 – 2023)



Proportion of Adult Smokers with a Quit Attempt in the Last Year by Race/Ethnicity

The proportion of current smokers reporting a quit attempt during 2021-2023 varies somewhat by race/ethnicity, with a high of 68.3% among NH-Black smokers and a low of 45.8% among NH-other smokers. Over time, the average annual percent change in past quit attempts was statistically significant for NH-Whites (-1.9%), NH-AI/AN (-1.3%), and NH-Other (-3.6%) smokers.


Figure 6.4 Three-year rolling prevalence of smokers with a quit attempt in the last year by race/ethnicity (Oklahoma BRFSS 2011 – 2023)



Proportion of Adult Smokers with a Quit Attempt in the Last Year by Education

When examining potential differences in quit attempts by education level, there were no significant differences between groups in the proportion of current smokers reporting a quit attempt in the last year. Smokers with some college education had the highest rate of quit attempts in 2021-2023 (52.4%), and smokers with a college education had the lowest rate of quit attempts (45.1%). There was a statistically significant decrease in past quit attempts for all income groups over the past 13 years.


Figure 6.5 Three-year rolling prevalence of smokers with a quit attempt in the last year by education (Oklahoma BRFSS 2011 – 2023)



Proportion of Adult Smokers with a Quit Attempt in the Last Year by Income

Current smokers in the \$35,000 to less than \$50,000 category had the highest percentage of smokers reporting a quit attempt in the 2021-2023-year period (57.2%). Smokers in the \$50,000 or more income category had the lowest percentage reporting a quit attempt (43.3%) in the same time period. The proportion of past quit attempts during the past 13 years changed significantly in every category from -0.2% in the \$35,000 to less than \$50,000 category to -2.6% in the less than \$15,000 category.


Figure 6.6 Three-year rolling prevalence of smokers with a quit attempt in the last year by income (Oklahoma BRFSS 2011 – 2023)



Proportion of Adult Smokers with a Quit Attempt in the Last Year by Area of Residence

Oklahoma smokers living in urban areas were slightly more likely to report a quit attempt in the last year compared to those in rural areas. In 2023, 52.9% of urban residents reported a quit attempt in the last year compared to 50.6% of smokers in rural areas. There was a significant change of in the proportion of smokers in both groups reporting a quit attempt over time (-2.1 for smokers in rural areas and -0.9 for smokers in urban areas).


Figure 6.7 Percent of smokers with a quit attempt in the last year by area of residence (Oklahoma BRFSS 2011 – 2023)



Proportion of Adult Smokers with a Quit Attempt in the Last Year by Physical Health

There were no significant differences in the proportion of smokers with a quit attempt in the last year by physical health status. In 2023, 65.0% of Oklahoma adult smokers with poor physical health and 48.4% of those without poor health reported a past quit attempt. There was significant change in past quit attempts (-1.9%) in those without poor physical health).

Figure 6.8 Percent of smokers with a quit attempt in the last year by poor physical health (Oklahoma BRFSS 2011 – 2023)




Poor physical health was defined as having 14 or more days of poor physical health in the past 30 days

Proportion of Adult Smokers with a Quit Attempt in the Last Year by Mental Health

Similar to physical health, there were no significant differences in the proportion of past quit attempts among smokers by mental health status. In 2023, 61.8% of adult smokers with frequent mental distress (FMD) reported a quit attempt in the past year, compared to 47.2% among those without FMD in 2023. Since 2011, there was significant decline in the proportion of smokers reporting quit attempts in the last year among those without FMD (-1.8%).

Figure 6.9 Percent of smokers with a quit attempt in the last year by Frequent Mental Distress (FMD)
(Oklahoma BRFSS 2011 – 2023)




14 or more days of poor mental health in the past 30 days is referred to as Frequent Mental Distress (FMD)

Proportion of Adult Smokers with a Quit Attempt in the Last Year by Health Insurance Coverage

The proportion of adult smokers with a quit attempt in the last year was not significantly different for those with health insurance coverage compared to those without health insurance coverage. Among those without insurance, the proportion of smokers who reported a quit attempt increased from the 2022 value of 40.3% to the 2023 value of 66.5%. The proportion of smokers with insurance who made a quit attempt only increased from 48.2% to 49.3%. Since 2011, there was only a significant change in the proportion of smokers reporting quit attempts in the last year among those with insurance (-1.7%).


Figure 6.10 Percent of smokers with a quit attempt in the last year by health insurance coverage (Oklahoma BRFSS 2011 – 2023)



Proportion of Adult Smokers with a Quit Attempt in the Last Year by Disability

Oklahoma smokers with no disability were less likely to report a quit attempt in 2023 (45.8%) compared to those with a disability (55.6%). During the past 13 years, there was only a significant change (-3.6%) in the proportion of smokers with no disabilities reporting a quit attempt.

Figure 6.11 Percent of smokers with a quit attempt in the last year by disability (Oklahoma BRFSS 2016 – 2023)




RESULTS: SHS EXPOSURE AT HOME*

*SHS questions were not asked on the 2022 BRFSS.

Proportion of Adults with a Home Smoking Ban

The proportion of Oklahoma adults reporting a ban on smoking in the home had a statistically significant average annual percent increase of 1.1% over the past 13 years. In 2023, 87.2% of Oklahoma adults reported a smoking ban at home.

Figure 7.1 Proportion of Oklahoma adults reporting a smoking ban at home (Oklahoma BRFSS 2011 – 2023)




For APC calculation, missing values for 2022 were imputed by taking the mean of the 2021 and 2023 values.

Proportion of Adults with a Home Smoking Ban by Sex

The proportion of males and females reporting a ban on smoking in the home increased similarly over the past 13 years, with an average annual percent increase of 1.3% and 0.9%, respectively. In 2023, 86.2% of males and 88.3% of females reported a smoking ban at home.

Figure 7.2 Proportion of Oklahoma adults reporting a smoking ban at home by sex (Oklahoma BRFSS 2011 – 2023)



For APC calculation, missing values for 2022 were imputed by taking the mean of the 2021 and 2023 values.

Proportion of Adults with a Home Smoking Ban by Age

The proportion of adults with a home smoking ban did not vary significantly by age but did increase over time in all age groups. The average annual percent change increased significantly by about 1% within each age group during the 13-year period.


Figure 7.3 Three-year rolling prevalence of Oklahoma adults reporting a smoking ban at home by age (Oklahoma BRFSS 2011 – 2023)



Proportion of Adults with a Home Smoking Ban by Race/Ethnicity

The proportion of adults with home smoking bans did differ significantly by race/ethnicity at times. NH-AI/AN had the highest average annual percent increase in proportion of adults reporting home smoking ban (APC 2%). Except for those reporting Hispanic ethnicity, all other groups had a significant increase in home smoking bans over the past 13 years.


Figure 7.4 Three-year rolling prevalence of Oklahoma adults reporting a smoking ban at home by race/ethnicity (Oklahoma BRFSS 2011 – 2023)



Proportion of Adults with a Home Smoking Ban by Education

There was a positive association between education and home smoking bans. Individuals with less than a high school diploma were less likely to report a home smoking ban (82.2%), while those with a college degree were most likely (93.0%) in 2021-2023. Over time, there were significant increases in the proportion of individuals with a home smoking ban for all education groups (ranging from 2.2% for less than high school to 0.2% for those with a college degree).


Figure 7.5 Three-year rolling prevalence of Oklahoma adults reporting a smoking ban at home by education (Oklahoma BRFSS 2011 – 2023)



Proportion of Adults with a Home Smoking Ban by Income

The proportion of adults with a home smoking ban was typically lower among populations with lower income; in 2023, 76.0% of adults making <\$15,000 had a home smoking ban compared to was 92.7% of adults making \$50,000 or more. However, home smoking bans among adults in income groups <\$15,000 and \$15,000 to <\$25,000 significantly increased by about 2% during the past 13 years.


Figure 7.6 Three-year rolling prevalence of Oklahoma adults reporting a smoking ban at home by income (Oklahoma BRFSS 2011 – 2023)



Proportion of Adults with a Home Smoking Ban by Area of Residence

Adults in Oklahoma living in urban areas were more likely to report a home smoking ban in 2023 (88.0%) compared to rural areas (86.6%). The proportion of adults with a home smoking ban increased significantly by an average of 0.9% each year for urban residents and 1.3% for rural Oklahomans.

Figure 7.7 Proportion of Oklahoma adults reporting a smoking ban at home by area of residence (Oklahoma BRFSS 2011 – 2023)




For APC calculation, missing values for 2022 were imputed by taking the mean of the 2021 and 2023 values.

Proportion of Adults with a Home Smoking Ban by Physical Health

Adults reporting poor physical health were less likely to report a home smoking policy compared to those without poor health. In 2023, 88.6% of adults without poor physical health reported a home smoking ban compared to 79.2% of those in poor physical health. The proportion of adults reporting a home smoking policy significantly increased by an average of 1.6% per year among those in poor health and by 1.0% among those without poor health.

Figure 7.8 Proportion of Oklahoma adults reporting a smoking ban at home by poor physical health (Oklahoma BRFSS 2011 – 2023)




For APC calculation, missing values for 2022 were imputed by taking the mean of the 2021 and 2023 values.

Proportion of Adults with a Home Smoking Ban by Mental Health

In 2023, Oklahomans with poor mental health, or Frequent Mental Distress (FMD), were less likely to report a home smoking ban (77.1%) compared to those without FMD (89.7%). The proportion of home smoking bans significantly increased an average of 1.0% annually for those without FMD and increased an average of 2.7% annually among those with FMD.

Figure 7.9 Proportion of Oklahoma adults reporting a smoking ban at home by Frequent Mental Distress (FMD) (Oklahoma BRFSS 2011 – 2023)




For APC calculation, missing values for 2022 were imputed by taking the mean of the 2021 and 2023 values.

Proportion of Adults with a Home Smoking Ban by Health Insurance Coverage

Oklahoma adults with health insurance coverage were more likely to report having a home smoking ban compared to those who were not insured. The proportion of adults with insurance having a home smoking ban increased an average of 1.0% annually and by 2023, 87.6% had policies. Among those without insurance, 84.3% had home smoking bans in 2021 and the proportion significantly increased an average of 1.7% annually during the past 13 years.

Figure 7.10 Proportion of Oklahoma adults reporting a smoking ban at home by health insurance coverage (Oklahoma BRFSS 2011 – 2023)




For APC calculation, missing values for 2022 were imputed by taking the mean of the 2021 and 2023 values.

Proportion of Adults with a Home Smoking Ban by Disability

Those with one or more disabilities consistently reported a lower prevalence of home smoking bans compared to those with no disability. In 2023, 82.0% of those with at least one disability and 90.8% of those without a disability had a home smoking ban. There was a significant increase of 1.4% in the average annual percentage of individuals with a home smoking ban among those with one or more disabilities.

Figure 7.11 Proportion of Oklahoma adults reporting a smoking ban at home by disability (Oklahoma BRFSS 2016 – 2023)



For APC calculation, missing values for 2022 were imputed by taking the mean of the 2021 and 2023 values.

Appendix A

Tables

Tobacco Use Patterns and Burden in Oklahoma 2011-2023

Section I. Cigarette smoking

Table 1. Current cigarette smoking status by year, Oklahoma BRFSS 2011 – 2023

Smoking Status	2011		2012		2013		2014		2015	
	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Current smoker	1857	26.1 (24.7, 27.5)	1636	23.3 (22.0, 24.6)	1567	23.7 (22.4, 25.0)	1427	21.0 (19.8, 22.3)	1148	22.2 (20.6, 23.7)
Former smoker	2482	24.7 (23.5, 25.9)	2201	24.4 (23.2, 25.5)	2327	24.5 (23.4, 25.7)	2404	25.9 (24.8, 27.1)	1992	24.7 (23.3, 26.1)
Never smoker	4169	49.2 (47.6, 50.7)	4111	52.4 (50.9, 53.8)	4240	51.8 (50.3, 53.2)	4366	53.0 (51.6, 54.4)	3613	53.4 (51.4, 54.9)

Smoking Status	2016		2017		2018		2019		2020	
	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Current smoker	1010	19.6 (18.2, 21.0)	1008	20.2 (18.8, 21.5)	839	19.7 (18.2, 21.2)	969	18.9 (17.6, 20.2)	798	19.1 (17.6, 20.6)
Former smoker	1914	24.8 (23.5, 26.2)	1856	24.5 (23.2, 25.8)	1439	24.8 (23.3, 26.3)	1699	25.3 (24.0, 26.6)	1307	25.2 (23.7, 26.7)
Never smoker	3766	55.57 (53.9, 57.2)	3507	55.3 (53.7, 56.9)	2782	55.5 (53.7, 57.3)	3368	55.8 (54.3, 57.4)	2617	55.7 (53.9, 57.5)

Smoking Status	2021		2022		2023	
	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Current smoker	767	16.9 (15.6, 18.3)	778	15.6 (14.4, 16.8)	900	15.8 (14.7, 16.9)
Former smoker	1408	25.7 (24.2, 27.2)	1555	27.2 (25.8, 28.6)	1820	26.1 (24.8, 27.4)
Never smoker	2918	57.4 (55.6, 59.1)	3037	57.2 (55.6, 58.8)	3637	58.1 (56.6, 59.6)

Table 2. Gender and current smoking status by year, Oklahoma BRFSS 2011 – 2023

Gender	Smoking Status	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Male	Current smoker	753	28.1 (25.8, 30.4)	685	24.4 (22.4, 26.4)	689	25.8 (23.8, 27.8)	648	23.0 (21.1, 25.0)	505	24.0 (21.4, 26.5)
	Former smoker	1166	28.8 (26.7, 30.8)	1150	29.6 (27.7, 31.5)	1114	27.6 (25.8, 29.5)	1158	29.6 (27.7, 31.5)	960	27.6 (25.4, 29.8)
	Never smoker	1244	43.2 (40.7, 45.6)	1371	46.0 (43.7, 48.3)	1428	46.6 (44.4, 48.8)	1530	47.4 (45.2, 49.5)	1209	48.5 (45.7, 51.2)
Female	Current smoker	1104	24.3 (22.6, 26.0)	951	22.2 (20.6, 23.8)	878	21.7 (20.0, 23.4)	779	19.1 (17.5, 20.7)	643	20.4 (18.5, 22.3)
	Former smoker	1316	20.9 (19.5, 22.2)	1051	19.4 (18.0, 20.7)	1213	21.6 (20.1, 23.0)	1246	22.4 (21.0, 23.9)	1032	21.9 (20.1, 23.7)
	Never smoker	2925	54.9 (53.0, 56.7)	2740	58.4 (56.6, 60.3)	2812	56.7 (54.9, 58.6)	2836	58.5 (56.6, 60.3)	2404	57.7 (55.4, 59.9)

Gender	Smoking Status	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Male	Current smoker	445	21.4 (19.1, 23.7)	447	21.2 (19.1, 23.3)	407	22.2 (19.9, 24.5)	465	21.1 (19.1, 23.1)	368	20.6 (18.3, 23.0)
	Former smoker	944	28.2 (26.1, 30.3)	916	27.9 (25.8, 29.9)	718	28.5 (26.2, 30.8)	883	29.1 (27.1, 31.2)	644	28.3 (25.9, 30.6)
	Never smoker	1316	50.4 (47.8, 52.9)	1232	50.9 (48.4, 53.4)	1001	49.3 (46.6, 52.0)	1260	49.8 (47.4, 52.2)	1013	51.1 (48.4, 53.8)
Female	Current smoker	565	17.9 (16.1, 19.6)	561	19.2 (17.4, 21.0)	432	17.2 (15.4, 19.1)	504	16.8 (15.1, 18.4)	430	17.6 (15.8, 19.4)
	Former smoker	970	21.6 (20.0, 23.2)	940	21.4 (19.7, 23.0)	720	21.3 (19.5, 23.1)	816	21.6 (19.9, 23.3)	663	22.3 (20.4, 24.2)
	Never smoker	2450	60.5 (58.4, 62.6)	2273	59.4	1781	61.5	2108	61.6 (59.5, 63.7)	1604	60.1 (57.7, 62.4)

(57.3, 61.6)(59.3, 63.8)

Gender	Smoking Status	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Male	Current smoker	340	17.1 (15.0, 19.1)	375	16.6 (14.8, 18.5)	403	16.4 (14.7, 18.1)
	Former smoker	718	29.0 (26.6, 31.4)	780	29.7 (27.6, 31.8)	948	29.9 (27.9, 31.9)
	Never smoker	1183	53.9 (51.3, 56.6)	1257	53.7 (51.3, 56.0)	1483	53.7 (51.5, 56.0)
Female	Current smoker	427	16.8 (15.0, 18.6)	403	14.6 (13.0, 16.2)	497	15.3 (13.8, 16.8)
	Former smoker	690	22.6 (20.6, 24.6)	775	24.7 (22.9, 26.5)	872	22.5 (20.9, 24.1)
	Never smoker	1735	60.6 (58.3, 62.9)	1780	60.7 (58.6, 62.8)	2154	62.3 (60.3, 64.2)

Table 3. Three-year rolling percentage of current cigarette smoking among Oklahoma adults by age, Oklahoma BRFSS 2011 – 2023

Age Group	Smoking Status	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
18-24	Current smoker	296	26.8 (23.7, 29.8)	294	26.1 (23.2, 29.0)	236	24.2 (21.1, 27.4)	195	22.5 (19.3, 25.7)	162	21.4 (18.1, 24.7)
	Former smoker	107	8.0 (6.4, 9.6)	101	7.6 (6.0, 9.2)	87	8.0 (6.1, 9.9)	70	7.0 (5.1, 8.8)	61	6.5 (4.6, 8.3)
	Never smoker	735	65.2 (62.0, 68.5)	795	66.3 (63.2, 69.4)	727	67.7 (64.4, 71.1)	708	70.5 (67.1, 74.0)	635	72.1 (68.6, 75.6)
25-34	Current smoker	734	30.7 (28.6, 32.9)	670	28.9 (26.8, 31.0)	572	28.6 (26.3, 30.9)	500	27.0 (24.7, 29.3)	466	27.0 (24.6, 29.4)
	Former smoker	477	18.3 (16.6, 20.1)	475	19.0 (17.3, 20.8)	377	17.5 (15.7, 19.3)	346	17.8 (15.9, 19.7)	311	16.4 (14.5, 18.3)
	Never smoker	1406	50.9 (48.7, 53.2)	1424	52.1 (49.8, 54.3)	1275	53.9 (51.4, 56.3)	1182	55.2 (52.7, 57.8)	1123	56.6 (54.0, 59.2)
35-44	Current smoker	758	27.2 (25.2, 29.2)	688	25.0 (23.1, 26.8)	593	24.3 (22.3, 26.3)	478	22.6 (20.4, 24.7)	426	22.6 (20.4, 24.8)
	Former smoker	531	18.9 (17.2, 20.7)	564	20.2 (18.5, 21.9)	525	22.0 (20.0, 23.9)	493	22.4 (20.3, 24.5)	437	22.2 (20.0, 24.4)
	Never smoker	1682	53.8 (51.7, 56.0)	1670	54.9 (52.7, 57.0)	1490	53.7 (51.4, 56.1)	1357	55.0 (52.5, 57.5)	1182	55.2 (52.6, 57.8)
45-54	Current smoker	1158	29.6 (27.9, 31.3)	982	26.2 (24.6, 27.8)	827	25.9 (24.0, 27.7)	676	24.2 (22.3, 26.1)	573	23.8 (21.7, 25.8)
	Former smoker	931	23.4 (21.8, 24.9)	888	23.7 (22.1, 25.2)	821	24.7 (23.0, 26.4)	745	25.2 (23.4, 27.1)	615	24.6 (22.6, 26.5)
	Never smoker	1965	47.1 (45.2, 48.9)	1986	50.1 (48.3, 52.0)	1827	49.4 (47.4, 51.4)	1716	50.6 (48.5, 52.7)	1517	51.7 (49.4, 53.9)
55-64	Current smoker	1181	22.7 (21.3, 24.1)	1129	21.5 (20.2, 22.9)	1044	21.5 (20.0, 22.9)	907	20.2 (18.8, 21.7)	785	20.4 (18.8, 21.9)
	Former smoker	1614	31.5 (30.0, 33.1)	1582	31.3 (29.9, 32.8)	1527	31.2 (29.6, 32.7)	1435	31.3 (29.7, 32.9)	1277	31.4 (29.7, 33.1)
	Never smoker	2537	45.8 (44.2, 47.4)	2612	47.1 (45.5, 48.7)	2524	47.4 (45.7, 49.0)	2403	48.5 (46.7, 50.2)	2153	48.2 (46.4, 50.1)
65 or more	Current smoker	933	10.7 (9.9, 11.5)	867	10.6 (9.7, 11.4)	870	11.4 (10.5, 12.3)	829	11.1 (10.2, 12.0)	754	10.8 (9.8, 11.7)
	Former smoker	3350	42.1 (40.8, 43.4)	3322	42.1 (40.8, 43.4)	3386	41.1 (39.8, 42.4)	3221	40.8 (39.5, 42.1)	3061	40.4 (39.1, 41.7)
	Never smoker	4195	47.2 (45.9, 48.5)	4227	47.3 (46.0, 48.6)	4373	47.5 (46.2, 48.8)	4376	48.0 (46.7, 49.3)	4276	48.9 (47.5, 50.2)

Age Group	Smoking Status	2016-18		2017-19		2018-20		2019-21		2020-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
18-24	Current smoker	154	19.8 (16.7, 22.9)	150	18.3 (15.4, 21.3)	121	15.0 (12.2, 17.8)	106	11.9 (9.6, 14.3)	89	10.0 (7.8, 12.1)
	Former smoker	57	5.6 (4.0, 7.2)	64	6.6 (4.8, 8.4)	71	7.5 (5.6, 9.5)	79	8.5 (6.5, 10.6)	86	8.6 (6.7, 10.6)
	Never smoker	661	74.6 (71.3, 78.0)	674	75.1 (71.9, 78.4)	691	77.5 (74.3, 80.6)	752	79.5 (76.6, 82.4)	789	81.4 (78.6, 84.2)
25-34	Current smoker	430	25.4 (23.1, 27.8)	404	24.0 (21.7, 26.4)	347	22.4 (20.1, 24.8)	318	20.0 (17.7, 22.2)	288	18.3 (16.1, 20.5)
	Former smoker	315	16.9 (14.9, 18.8)	312	16.5 (14.6, 18.4)	309	17.4 (15.4, 19.5)	330	19.6 (17.4, 21.8)	339	20.7 (18.4, 22.9)
	Never smoker	1107	57.7 (55.0, 60.3)	1159	59.5 (56.9, 62.0)	1088	60.1 (57.5, 62.8)	1120	60.5 (57.8, 63.1)	1120	61.0 (58.3, 63.7)
35-44	Current smoker	408	21.2 (19.1, 23.4)	452	21.6 (19.6, 23.6)	433	21.8 (19.7, 23.9)	457	22.9 (20.8, 25.1)	421	22.0 (19.9, 24.1)
	Former smoker	454	23.4 (21.2, 25.6)	487	24.0 (21.8, 26.1)	488	25.6 (23.3, 27.8)	518	25.0 (22.8, 27.1)	543	26.4 (24.2, 28.6)
	Never smoker	1118	55.3 (52.7, 57.9)	1150	54.5 (52.0, 56.9)	1077	52.6 (50.1, 55.2)	1159	52.1 (49.6, 54.6)	1123	51.6 (49.1, 54.1)
45-54	Current smoker	535	23.7 (21.6, 25.7)	521	23.4 (21.4, 25.5)	502	23.7 (21.6, 25.8)	469	21.9 (19.9, 24.0)	424	20.2 (18.2, 22.2)
	Former smoker	548	23.9 (21.8, 25.9)	532	23.5 (21.4, 25.5)	511	23.0 (20.9, 25.0)	532	23.7 (21.7, 25.8)	552	25.4 (23.3, 27.5)
	Never smoker	1390	52.5 (50.2, 54.8)	1323	53.1 (50.7, 55.4)	1245	53.3 (50.9, 55.7)	1275	54.3 (51.9, 56.8)	1242	54.4 (52.0, 56.8)
55-64	Current smoker	670	20.5 (18.9, 22.2)	638	20.9 (19.2, 22.6)	583	21.8 (19.9, 23.7)	574	21.9 (20.0, 23.8)	548	21.5 (19.6, 23.4)
	Former smoker	1107	30.9 (29.1, 32.7)	1017	31.1 (29.2, 32.9)	851	29.9 (27.9, 31.9)	813	29.6 (27.5, 31.6)	763	29.4 (27.3, 31.4)
	Never smoker	1932	48.6 (46.6, 50.5)	1733	48.1 (46.1, 50.0)	1503	48.3 (46.2, 50.4)	1421	48.5 (46.3, 50.7)	1373	49.1 (46.9, 51.4)
65 or more	Current smoker	660	10.3 (9.4, 11.3)	651	11.1 (10.2, 12.1)	620	12.0 (10.9, 13.0)	610	12.1 (11.0, 13.2)	573	11.9 (10.7, 13.0)
	Former smoker	2728	40.5 (39.1, 41.9)	2582	40.4 (38.9, 41.8)	2215	39.7 (38.2, 41.2)	2142	38.6 (37.0, 40.2)	1987	38.2 (36.6, 39.8)
	Never smoker	3847	49.2 (47.8, 50.6)	3618	48.5 (47.1, 49.9)	3163	48.3 (46.8, 49.9)	3176	49.3 (47.7, 50.9)	2925	50.0 (48.3, 51.6)

Age Group	Smoking Status	2021-23	
		N	Weighted% (95%CI)
18-24	Current smoker	90	9.3 (7.3, 11.4)
	Former smoker	86	8.3 (6.4, 10.2)
	Never smoker	867	82.3 (79.7, 84.9)
25-34	Current smoker	280	15.7 (13.7, 17.6)
	Former smoker	414	22.9 (20.7, 25.2)
	Never smoker	1262	61.4 (58.9, 63.9)
35-44	Current smoker	461	20.5 (18.6, 22.4)
	Former smoker	643	26.3 (24.3, 28.3)
	Never smoker	1358	53.2 (50.9, 55.5)
45-54	Current smoker	440	19.2 (17.3, 21.1)
	Former smoker	636	26.1 (24.1, 28.2)
	Never smoker	1393	54.7 (52.4, 57.0)
55-64	Current smoker	576	21.2 (19.4, 23.0)
	Former smoker	825	28.2 (26.3, 30.1)
	Never smoker	1555	50.6 (48.4, 52.7)
65 or more	Current smoker	598	11.6 (10.5, 12.7)
	Former smoker	2179	38.0 (36.5, 39.6)
	Never smoker	3157	50.3 (48.8, 51.9)

Table 4. Three-year rolling percentage of current cigarette smoking among Oklahoma adults by race/ethnicity, Oklahoma BRFSS 2011 – 2023

Race/Ethnicity	Smoking Status	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
NH-White	Current smoker	3521	23.7 (22.8, 24.6)	3240	21.7 (20.9, 22.6)	2907	21.4 (20.5, 22.2)	2585	20.3 (19.4, 21.3)	2280	19.9 (18.9, 20.9)
	Former smoker	5464	26.3 (25.5, 27.1)	5456	27.4 (26.6, 28.2)	5340	27.8 (27.0, 28.7)	5038	27.8 (26.9, 28.7)	4604	27.1 (26.2, 28.1)
	Never smoker	9320	50.1 (49.1, 51.0)	9447	50.9 (50.0, 51.8)	9118	50.8 (49.8, 51.8)	8916	51.9 (50.8, 52.9)	8352	53.0 (51.9, 54.1)
NH-Black	Current smoker	387	26.7 (23.5, 29.9)	318	25.0 (22.0, 28.0)	295	25.5 (22.1, 28.8)	227	23.0 (19.5, 26.4)	189	20.7 (17.2, 24.2)
	Former smoker	307	17.4 (14.8, 20.0)	295	17.1 (14.8, 19.4)	318	18.8 (16.2, 21.3)	300	19.4 (16.7, 22.1)	271	19.0 (16.1, 21.8)
	Never smoker	777	55.9 (52.3, 59.5)	781	57.9 (54.6, 61.3)	751	55.8 (52.2, 59.3)	667	57.7 (53.9, 61.5)	624	60.3 (56.4, 64.3)
NH-AI/AN	Current smoker	451	33.6 (30.3, 36.8)	417	31.2 (28.0, 34.3)	405	32.3 (29.0, 35.6)	349	27.8 (24.6, 31.0)	321	28.9 (25.5, 32.2)
	Former smoker	428	24.1 (21.5, 26.7)	429	23.3 (20.8, 25.7)	388	20.9 (18.5, 23.4)	390	23.3 (20.6, 26.0)	333	22.1 (19.4, 24.9)
	Never smoker	622	42.3 (39.0, 45.6)	694	45.6 (42.3, 48.8)	707	46.8 (43.4, 50.2)	697	48.8 (45.4, 52.3)	623	49.0 (45.4, 52.6)
NH-Other	Current smoker	455	24.4 (21.5, 27.3)	439	27.2 (23.8, 30.5)	350	26.3 (22.7, 30.0)	268	24.0 (20.2, 27.7)	210	21.8 (18.2, 25.4)
	Former smoker	540	21.4 (18.8, 24.0)	488	18.8 (16.4, 21.2)	421	17.7 (15.2, 20.2)	360	17.6 (15.0, 20.1)	332	19.1 (16.4, 21.8)
	Never smoker	935	54.2 (50.8, 57.7)	890	54.0 (50.5, 57.6)	809	56.0 (52.1, 59.9)	722	58.5 (54.4, 62.5)	643	59.1 (55.1, 63.2)
Hispanic	Current smoker	229	19.9 (17.0, 22.8)	200	18.2 (15.5, 20.9)	163	16.4 (13.4, 19.4)	128	16.1 (12.9, 19.4)	131	18.2 (14.7, 21.8)
	Former smoker	244	17.9 (15.3, 20.5)	234	16.7 (14.3, 19.2)	214	15.6 (13.1, 18.1)	181	15.1 (12.5, 17.6)	174	16.1 (13.3, 18.9)
	Never smoker	810	62.2 (58.8, 65.7)	835	65.1 (61.8, 68.3)	762	68.0 (64.5, 71.5)	664	68.8 (65.1, 72.5)	551	65.7 (61.7, 69.7)

Race/Ethnicity	Smoking Status	2016-18		2017-19		2018-20		2019-21		2020-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
NH-White	Current smoker	2052	19.3 (18.4, 20.3)	1969	19.0 (18.1, 19.9)	1805	19.0 (18.0, 20.0)	1737	18.0 (17.0, 18.9)	1623	17.1 (16.2, 18.0)
	Former smoker	4142	27.0 (26.1, 27.9)	3935	27.2 (26.2, 28.1)	3470	27.6 (26.6, 28.6)	3399	27.9 (26.9, 28.9)	3253	28.6 (27.5, 29.6)
	Never smoker	7660	53.7 (52.6, 54.8)	7225	53.8 (52.7, 54.9)	6432	53.4 (52.3, 54.6)	6446	54.2 (53.0, 55.3)	6120	54.3 (53.2, 55.5)
NH-Black	Current smoker	167	19.4 (16.2, 22.7)	165	18.9 (15.8, 21.9)	151	19.9 (16.6, 23.2)	139	19.1 (15.8, 22.4)	125	18.2 (14.9, 21.5)
	Former smoker	227	18.1 (15.2, 21.0)	204	18.3 (15.3, 21.2)	175	17.3 (14.3, 20.3)	174	17.6 (14.6, 20.6)	170	18.5 (15.4, 21.6)
	Never smoker	569	62.4 (58.5, 66.3)	552	62.9 (59.1, 66.7)	495	62.8 (58.8, 66.8)	505	63.3 (59.3, 67.2)	479	63.3 (59.3, 67.3)
NH-AI/AN	Current smoker	256	25.3 (22.0, 28.6)	279	27.4 (24.1, 30.7)	277	26.4 (23.0, 29.8)	304	27.8 (24.5, 31.1)	269	24.2 (21.1, 27.3)
	Former smoker	336	25.3 (22.3, 28.3)	330	24.9 (21.9, 27.9)	335	26.3 (23.2, 29.3)	322	25.1 (22.1, 28.1)	323	26.2 (23.2, 29.1)
	Never smoker	559	49.5 (45.8, 53.1)	546	47.7 (44.1, 51.3)	544	47.4 (43.7, 51.0)	567	47.1 (43.6, 50.6)	567	49.7 (46.1, 53.2)
NH-Other	Current smoker	223	23.5 (20.0, 27.0)	236	22.6 (19.3, 26.0)	222	22.3 (18.9, 25.8)	190	18.3 (15.1, 21.4)	176	18.6 (15.4, 21.7)
	Former smoker	301	18.7 (16.1, 21.4)	293	20.6 (17.8, 23.5)	253	22.0 (18.8, 25.1)	261	22.6 (19.3, 25.8)	262	23.4 (20.3, 26.6)
	Never smoker	611	57.7 (53.8, 61.7)	595	56.7 (52.9, 60.6)	529	55.7 (51.7, 59.8)	559	59.2 (55.2, 63.2)	560	58.0 (54.1, 61.9)
Hispanic	Current smoker	129	16.4 (13.2, 19.5)	136	15.3 (12.4, 18.2)	125	12.0 (9.7, 14.4)	128	12.0 (9.6, 14.4)	115	10.5 (8.3, 12.8)
	Former smoker	164	16.3 (13.4, 19.2)	176	15.6 (12.9, 18.3)	164	13.9 (11.4, 16.4)	180	14.9 (12.3, 17.5)	184	15.9 (13.3, 18.6)
	Never smoker	561	67.3 (63.5, 71.2)	635	69.1 (65.5, 72.7)	673	74.1 (70.9, 77.3)	708	73.1 (69.8, 76.3)	716	73.5 (70.3, 76.7)

Race/Ethnicity	Smoking Status	2021-23	
		N	Weighted% (95%CI)
NH-White	Current smoker	1687	15.9 (15.1, 16.8)
	Former smoker	3620	28.8 (27.8, 29.8)
	Never smoker	6786	55.3 (54.2, 56.4)
NH-Black	Current smoker	125	16.0 (13.1, 18.9)
	Former smoker	183	19.5 (16.5, 22.6)
	Never smoker	521	64.5 (60.7, 68.3)
NH-AI/AN	Current smoker	263	23.0 (20.1, 25.8)
	Former smoker	353	27.0 (24.0, 29.9)
	Never smoker	618	50.1 (46.7, 53.4)
NH-Other	Current smoker	192	17.4 (14.7, 20.2)
	Former smoker	318	23.8 (21.0, 26.6)
	Never smoker	685	58.8 (55.3, 62.2)
Hispanic	Current smoker	139	11.4 (9.3, 13.5)
	Former smoker	221	16.8 (14.4, 19.2)
	Never smoker	822	71.8 (68.8, 74.8)

Table 5. Three-year rolling percentage of current cigarette smoking among Oklahoma adults by level of education, Oklahoma BRFSS 2011 – 2023

Education	Smoking Status	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than high school	Current smoker	949	40.0 (37.4, 42.6)	824	37.4 (34.9, 40.0)	671	37.3 (34.3, 40.2)	520	36.3 (33.1, 39.4)	423	35.8 (32.4, 39.2)
	Former smoker	800	23.1 (21.1, 25.0)	740	23.7 (21.7, 25.7)	661	23.7 (21.5, 25.9)	570	24.8 (22.4, 27.3)	489	24.9 (22.2, 27.5)
	Never smoker	1119	36.9 (34.4, 39.4)	1066	38.9 (36.4, 41.4)	891	39.0 (36.2, 41.8)	734	38.9 (35.9, 41.9)	579	39.3 (35.9, 42.7)
High school	Current smoker	1932	27.7 (26.3, 29.0)	1783	26.3 (25.0, 27.6)	1571	25.7 (24.3, 27.1)	1369	24.2 (22.7, 25.6)	1192	23.7 (22.2, 25.2)
	Former smoker	2240	24.6 (23.4, 25.8)	2255	25.4 (24.2, 26.5)	2206	26.0 (24.7, 27.3)	2076	26.2 (24.8, 27.5)	1808	25.0 (23.6, 26.4)
	Never smoker	3588	47.7 (46.2, 49.3)	3590	48.4 (46.9, 49.8)	3440	48.3 (46.7, 49.9)	3273	49.7 (48.0, 51.3)	3011	51.3 (49.6, 53.0)
Some college	Current smoker	1497	23.3 (22.0, 24.6)	1388	21.6 (20.3, 22.8)	1317	21.3 (20.0, 22.6)	1182	19.5 (18.2, 20.8)	1086	19.5 (18.2, 20.9)
	Former smoker	2089	25.7 (24.4, 26.9)	2052	25.8 (24.6, 27.0)	1996	25.7 (24.4, 27.0)	1942	26.1 (24.7, 27.4)	1860	26.0 (24.6, 27.4)
	Never smoker	3431	51.0 (49.5, 52.6)	3489	52.6 (51.1, 54.1)	3378	52.9 (51.4, 54.5)	3337	54.4 (52.8, 56.0)	3238	54.4 (52.8, 56.1)
College graduate	Current smoker	678	9.5 (8.6, 10.4)	633	8.5 (7.7, 9.4)	580	8.5 (7.6, 9.4)	511	8.3 (7.4, 9.3)	461	8.4 (7.4, 9.3)
	Former smoker	1871	23.7 (22.5, 25.0)	1876	23.9 (22.7, 25.1)	1851	23.5 (22.3, 24.7)	1712	22.5 (21.3, 23.8)	1592	22.1 (20.8, 23.3)
	Never smoker	4364	66.7 (65.4, 68.1)	4548	67.6 (66.3, 68.9)	4483	68.0 (66.6, 69.4)	4371	69.1 (67.7, 70.6)	4032	69.6 (68.1, 71.0)

Education	Smoking Status	2016-18		2017-19		2018-20		2019-21		2020-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than high school	Current smoker	388	33.9 (30.6, 37.1)	397	33.6 (30.3, 36.8)	377	33.6 (30.3, 36.9)	362	33.2 (29.9, 36.5)	323	30.6 (27.3, 33.9)
	Former smoker	429	25.6 (22.8, 28.4)	401	24.9 (22.2, 27.7)	347	24.4 (21.6, 27.2)	321	23.7 (20.8, 26.5)	294	24.5 (21.6, 27.5)
	Never smoker	521	40.5 (37.1, 44.0)	530	41.5 (38.2, 44.9)	503	42.0 (38.6, 45.4)	496	43.1 (39.7, 46.6)	471	44.9 (41.3, 48.4)
High school	Current smoker	1062	23.1 (21.5, 24.6)	1064	23.3 (21.8, 24.8)	968	22.7 (21.2, 24.2)	920	21.7 (20.2, 23.1)	799	20.1 (18.6, 21.6)
	Former smoker	1573	24.7 (23.3, 26.1)	1513	25.7 (24.2, 27.1)	1364	26.6 (25.0, 28.1)	1306	26.6 (25.1, 28.2)	1229	27.2 (25.6, 28.8)
	Never smoker	2681	52.3 (50.5, 54.0)	2490	51.0 (49.2, 52.8)	2191	50.7 (48.9, 52.6)	2187	51.7 (49.9, 53.5)	2058	52.7 (50.9, 54.6)
Some college	Current smoker	973	18.9 (17.6, 20.3)	930	18.8 (17.4, 20.1)	875	18.6 (17.3, 19.9)	884	17.8 (16.5, 19.1)	866	17.3 (16.0, 18.5)
	Former smoker	1733	26.6 (25.1, 28.0)	1653	26.4 (25.0, 27.8)	1473	26.6 (25.1, 28.0)	1497	27.4 (25.9, 28.9)	1475	28.3 (26.7, 29.8)
	Never smoker	3013	54.5 (52.9, 56.2)	2841	54.9 (53.2, 56.5)	2546	54.8 (53.1, 56.5)	2556	54.8 (53.1, 56.5)	2500	54.5 (52.8, 56.2)
College graduate	Current smoker	429	8.1 (7.2, 9.1)	420	7.6 (6.7, 8.5)	382	7.5 (6.6, 8.4)	364	6.6 (5.8, 7.4)	353	6.5 (5.7, 7.3)
	Former smoker	1462	21.6 (20.4, 22.9)	1417	21.6 (20.4, 22.9)	1253	21.6 (20.3, 22.9)	1278	22.0 (20.7, 23.3)	1256	22.4 (21.0, 23.7)
	Never smoker	3818	70.2 (68.8, 71.7)	3775	70.8 (69.3, 72.2)	3502	70.9 (69.5, 72.4)	3628	71.4 (69.9, 72.9)	3494	71.1 (69.7, 72.6)

Education	Smoking Status	2021-23	
		N	Weighted% (95%CI)
Less than high school	Current smoker	327	29.4 (26.3, 32.5)
	Former smoker	301	24.9 (22.0, 27.9)
	Never smoker	470	45.7 (42.2, 49.2)
High school	Current smoker	825	18.9 (17.5, 20.3)
	Former smoker	1323	27.4 (25.8, 28.9)
	Never smoker	2235	53.7 (51.9, 55.5)
Some college	Current smoker	892	16.1 (15.0, 17.2)
	Former smoker	1649	28.7 (27.3, 30.1)
	Never smoker	2753	55.2 (53.6, 56.8)
College graduate	Current smoker	398	6.0 (5.3, 6.6)
	Former smoker	1496	22.6 (21.4, 23.8)
	Never smoker	4084	71.5 (70.2, 72.8)

Table 6. Three-year rolling percentage of current cigarette smoking among Oklahoma adults by income level, Oklahoma BRFSS 2011 – 2023

Income	Smoking Status	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than 15,000	Current smoker	1313	43.1 (40.7, 45.6)	1116	39.7 (37.3, 42.0)	927	39.0 (36.3, 41.7)	698	36.5 (33.6, 39.5)	574	35.4 (32.2, 38.5)
	Former smoker	850	19.6 (17.9, 21.4)	798	20.7 (18.9, 22.5)	698	21.6 (19.5, 23.6)	572	22.1 (19.8, 24.3)	470	21.9 (19.4, 24.3)
	Never smoker	1256	37.3 (34.8, 39.7)	1184	39.7 (37.3, 42.1)	1014	39.4 (36.7, 42.2)	875	41.4 (38.3, 44.5)	752	42.8 (39.5, 46.1)
15,000 to Less than 25,000	Current smoker	1199	32.4 (30.4, 34.4)	1106	31.1 (29.1, 33.1)	978	31.5 (29.3, 33.7)	826	30.3 (28.0, 32.6)	693	30.1 (27.6, 32.5)
	Former smoker	1333	24.1 (22.5, 25.6)	1262	24.1 (22.6, 25.7)	1182	24.5 (22.7, 26.2)	1074	25.2 (23.3, 27.1)	931	24.0 (22.0, 25.9)
	Never smoker	2004	43.6 (41.6, 45.6)	1960	44.7 (42.7, 46.7)	1775	44.0 (41.8, 46.2)	1611	44.6 (42.2, 46.9)	1461	46.0 (43.4, 48.5)
25,000 to Less than 35,000	Current smoker	603	26.7 (24.3, 29.0)	556	25.9 (23.6, 28.3)	454	23.2 (20.7, 25.6)	403	22.5 (19.9, 25.0)	348	22.0 (19.3, 24.7)
	Former smoker	806	26.3 (24.1, 28.5)	813	26.3 (24.2, 28.4)	777	26.0 (23.8, 28.2)	715	26.1 (23.8, 28.5)	633	25.8 (23.4, 28.3)
	Never smoker	1306	47.0 (44.4, 49.6)	1317	47.7 (45.2, 50.3)	1257	50.8 (48.1, 53.6)	1145	51.4 (48.5, 54.3)	1002	52.1 (49.0, 55.2)
35,000 to Less than 50,000	Current smoker	592	21.5 (19.5, 23.4)	534	19.9 (18.1, 21.7)	484	19.7 (17.7, 21.6)	415	19.1 (16.9, 21.2)	383	20.0 (17.7, 22.3)
	Former smoker	1053	27.9 (26.0, 29.8)	997	27.8 (26.0, 29.7)	964	27.8 (25.8, 29.8)	893	27.2 (25.1, 29.3)	814	27.7 (25.5, 30.0)
	Never smoker	1674	50.6 (48.4, 52.9)	1686	52.2 (50.0, 54.4)	1613	52.5 (50.2, 54.9)	1511	53.7 (51.2, 56.3)	1343	52.3 (49.6, 55.0)
50,000 or more	Current smoker	922	14.3 (13.3, 15.4)	879	13.0 (12.1, 14.0)	832	13.9 (12.7, 15.0)	779	13.6 (12.5, 14.8)	691	13.7 (12.4, 14.9)
	Former smoker	2208	26.3 (25.1, 27.5)	2325	27.3 (26.1, 28.4)	2254	26.9 (25.6, 28.1)	2128	26.8 (25.5, 28.2)	1921	26.2 (24.9, 27.6)
	Never smoker	4580	59.4 (58.0, 60.7)	4785	59.7 (58.4, 61.0)	4635	59.3 (57.8, 60.7)	4488	59.5 (58.0, 61.0)	4084	60.1 (58.5, 61.7)

Income	Smoking Status	2016-18		2017-19		2018-20		2019-21		2020-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than 15,000	Current smoker	479	34.8 (31.6, 38.0)	463	34.5 (31.3, 37.7)	402	34.7 (31.2, 38.1)	363	33.0 (29.5, 36.5)	299	32.3 (28.6, 36.0)
	Former smoker	405	21.3 (18.8, 23.7)	376	21.7 (19.2, 24.3)	299	21.2 (18.4, 24.0)	285	23.3 (20.2, 26.4)	258	24.2 (20.9, 27.5)
	Never smoker	663	43.9 (40.6, 47.3)	573	43.7 (40.3, 47.2)	501	44.1 (40.5, 47.7)	460	43.7 (40.0, 47.4)	414	43.5 (39.6, 47.3)
15,000 to Less than 25,000	Current smoker	602	29.0 (26.5, 31.5)	608	29.8 (27.3, 32.3)	571	28.4 (25.9, 30.8)	524	27.7 (25.2, 30.2)	431	26.1 (23.5, 28.7)
	Former smoker	799	23.4 (21.4, 25.4)	731	22.6 (20.7, 24.6)	665	23.7 (21.6, 25.7)	585	23.7 (21.6, 25.9)	491	25.0 (22.6, 27.5)
	Never smoker	1339	47.7 (45.1, 50.2)	1281	47.6 (45.1, 50.2)	1144	48.0 (45.4, 50.6)	1031	48.6 (45.9, 51.3)	799	48.9 (45.9, 51.9)
25,000 to Less than 35,000	Current smoker	351	24.6 (21.8, 27.4)	324	23.7 (20.9, 26.5)	313	24.4 (21.6, 27.3)	316	23.5 (20.8, 26.3)	346	22.6 (20.1, 25.1)
	Former smoker	566	25.7 (23.1, 28.3)	533	26.1 (23.5, 28.8)	463	26.9 (24.1, 29.7)	471	27.2 (24.4, 29.9)	514	27.7 (25.1, 30.3)
	Never smoker	886	49.7 (46.5, 52.8)	848	50.2 (47.0, 53.3)	761	48.7 (45.4, 51.9)	796	49.3 (46.1, 52.4)	852	49.7 (46.7, 52.7)
35,000 to Less than 50,000	Current smoker	364	20.3 (18.0, 22.6)	345	19.2 (16.9, 21.4)	307	18.7 (16.3, 21.1)	312	18.9 (16.5, 21.3)	293	18.1 (15.7, 20.5)
	Former smoker	732	27.6 (25.3, 29.9)	694	28.2 (25.9, 30.5)	627	27.1 (24.7, 29.4)	573	25.9 (23.5, 28.2)	546	25.8 (23.4, 28.2)
	Never smoker	1221	52.1 (49.3, 54.8)	1148	52.6 (49.9, 55.4)	1107	54.2 (51.4, 57.0)	1110	55.2 (52.4, 58.1)	1099	56.1 (53.3, 58.9)
50,000 or more	Current smoker	639	12.2 (11.1, 13.3)	664	12.2 (11.1, 13.3)	641	12.0 (10.9, 13.1)	634	11.6 (10.6, 12.7)	602	10.6 (9.6, 11.6)
	Former smoker	1847	27.0 (25.6, 28.3)	1865	26.8 (25.5, 28.2)	1702	26.8 (25.5, 28.2)	1742	27.4 (26.0, 28.8)	1745	27.9 (26.6, 29.3)
	Never smoker	3933	60.8 (59.3, 62.4)	3977	60.9 (59.4, 62.4)	3691	61.2 (59.6, 62.7)	3756	60.9 (59.4, 62.5)	3745	61.5 (60.0, 63.0)

Income	Smoking Status	2021-23	
		N	Weighted% (95%CI)
Less than 15,000	Current smoker	285	33.2 (29.4, 37.1)
	Former smoker	242	24.3 (20.9, 27.8)
	Never smoker	365	42.4 (38.4, 46.5)
15,000 to Less than 25,000	Current smoker	357	25.6 (22.8, 28.4)
	Former smoker	429	26.4 (23.6, 29.2)
	Never smoker	647	48.0 (44.7, 51.3)
25,000 to Less than 35,000	Current smoker	369	22.1 (19.7, 24.4)
	Former smoker	562	27.9 (25.4, 30.3)
	Never smoker	918	50.1 (47.2, 52.9)
35,000 to Less than 50,000	Current smoker	363	17.5 (15.5, 19.4)
	Former smoker	648	26.2 (24.0, 28.4)
	Never smoker	1244	56.3 (53.8, 58.9)
50,000 or more	Current smoker	705	10.4 (9.5, 11.3)
	Former smoker	2150	27.9 (26.7, 29.2)
	Never smoker	4645	61.7 (60.3, 63.0)

Table 7. Area of residence and current smoking status by year, Oklahoma BRFSS 2011 – 2023

Area of Residence	Smoking Status	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Urban	Current smoker	807	25.7 (23.6, 27.8)	508	19.3 (17.4, 21.1)	581	20.9 (19.0, 22.8)	466	18.4 (16.6, 20.3)	378	20.3 (17.7, 22.8)
	Former smoker	1057	23.2 (21.4, 25)	748	24.2 (22.4, 26.1)	937	24.2 (22.4, 26.0)	870	25.1 (23.3, 27.0)	699	23.3 (21.1, 25.4)
	Never smoker	1865	51.1 (48.8, 53.4)	1537	56.5 (54.2, 58.8)	1810	54.9 (52.7, 57.1)	1657	56.4 (54.2, 58.7)	1351	56.5 (53.6, 59.3)
Rural	Current smoker	1050	26.4 (24.5, 28.3)	1128	26.2 (24.4, 28)	986	25.8 (24, 27.5)	961	22.9 (21.3, 24.6)	770	23.6 (21.6, 25.6)
	Former smoker	1425	25.8 (24.2, 27.4)	1453	24.5 (23, 25.9)	1390	24.8 (23.3, 26.3)	1534	26.5 (25.0, 28.0)	1293	25.8 (23.9, 27.6)
	Never smoker	2304	47.8 (45.7, 49.8)	2574	49.3 (47.4, 51.2)	2430	49.5 (47.6, 51.4)	2709	50.5 (48.7, 52.3)	2262	50.6 (48.4, 52.9)

Area of Residence	Smoking Status	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Urban	Current smoker	308	16.8 (14.6, 19.0)	322	18.5 (16.2, 20.7)	251	16.2 (14.1, 18.4)	272	16.6 (14.5, 18.7)	236	17.3 (15.0, 19.7)
	Former smoker	662	22.2 (20.2, 24.2)	642	24.2 (22.0, 26.4)	519	23.9 (21.6, 26.3)	551	24.1 (22.0, 26.2)	429	23.3 (20.9, 25.7)
	Never smoker	1441	61.1 (58.4, 63.7)	1261	57.3 (54.6, 60.0)	1084	59.8 (57.1, 62.6)	1208	59.3 (56.7, 61.9)	962	59.4 (56.5, 62.3)
Rural	Current smoker	702	21.7 (19.8, 23.6)	686	21.4 (19.7, 23.2)	588	22.3 (20.3, 24.3)	697	20.6 (19.0, 22.3)	562	20.4 (18.6, 22.3)
	Former smoker	1252	26.8 (25.1, 28.6)	1214	24.8 (23.2, 26.4)	920	25.5 (23.7, 27.4)	1148	26.2 (24.5, 27.9)	878	26.7 (24.8, 28.6)
	Never smoker	2325	51.5 (49.4, 53.5)	2246	53.8 (51.8, 55.8)	1698	52.1 (49.9, 54.4)	2160	53.2 (51.2, 55.2)	1655	52.9 (50.7, 55.1)

Area of Residence	Smoking Status	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Urban	Current smoker	191	13.0 (10.9, 15.0)	242	13.5 (11.7, 15.3)	262	13.0 (11.4, 14.7)
	Former smoker	417	26.9 (24.3, 29.6)	522	26.3 (24.1, 28.5)	644	24.8 (22.8, 26.7)
	Never smoker	934	60.1 (57.1, 63.1)	1099	60.2 (57.6, 62.8)	1405	62.2 (59.9, 64.5)
Rural	Current smoker	576	20.0 (18.2, 21.7)	536	17.3 (15.7, 18.9)	638	18.0 (16.5, 19.6)
	Former smoker	991	24.7 (23.0, 26.5)	1033	27.9 (26.1, 29.6)	1176	27.1 (25.5, 28.8)
	Never smoker	1984	55.3 (53.2, 57.4)	1938	54.9 (52.9, 56.9)	2232	54.8 (52.9, 56.8)

Table 8. Physical health (14 or more physically unhealthy days) and current smoking status by year, Oklahoma BRFSS 2011 – 2023

Poor Physical Health	Smoking Status	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Current smoker	466	36.7 (33.2, 40.2)	403	34.4 (30.9, 38.0)	388	34.2 (30.8, 37.7)	350	31.2 (27.8, 34.5)	281	32.4 (28.0, 36.8)
	Former smoker	491	30.2 (27.0, 33.3)	411	29.7 (26.5, 32.9)	431	29.3 (26.3, 32.4)	494	33.7 (30.5, 36.8)	400	32.7 (28.8, 36.6)
	Never smoker	507	33.1 (29.8, 36.5)	519	35.8 (32.4, 39.3)	516	36.4 (33.1, 39.8)	494	35.2 (31.9, 38.4)	417	34.9 (30.7, 39.1)
No	Current smoker	1391	24.5 (22.9, 26.0)	1233	21.5 (20.1, 22.9)	1179	22.0 (20.6, 23.4)	1077	19.4 (18.0, 20.7)	867	20.4 (18.7, 22.1)
	Former smoker	1991	23.9 (22.5, 25.2)	1790	23.5 (22.3, 24.8)	1896	23.8 (22.5, 25.0)	1910	24.7 (23.4, 25.9)	1592	23.3 (21.8, 24.8)
	Never smoker	3662	51.7 (50.0, 53.4)	3592	55.0 (53.4, 56.6)	3724	54.2 (52.7, 55.8)	3872	56.0 (54.4, 57.5)	3196	56.3 (54.4, 58.2)

Poor Physical Health	Smoking Status	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Current smoker	260	31.8 (27.8, 35.9)	250	29.8 (25.9, 33.7)	206	31.6 (27.3, 35.9)	242	30.8 (26.8, 34.8)	186	31.6 (27.0, 36.1)
	Former smoker	358	30.9 (27.3, 34.4)	348	30.8 (27.2, 34.4)	268	28.1 (24.2, 31.9)	295	29.0 (25.5, 32.5)	204	30.9 (26.4, 35.3)
	Never smoker	464	37.3 (33.5, 41.1)	413	39.4 (35.3, 43.6)	339	40.3 (35.9, 44.7)	378	40.2 (36.2, 44.2)	268	37.6 (33.0, 42.2)
No	Current smoker	707	17.0 (15.5, 18.5)	721	18.2 (16.7, 19.7)	605	17.3 (15.7, 18.8)	680	16.6 (15.2, 18.0)	580	17.0 (15.5, 18.6)
	Former smoker	1504	23.7 (22.3, 25.2)	1432	23.2 (21.7, 24.6)	1131	24.1 (22.5, 25.7)	1338	24.6 (23.1, 26.1)	1055	24.4 (22.7, 26.0)
	Never smoker	3205	59.3 (57.5, 61.1)	3016	58.7 (56.9, 60.4)	2389	58.6 (56.7, 60.5)	2862	58.9 (57.1, 60.6)	2261	58.6 (56.7, 60.6)

Poor Physical Health	Smoking Status	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Current smoker	167	29.0 (24.6, 33.4)	201	27.4 (23.7, 31.1)	202	26.6 (22.9, 30.4)
	Former smoker	214	28.8 (24.6, 33.0)	275	32.8 (29.0, 36.5)	304	32.6 (28.8, 36.4)
	Never smoker	297	42.2 (37.6, 46.9)	337	39.9 (35.9, 43.9)	370	40.8 (36.8, 44.8)
No	Current smoker	557	14.9 (13.5, 16.3)	548	13.5 (12.3, 14.8)	651	13.7 (12.5, 14.8)
	Former smoker	1131	25.2 (23.6, 26.9)	1222	26.1 (24.6, 27.6)	1423	24.7 (23.3, 26.1)
	Never smoker	2499	59.9 (57.9, 61.8)	2617	60.4 (58.7, 62.1)	3146	61.6 (60.0, 63.2)

Table 9. Frequent Mental Distress (FMD) and current smoking status by year, Oklahoma BRFSS 2011 – 2023

Poor mental health	Smoking Status	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Current smoker	493	43.2 (39.1, 47.3)	414	42.9 (38.8, 47.0)	403	45.2 (41.1, 49.4)	348	38.9 (35.0, 42.8)	265	41.1 (36.1, 46.2)
	Former smoker	328	22.6 (19.5, 25.7)	248	20.8 (17.7, 23.9)	276	21.9 (18.8, 25.0)	301	24.7 (21.6, 27.9)	211	22.5 (18.6, 26.4)
	Never smoker	395	34.2 (30.2, 38.1)	369	36.3 (32.3, 40.3)	375	32.9 (29.0, 36.7)	381	36.3 (32.6, 40.1)	280	36.4 (31.4, 41.3)
No	Current smoker	1364	23.2 (21.7, 24.7)	1222	20.3 (19.0, 21.7)	1164	20.2 (18.9, 21.5)	1079	18.3 (17.1, 19.6)	883	19.4 (17.7, 21.0)
	Former smoker	2154	25.1 (23.8, 26.4)	1953	24.9 (23.7, 26.2)	2051	25.0 (23.7, 26.2)	2103	26.1 (24.9, 27.4)	1781	25.0 (23.5, 26.5)
	Never smoker	3774	51.7 (50.1, 53.4)	3742	54.8 (53.2, 56.3)	3865	54.9 (53.3, 56.4)	3985	55.5 (54.0, 57.0)	3333	55.6 (53.7, 57.5)

Poor mental health	Smoking Status	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Current smoker	242	36.0 (31.4, 40.7)	234	33.8 (29.5, 38.2)	215	34.9 (30.3, 39.6)	243	33.8 (29.6, 37.9)	219	33.8 (29.6, 37.9)
	Former smoker	234	24.5 (20.9, 28.2)	233	22.9 (19.6, 26.2)	176	21.7 (17.9, 25.4)	220	24.4 (20.8, 27.9)	181	24.4 (20.8, 27.9)
	Never smoker	333	39.4 (34.9, 43.9)	338	43.3 (38.8, 47.8)	276	43.4 (38.5, 48.3)	324	41.9 (37.6, 46.2)	268	41.9 (37.6, 46.2)
No	Current smoker	741	16.6 (15.2, 18.1)	744	17.4 (15.9, 18.8)	591	16.5 (14.9, 18.0)	678	15.8 (14.4, 17.1)	543	15.8 (14.4, 17.1)
	Former smoker	1643	25.0 (23.5, 26.4)	1577	24.9 (23.4, 26.3)	1237	25.6 (23.9, 27.2)	1426	25.6 (24.1, 27.1)	1080	25.6 (24.1, 27.1)
	Never smoker	3362	58.4 (56.6, 60.1)	3105	57.8 (56.0, 59.5)	2449	58.0 (56.1, 59.9)	2930	58.6 (56.9, 60.3)	2261	58.6 (56.9, 60.3)

Frequent Mental Distress = 14 or more days of poor mental health

Poor mental health	Smoking Status	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Current smoker	195	27.9 (23.9, 32.0)	187	23.5 (20.1, 27.0)	245	25.9 (22.5, 29.4)
	Former smoker	213	26.4 (22.5, 30.3)	254	27.1 (23.7, 30.5)	293	29.3 (25.8, 32.7)
	Never smoker	333	45.6 (41.2, 50.1)	415	49.3 (45.4, 53.3)	404	44.8 (40.9, 48.7)
No	Current smoker	534	14.3 (12.9, 15.7)	559	13.5 (12.3, 14.8)	622	13.6 (12.5, 14.8)
	Former smoker	1148	25.6 (23.9, 27.3)	1252	27.2 (25.6, 28.8)	1467	25.5 (24.1, 26.9)
	Never smoker	2503	60.2 (58.2, 62.1)	2536	59.3 (57.5, 61.0)	3120	60.9 (59.2, 62.5)

Frequent Mental Distress = 14 or more days of poor mental health

Table 10. Health insurance coverage and current smoking status by year, Oklahoma BRFSS 2011 – 2023

Health Insurance Status	Smoking Status	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Insured	Current smoker	1392	22.3 (20.9, 23.8)	1246	19.8 (18.6, 21.1)	1173	20.0 (18.7, 21.4)	1154	18.6 (17.4, 19.9)	955	19.8 (18.2, 21.4)
	Former smoker	2227	26.9 (25.5, 28.2)	2023	26.6 (25.3, 27.9)	2132	26.2 (24.9, 27.4)	2256	27.3 (26.0, 28.5)	1897	26.4 (24.9, 27.9)
	Never smoker	3644	50.8 (49.2, 52.4)	3615	53.6 (52.0, 55.1)	3791	53.8 (52.3, 55.3)	4003	54.1 (52.6, 55.6)	3374	53.8 (52.0, 55.7)
Uninsured	Current smoker	461	39.9 (36.0, 43.7)	387	39.3 (35.5, 43.1)	389	39.8 (36.1, 43.6)	269	36.0 (31.8, 40.2)	189	38.2 (32.8, 43.7)
	Former smoker	248	17.0 (14.3, 19.6)	176	14.8 (12.2, 17.3)	189	17.9 (15.0, 20.7)	142	17.6 (14.3, 20.8)	93	14.5 (10.9, 18.0)
	Never smoker	516	43.1 (39.3, 47.0)	483	45.9 (42.1, 49.7)	437	42.3 (38.6, 46.1)	352	46.4 (42.1, 50.8)	224	47.3 (41.6, 53.0)

Health Insurance Status	Smoking Status	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Insured	Current smoker	833	17.3 (15.9, 18.7)	829	18.2 (16.8, 19.6)	678	17.5 (16.0, 19.0)	774	17.2 (15.8, 18.5)	659	18.3 (16.7, 19.8)
	Former smoker	1794	26.1 (24.7, 27.6)	1748	26.6 (25.1, 28.0)	1335	26.0 (24.4, 27.5)	1573	26.7 (25.2, 28.1)	1182	26.1 (24.5, 27.7)
	Never smoker	3486	56.6 (54.9, 58.3)	3209	55.2 (53.5, 56.9)	2528	56.5 (54.6, 58.4)	3019	56.2 (54.5, 57.9)	2333	55.6 (53.7, 57.5)
Uninsured	Current smoker	174	33.5 (28.4, 38.5)	178	31.6 (26.8, 36.4)	159	33.2 (28.2, 38.2)	194	28.8 (24.6, 32.9)	136	24.1 (19.8, 28.3)
	Former smoker	114	17.7 (14.1, 21.3)	105	13.4 (10.4, 16.5)	99	18.1 (14.1, 22.1)	123	17.9 (14.5, 21.3)	121	21.1 (17.0, 25.3)
	Never smoker	268	48.8 (43.6, 54.0)	282	55.0 (49.8, 60.1)	245	48.7 (43.5, 54.0)	340	53.4 (48.8, 57.9)	267	54.8 (49.7, 59.9)

Health Insurance Status	Smoking Status	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Insured	Current smoker	635	15.7 (14.3, 17.1)	686	15.1 (13.8, 16.3)	790	15.3 (14.1, 16.5)
	Former smoker	1310	27.0 (25.3, 28.6)	1449	28.0 (26.5, 29.5)	1709	27.5 (26.1, 28.8)
	Never smoker	2661	57.3 (55.5, 59.2)	2775	56.9 (55.3, 58.6)	3307	57.2 (55.7, 58.8)
Uninsured	Current smoker	113	27.7 (22.4, 32.9)	81	21.0 (16.3, 25.6)	93	21.8 (17.3, 26.3)
	Former smoker	75	18.8 (14.1, 23.6)	80	20.9 (16.3, 25.6)	75	15.8 (11.9, 19.7)
	Never smoker	176	53.5 (47.4, 59.5)	208	58.1 (52.3, 63.9)	238	62.4 (57.0, 67.7)

Table 11. Disability status and current smoking status by year, Oklahoma BRFSS 2016 – 2023

Disability Status	Smoking Status	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
One or more disability	Current smoker	503	27.9 (25.3, 30.5)	510	27.9 (25.4, 30.5)	412	27.0 (24.2, 29.7)	493	25.1 (22.8, 27.5)	429	28.2 (25.4, 30.9)
	Former smoker	798	29.8 (27.4, 32.1)	861	29.2 (27.0, 31.5)	670	30.7 (28.1, 33.4)	804	31.8 (29.4, 34.1)	586	30.4 (27.7, 33.0)
	Never smoker	1193	42.3 (39.6, 44.9)	1136	42.9 (40.1, 45.6)	900	42.3 (39.4, 45.2)	1104	43.1 (40.5, 45.6)	795	41.5 (38.5, 44.4)
No disability	Current smoker	486	15.2 (13.6, 16.9)	486	16.0 (14.4, 17.6)	423	15.8 (14.1, 17.5)	461	15.3 (13.7, 16.9)	355	14.1 (12.4, 15.8)
	Former smoker	1087	22.4 (20.8, 24.0)	976	22.0 (20.4, 23.7)	762	21.8 (20.0, 23.5)	873	21.4 (19.8, 23.0)	696	22.3 (20.5, 24.1)
	Never smoker	2525	62.4 (60.4, 64.4)	2334	62.0 (60.0, 64.0)	1874	62.5 (60.3, 64.6)	2225	63.3 (61.3, 65.3)	1797	63.6 (61.4, 65.8)

Disability Status	Smoking Status	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
One or more disability	Current smoker	392	24.2 (21.7, 26.8)	422	21.3 (19.2, 23.5)	488	22.6 (20.5, 24.8)
	Former smoker	637	29.8 (27.2, 32.4)	778	33.0 (30.7, 35.3)	808	30.0 (27.8, 32.1)
	Never smoker	909	46.0 (43.1, 48.9)	1045	45.7 (43.1, 48.2)	1158	47.4 (44.9, 49.9)
No disability	Current smoker	354	12.8 (11.2, 14.3)	348	12.0 (10.6, 13.4)	390	11.3 (10.0, 12.5)
	Former smoker	750	23.8 (21.8, 25.7)	756	23.2 (21.5, 24.9)	969	23.4 (21.8, 25.0)
	Never smoker	1948	63.5 (61.3, 65.7)	1949	64.8 (62.8, 66.8)	2418	65.3 (63.5, 67.2)

Section II. Pattern of cigarette smoking (Daily, Someday, Former, and Never smokers)

Table 1. Proportion of everyday and someday smokers among Oklahoma adults, Oklahoma BRFSS 2011 – 2023

Smoking Status	2011		2012		2013		2014		2015	
	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Daily smoker	1394	19.9 (18.6, 21.2)	1214	17.0 (15.8, 18.1)	1140	17.4 (16.2, 18.5)	1030	15.1 (14.0, 16.2)	842	16.0 (14.6, 17.4)
Someday smoker	463	6.2 (5.4, 7.0)	422	6.3 (5.5, 7.1)	427	6.3 (5.6, 7.1)	397	6.0 (5.2, 6.7)	306	6.2 (5.2, 7.2)
Former smoker	2482	24.7 (23.5, 25.9)	2201	24.4 (23.2, 25.5)	2327	24.5 (23.4, 25.7)	2404	25.9 (24.8, 27.1)	1992	24.7 (23.3, 26.1)
Never smoker	4169	49.2 (47.6, 50.7)	4111	52.4 (50.9, 53.8)	4240	51.8 (50.3, 53.2)	4366	53.0 (51.6, 54.4)	3613	53.4 (51.4, 54.9)

Smoking Status	2016		2017		2018		2019		2020	
	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Daily smoker	729	14.0 (12.8, 15.3)	702	13.3 (12.1, 14.4)	595	13.7 (12.4, 15.0)	683	13.2 (12.0, 14.3)	608	14.0 (12.8, 15.3)
Someday smoker	281	5.6 (4.7, 6.4)	306	6.9 (5.9, 7.8)	244	6.0 (5.1, 6.9)	286	5.7 (4.9, 6.5)	190	5.1 (4.1, 6.0)
Former smoker	1914	24.8 (23.5, 26.2)	1856	24.5 (23.2, 25.8)	1439	24.8 (23.4, 26.3)	1699	25.3 (24.0, 26.6)	1307	25.2 (23.7, 26.7)
Never smoker	3766	55.6 (53.9, 57.2)	3507	55.3 (53.7, 56.9)	2782	55.5 (53.7, 57.3)	3368	55.8 (54.3, 57.4)	2617	55.7 (53.9, 57.5)

Smoking Status	2021		2022		2023	
	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Daily smoker	559	12.2 (11.0, 13.3)	577	11.6 (10.5, 12.6)	642	11.1 (10.1, 12.1)
Someday smoker	208	4.8 (4.0, 5.6)	201	4.0 (3.4, 4.7)	258	4.7 (4.1, 5.4)
Former smoker	1408	25.7 (24.2, 27.2)	1555	27.2 (25.8, 28.6)	1820	26.1 (24.8, 27.4)
Never smoker	2918	57.4 (55.6, 59.1)	3037	57.2 (55.6, 58.8)	3637	58.1 (56.6, 59.6)

Table 2. Proportion of everyday and someday smokers among Oklahoma adults by sex, Oklahoma BRFSS 2016 – 2023

Sex	Smoking Status	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Male	Daily smoker	584	21.8 (19.7, 23.9)	500	17.1 (15.4, 18.9)	508	18.7 (16.9, 20.5)	480	16.2 (14.6, 17.9)	370	17.1 (14.9, 19.3)
	Someday smoker	169	6.3 (4.9, 7.6)	185	7.3 (6.0, 8.6)	181	7.0 (5.8, 8.2)	168	6.8 (5.6, 8.1)	135	6.8 (5.2, 8.4)
	Former smoker	1166	28.8 (26.7, 30.8)	1150	29.6 (27.7, 31.5)	1114	27.6 (25.8, 29.5)	1158	29.6 (27.7, 31.5)	960	27.6 (25.4, 29.8)
	Never smoker	1244	43.2 (40.7, 45.6)	1371	46.0 (43.7, 48.3)	1428	46.6 (44.4, 48.8)	1530	47.4 (45.2, 49.5)	1209	48.5 (45.7, 51.2)
Female	Daily smoker	810	18.1 (16.6, 19.6)	714	16.8 (15.3, 18.2)	632	16.1 (14.6, 17.6)	550	14.0 (12.5, 15.4)	472	14.9 (13.2, 16.6)
	Someday smoker	294	6.2 (5.2, 7.2)	237	5.4 (4.5, 6.3)	246	5.6 (4.7, 6.6)	229	5.1 (4.3, 6.0)	171	5.5 (4.4, 6.7)
	Former smoker	1316	20.9 (19.5, 22.2)	1051	19.4 (18.0, 20.7)	1213	21.6 (20.1, 23.0)	1246	22.4 (21.0, 23.9)	1032	21.9 (20.1, 23.7)
	Never smoker	2925	54.9 (53.0, 56.7)	2740	58.4 (56.6, 60.3)	2812	56.7 (54.9, 58.6)	2836	58.5 (56.6, 60.3)	2404	57.7 (55.4, 59.9)
Sex	Smoking Status	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Male	Daily smoker	336	15.4 (13.4, 17.4)	307	13.7 (12.0, 15.4)	293	15.3 (13.3, 17.3)	320	14.2 (12.4, 15.9)	282	14.8 (12.8, 16.7)
	Someday smoker	109	6.0 (4.6, 7.4)	140	7.5 (6.0, 9.0)	114	6.9 (5.5, 8.4)	145	6.9 (5.6, 8.2)	86	5.9 (4.3, 7.5)
	Former smoker	944	28.2 (26.1, 30.3)	916	27.9 (25.8, 29.9)	718	28.5 (26.2, 30.8)	883	29.1 (27.1, 31.2)	644	28.3 (25.9, 30.6)
	Never smoker	1316	50.4 (47.8, 52.9)	1232	50.9 (48.4, 53.4)	1001	49.3 (46.6, 52.0)	1260	49.8 (47.4, 52.2)	1013	51.1 (48.4, 53.8)
Female	Daily smoker	393	12.8 (11.2, 14.3)	395	12.9 (11.4, 14.4)	302	12.1 (10.5, 13.7)	363	12.2 (10.7, 13.7)	326	13.4 (11.7, 15.0)
	Someday smoker	172	5.1 (4.2, 6.1)	166	6.3 (5.1, 7.5)	130	5.1 (4.0, 6.2)	141	4.6 (3.7, 5.5)	104	4.3 (3.3, 5.2)
	Former smoker	970	21.6 (20.0, 23.2)	940	21.4 (19.7, 23.0)	720	21.3 (19.5, 23.1)	816	21.6 (19.9, 23.3)	663	22.3 (20.4, 24.2)
	Never smoker	2450	60.5 (58.4, 62.6)	2273	59.4 (57.3, 61.6)	1781	61.5 (59.2, 63.8)	2108	61.6 (59.5, 63.7)	1604	60.1 (57.7, 62.4)

Sex	Smoking Status	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Male	Daily smoker	247	11.6 (10.0, 13.3)	288	12.6 (11.0, 14.2)	286	11.3 (9.9, 12.8)
	Someday smoker	93	5.4 (4.1, 6.8)	87	4.1 (3.1, 5.1)	117	5.1 (4.0, 6.1)
	Former smoker	718	29.0 (26.6, 31.4)	780	29.7 (27.6, 31.8)	948	29.9 (27.9, 31.9)
	Never smoker	1183	53.9 (51.3, 56.6)	1257	53.7 (51.3, 56.0)	1483	53.7 (51.5, 56.0)
Female	Daily smoker	312	12.7 (11.1, 14.3)	289	10.7 (9.3, 12.0)	356	10.9 (9.6, 12.2)
	Someday smoker	115	4.1 (3.2, 5.0)	114	3.9 (3.1, 4.8)	141	4.4 (3.6, 5.2)
	Former smoker	690	22.6 (20.6, 24.6)	775	24.7 (22.9, 26.5)	872	22.5 (20.9, 24.1)
	Never smoker	1735	60.6 (58.3, 62.9)	1780	60.7 (58.6, 62.8)	2154	62.3 (60.3, 64.2)

Table 3. Three-year rolling weighted percentages of everyday and someday smokers among Oklahoma adults by age, Oklahoma BRFSS 2011 – 2023

Age Group	Smoking Status	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
18-24	Daily smoker	200	18.1 (15.5, 20.8)	195	17.2 (14.6, 19.7)	155	15.7 (13.0, 18.3)	125	14.4 (11.7, 17.1)	107	14.0 (11.2, 16.8)
	Someday smoker	96	8.6 (6.7, 10.6)	99	8.9 (7.0, 10.9)	81	8.6 (6.5, 10.7)	70	8.1 (5.9, 10.3)	55	7.4 (5.2, 9.6)
	Former smoker	107	8.0 (6.4, 9.6)	101	7.6 (6.0, 9.2)	87	8.0 (6.1, 9.9)	70	7.0 (5.1, 8.8)	61	6.5 (4.6, 8.3)
	Never smoker	735	65.2 (62.0, 68.5)	795	66.3 (63.2, 69.4)	727	67.7 (64.4, 71.1)	708	70.5 (67.1, 74.0)	635	72.1 (68.6, 75.6)
25-34	Daily smoker	513	21.9 (20.0, 23.9)	477	21.0 (19.1, 22.9)	401	20.3 (18.2, 22.3)	352	19.2 (17.1, 21.3)	303	17.1 (15.0, 19.1)
	Someday smoker	221	8.8 (7.5, 10.1)	193	7.9 (6.7, 9.1)	171	8.3 (6.9, 9.8)	148	7.8 (6.4, 9.3)	163	9.9 (8.2, 11.6)
	Former smoker	477	18.3 (16.6, 20.1)	475	19.0 (17.3, 20.8)	377	17.5 (15.7, 19.3)	346	17.8 (15.9, 19.7)	311	16.4 (14.5, 18.3)
	Never smoker	1406	50.9 (48.7, 53.2)	1424	52.1 (49.8, 54.3)	1275	53.9 (51.4, 56.3)	1182	55.2 (52.7, 57.8)	1123	56.6 (54.0, 59.2)
35-44	Daily smoker	568	20.8 (19.0, 22.6)	496	18.4 (16.7, 20.1)	434	18.2 (16.4, 20.1)	346	16.6 (14.6, 18.5)	309	16.4 (14.4, 18.4)
	Someday smoker	190	6.4 (5.3, 7.4)	192	6.6 (5.5, 7.6)	159	6.1 (5.0, 7.2)	132	6.0 (4.8, 7.2)	117	6.2 (4.9, 7.4)
	Former smoker	531	18.9 (17.2, 20.7)	564	20.2 (18.5, 21.9)	525	22.0 (20.0, 23.9)	493	22.4 (20.3, 24.5)	437	22.2 (20.0, 24.4)
	Never smoker	1682	53.8 (51.7, 56.0)	1670	54.9 (52.7, 57.0)	1490	53.7 (51.4, 56.1)	1357	55.0 (52.5, 57.5)	1182	55.2 (52.6, 57.8)
45-54	Daily smoker	882	22.9 (21.4, 24.5)	729	19.3 (17.9, 20.8)	613	19.3 (17.6, 20.9)	500	18.0 (16.3, 19.7)	423	17.6 (15.8, 19.5)
	Someday smoker	276	6.6 (5.7, 7.6)	253	6.9 (5.9, 7.8)	214	6.6 (5.5, 7.7)	176	6.2 (5.1, 7.3)	150	6.1 (4.9, 7.3)
	Former smoker	931	23.4 (21.8, 24.9)	888	23.7 (22.1, 25.2)	821	24.7 (23.0, 26.4)	745	25.2 (23.4, 27.1)	615	24.6 (22.6, 26.5)
	Never smoker	1965	47.1 (45.2, 48.9)	1986	50.1 (48.3, 52.0)	1827	49.4 (47.4, 51.4)	1716	50.6 (48.5, 52.7)	1517	51.7 (49.4, 53.9)
55-64	Daily smoker	890	17.2 (15.9, 18.5)	840	16.0 (14.8, 17.2)	769	15.8 (14.6, 17.1)	671	15.0 (13.7, 16.3)	581	15.1 (13.7, 16.4)
	Someday smoker	291	5.5 (4.7, 6.3)	289	5.5 (4.8, 6.3)	275	5.6 (4.8, 6.4)	236	5.3 (4.5, 6.1)	204	5.3 (4.4, 6.2)
	Former smoker	1614	31.5 (30.0, 33.1)	1582	31.3 (29.9, 32.8)	1527	31.2 (29.6, 32.7)	1435	31.3 (29.7, 32.9)	1277	31.4 (29.7, 33.1)

Age Group	Smoking Status	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
65 or more	Never smoker	2537	45.8 (44.2, 47.4)	2612	47.1 (45.5, 48.7)	2524	47.4 (45.7, 49.0)	2403	48.5 (46.7, 50.2)	2153	48.2 (46.4, 50.1)
	Daily smoker	695	8.2 (7.5, 8.9)	647	8.1 (7.4, 8.8)	640	8.6 (7.8, 9.5)	607	8.2 (7.4, 9.0)	550	7.8 (7.0, 8.6)
	Someday smoker	238	2.5 (2.1, 2.9)	220	2.5 (2.1, 2.9)	230	2.7 (2.3, 3.2)	222	2.9 (2.4, 3.4)	204	3.0 (2.4, 3.5)
	Former smoker	3350	42.1 (40.8, 43.4)	3322	42.1 (40.8, 43.4)	3386	41.1 (39.8, 42.4)	3221	40.8 (39.5, 42.1)	3061	40.4 (39.1, 41.7)
	Never smoker	4195	47.2 (45.9, 48.5)	4227	47.3 (46.0, 48.6)	4373	47.5 (46.2, 48.8)	4376	48.0 (46.7, 49.3)	4276	48.9 (47.5, 50.2)

Age Group	Smoking Status	2016-18		2017-19		2018-20		2019-21		2020-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
18-24	Daily smoker	96	12.3 (9.7, 14.9)	94	11.6 (9.1, 14.1)	71	9.1 (6.8, 11.3)	66	7.6 (5.7, 9.5)	50	5.7 (4.0, 7.4)
	Someday smoker	58	7.5 (5.4, 9.6)	56	6.7 (4.8, 8.6)	50	5.9 (4.2, 7.7)	40	4.3 (2.9, 5.8)	39	4.2 (2.8, 5.7)
	Former smoker	57	5.6 (4.0, 7.2)	64	6.6 (4.8, 8.4)	71	7.5 (5.6, 9.5)	79	8.5 (6.5, 10.6)	86	8.6 (6.7, 10.6)
	Never smoker	661	74.6 (71.3, 78.0)	674	75.1 (71.9, 78.4)	691	77.5 (74.3, 80.6)	752	79.5 (76.6, 82.4)	789	81.4 (78.6, 84.2)
25-34	Daily smoker	270	15.5 (13.5, 17.4)	249	14.1 (12.3, 15.9)	224	14.4 (12.4, 16.3)	212	13.0 (11.1, 14.8)	193	12.0 (10.2, 13.8)
	Someday smoker	160	10.0 (8.3, 11.7)	155	9.9 (8.2, 11.7)	123	8.1 (6.5, 9.7)	106	7.0 (5.4, 8.5)	95	6.3 (4.8, 7.8)
	Former smoker	315	16.9 (14.9, 18.8)	312	16.5 (14.6, 18.4)	309	17.4 (15.4, 19.5)	330	19.6 (17.4, 21.8)	339	20.7 (18.4, 22.9)
	Never smoker	1107	57.7 (55.0, 60.3)	1159	59.5 (56.9, 62.0)	1088	60.1 (57.5, 62.8)	1120	60.5 (57.8, 63.1)	1120	61.0 (58.3, 63.7)
35-44	Daily smoker	299	15.5 (13.6, 17.3)	327	15.7 (13.9, 17.5)	314	15.7 (13.9, 17.5)	330	16.8 (14.9, 18.7)	312	16.3 (14.5, 18.2)
	Someday smoker	109	5.8 (4.6, 7.0)	125	5.9 (4.7, 7.0)	119	6.1 (4.9, 7.4)	127	6.2 (4.9, 7.4)	109	5.7 (4.4, 6.9)
	Former smoker	454	23.4 (21.2, 25.6)	487	24.0 (21.8, 26.1)	488	25.6 (23.3, 27.8)	518	25.0 (22.8, 27.1)	543	26.4 (24.2, 28.6)

Age Group	Smoking Status	2016-18		2017-19		2018-20		2019-21		2020-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
45-54	Never smoker	1118	55.3 (52.7, 57.9)	1150	54.5 (52.0, 56.9)	1077	52.6 (50.1, 55.2)	1159	52.1 (49.6, 54.6)	1123	51.6 (49.1, 54.1)
	Daily smoker	385	17.3 (15.5, 19.1)	373	16.9 (15.1, 18.7)	385	18.3 (16.4, 20.2)	358	16.8 (15.0, 18.7)	328	15.9 (14.1, 17.8)
	Someday smoker	150	6.3 (5.2, 7.5)	148	6.6 (5.4, 7.7)	117	5.4 (4.3, 6.5)	111	5.1 (4.0, 6.2)	96	4.3 (3.3, 5.3)
	Former smoker	548	23.9 (21.8, 25.9)	532	23.5 (21.4, 25.5)	511	23.0 (20.9, 25.0)	532	23.7 (21.7, 25.8)	552	25.4 (23.3, 27.5)
	Never smoker	1390	52.5 (50.2, 54.8)	1323	53.1 (50.7, 55.4)	1245	53.3 (50.9, 55.7)	1275	54.3 (51.9, 56.8)	1242	54.4 (52.0, 56.8)
55-64	Daily smoker	500	15.3 (13.8, 16.8)	468	15.1 (13.6, 16.6)	440	16.3 (14.6, 18.0)	434	16.3 (14.7, 18.0)	429	17.0 (15.3, 18.7)
	Someday smoker	170	5.2 (4.3, 6.1)	170	5.8 (4.8, 6.8)	143	5.5 (4.4, 6.6)	140	5.6 (4.5, 6.7)	119	4.5 (3.6, 5.4)
	Former smoker	1107	30.9 (29.1, 32.7)	1017	31.1 (29.2, 32.9)	851	29.9 (27.9, 31.9)	813	29.6 (27.5, 31.6)	763	29.4 (27.3, 31.4)
	Never smoker	1932	48.6 (46.6, 50.5)	1733	48.1 (46.1, 50.0)	1503	48.3 (46.2, 50.4)	1421	48.5 (46.3, 50.7)	1373	49.1 (46.9, 51.4)
65 or more	Daily smoker	476	7.5 (6.8, 8.3)	469	8.1 (7.3, 9.0)	452	8.8 (7.9, 9.8)	450	8.8 (7.9, 9.8)	432	8.9 (7.9, 9.9)
	Someday smoker	184	2.8 (2.3, 3.3)	182	3.0 (2.5, 3.5)	168	3.1 (2.6, 3.7)	160	3.3 (2.7, 3.9)	141	3.0 (2.4, 3.6)
	Former smoker	2728	40.5 (39.1, 41.9)	2582	40.4 (38.9, 41.8)	2215	39.7 (38.2, 41.2)	2142	38.6 (37.0, 40.2)	1987	38.2 (36.6, 39.8)
	Never smoker	3847	49.2 (47.8, 50.6)	3618	48.5 (47.1, 49.9)	3163	48.3 (46.8, 49.9)	3176	49.3 (47.7, 50.9)	2925	50.0 (48.3, 51.6)

Age Group	Smoking Status	2021-23	
		N	Weighted% (95%CI)
18-24	Daily smoker	54	5.8 (4.1, 7.4)
	Someday smoker	36	3.6 (2.3, 4.8)
	Former smoker	86	8.3 (6.4, 10.2)
	Never smoker	867	82.3 (79.7, 84.9)
25-34	Daily smoker	172	9.7 (8.1, 11.2)
	Someday smoker	108	6.0 (4.7, 7.2)
	Former smoker	414	22.9 (20.7, 25.2)
	Never smoker	1262	61.4 (58.9, 63.9)
35-44	Daily smoker	334	15.2 (13.5, 16.9)
	Someday smoker	127	5.3 (4.2, 6.3)
	Former smoker	643	26.3 (24.3, 28.3)
	Never smoker	1358	53.2 (50.9, 55.5)
45-54	Daily smoker	313	13.9 (12.3, 15.6)
	Someday smoker	127	5.3 (4.2, 6.3)
	Former smoker	636	26.1 (24.1, 28.2)
	Never smoker	1393	54.7 (52.4, 57.0)
55-64	Daily smoker	447	16.6 (15.0, 18.2)
	Someday smoker	129	4.6 (3.7, 5.5)
	Former smoker	825	28.2 (26.3, 30.1)
	Never smoker	1555	50.6 (48.4, 52.7)

Age Group	Smoking Status	2021-23	
		N	Weighted% (95%CI)
65 or more	Daily smoker	458	8.8 (7.9, 9.8)
	Someday smoker	140	2.8 (2.3, 3.4)
	Former smoker	2179	38.0 (36.5, 39.6)
	Never smoker	3157	50.3 (48.8, 51.9)

Table 4. Three-year rolling weighted percentages of everyday and someday smokers among Oklahoma adults by race/ethnicity, Oklahoma BRFSS 2011 – 2023

Race/Ethnicity	Smoking Status	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
NH-White	Daily smoker	2711	18.2 (17.4, 19.0)	2475	16.6 (15.8, 17.3)	2191	16.1 (15.3, 16.9)	1943	15.2 (14.4, 16.1)	1707	14.7 (13.9, 15.6)
	Someday smoker	810	5.5 (5.0, 6.0)	765	5.2 (4.7, 5.6)	716	5.2 (4.7, 5.8)	642	5.1 (4.6, 5.6)	573	5.2 (4.6, 5.7)
	Former smoker	5464	26.3 (25.5, 27.1)	5456	27.4 (26.6, 28.2)	5340	27.8 (27.0, 28.7)	5038	27.8 (26.9, 28.7)	4604	27.1 (26.2, 28.1)
	Never smoker	9320	50.1 (49.1, 51.0)	9447	50.9 (50.0, 51.8)	9118	50.8 (49.8, 51.8)	8916	51.9 (50.8, 52.9)	8352	53.0 (51.9, 54.1)
NH-Black	Daily smoker	245	18.0 (15.1, 20.8)	188	15.4 (12.9, 18.0)	180	16.0 (13.1, 18.9)	141	14.6 (11.6, 17.7)	106	12.0 (9.0, 15.0)
	Someday smoker	142	8.7 (6.8, 10.6)	130	9.5 (7.5, 11.6)	115	9.5 (7.3, 11.7)	86	8.3 (6.2, 10.5)	83	8.7 (6.4, 11.0)
	Former smoker	307	17.4 (14.8, 20.0)	295	17.1 (14.8, 19.4)	318	18.8 (16.2, 21.3)	300	19.4 (16.7, 22.1)	271	19.0 (16.1, 21.8)
	Never smoker	777	55.9 (52.3, 59.5)	781	57.9 (54.6, 61.3)	751	55.8 (52.2, 59.3)	667	57.7 (53.9, 61.5)	624	60.3 (56.4, 64.3)
NH-AI/AN	Daily smoker	306	23.6 (20.5, 26.7)	278	22.0 (19.0, 24.9)	278	23.6 (20.5, 26.7)	239	20.2 (17.3, 23.1)	210	18.8 (15.9, 21.8)
	Someday smoker	145	10.0 (8.0, 11.9)	139	9.2 (7.4, 11.0)	127	8.7 (6.9, 10.5)	110	7.6 (5.8, 9.5)	111	10.0 (7.7, 12.3)
	Former smoker	428	24.1 (21.5, 26.7)	429	23.3 (20.8, 25.7)	388	20.9 (18.5, 23.4)	390	23.3 (20.6, 26.0)	333	22.1 (19.4, 24.9)
	Never smoker	622	42.3 (39.0, 45.6)	694	45.6 (42.3, 48.8)	707	46.8 (43.4, 50.2)	697	48.8 (45.4, 52.3)	623	49.0 (45.4, 52.6)
NH-Other	Daily smoker	343	18.9 (16.3, 21.6)	330	19.5 (16.6, 22.5)	257	17.8 (14.7, 20.9)	186	15.9 (12.8, 19.1)	153	15.2 (12.2, 18.2)
	Someday smoker	112	5.4 (4.0, 6.9)	109	7.7 (5.5, 9.8)	93	8.5 (5.9, 11.1)	82	8.1 (5.5, 10.6)	57	6.6 (4.1, 9.0)
	Former smoker	540	21.4 (18.8, 24.0)	488	18.8 (16.4, 21.2)	421	17.7 (15.2, 20.2)	360	17.6 (15.0, 20.1)	332	19.1 (16.4, 21.8)
	Never smoker	935	54.2 (50.8, 57.7)	890	54.0 (50.5, 57.6)	809	56.0 (52.1, 59.9)	722	58.5 (54.4, 62.5)	643	59.1 (55.1, 63.2)
Hispanic	Daily smoker	131	11.5 (9.2, 13.8)	103	9.5 (7.4, 11.6)	93	9.4 (7.0, 11.7)	72	8.7 (6.3, 11.2)	68	8.7 (6.2, 11.2)
	Someday smoker	98	8.4 (6.4, 10.4)	97	8.7 (6.7, 10.6)	70	7.1 (4.9, 9.2)	56	7.4 (5.0, 9.7)	63	9.5 (6.7, 12.3)
	Former smoker	244	17.9	234	16.7	214	15.6	181	15.1	174	16.1

Race/Ethnicity	Smoking Status	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
			(15.3, 20.5)		(14.3, 19.2)		(13.1, 18.1)		(12.5, 17.6)		(13.3, 18.9)
	Never smoker	810	62.2 (58.8, 65.7)	835	65.1 (61.8, 68.3)	762	68.0 (64.5, 71.5)	664	68.8 (65.1, 72.5)	551	65.7 (61.7, 69.7)

Race/Ethnicity	Smoking Status	2016-18		2017-19		2018-20		2019-21		2020-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
NH-White	Daily smoker	1523	14.2 (13.3, 15.0)	1459	13.9 (13.1, 14.7)	1368	14.2 (13.3, 15.0)	1329	13.7 (12.8, 14.5)	1255	13.2 (12.4, 14.0)
	Someday smoker	529	5.2 (4.6, 5.7)	510	5.1 (4.6, 5.7)	437	4.8 (4.2, 5.4)	408	4.3 (3.8, 4.8)	368	3.9 (3.4, 4.4)
	Former smoker	4142	27.0 (26.1, 27.9)	3935	27.2 (26.2, 28.1)	3470	27.6 (26.6, 28.6)	3399	27.9 (26.9, 28.9)	3253	28.6 (27.5, 29.6)
	Never smoker	7660	53.7 (52.6, 54.8)	7225	53.8 (52.7, 54.9)	6432	53.4 (52.3, 54.6)	6446	54.2 (53.0, 55.3)	6120	54.3 (53.2, 55.5)
NH-Black	Daily smoker	90	10.6 (8.0, 13.2)	91	10.3 (8.0, 12.6)	93	12.6 (9.8, 15.4)	88	12.6 (9.8, 15.4)	81	12.3 (9.4, 15.1)
	Someday smoker	77	8.8 (6.5, 11.1)	74	8.6 (6.3, 10.8)	58	7.3 (5.1, 9.4)	51	6.5 (4.6, 8.5)	44	5.9 (4.0, 7.9)
	Former smoker	227	18.1 (15.2, 21.0)	204	18.3 (15.3, 21.2)	175	17.3 (14.3, 20.3)	174	17.6 (14.6, 20.6)	170	18.5 (15.4, 21.6)
	Never smoker	569	62.4 (58.5, 66.3)	552	62.9 (59.1, 66.7)	495	62.8 (58.8, 66.8)	505	63.3 (59.3, 67.2)	479	63.3 (59.3, 67.3)
NH-AI/AN	Daily smoker	167	16.2 (13.4, 19.0)	180	18.1 (15.2, 21.0)	183	17.5 (14.7, 20.4)	194	17.5 (14.8, 20.2)	178	14.8 (12.3, 17.2)
	Someday smoker	89	9.1 (6.8, 11.4)	99	9.3 (7.1, 11.5)	94	8.9 (6.5, 11.2)	110	10.3 (7.9, 12.7)	91	9.4 (7.1, 11.8)
	Former smoker	336	25.3 (22.3, 28.3)	330	24.9 (21.9, 27.9)	335	26.3 (23.2, 29.3)	322	25.1 (22.1, 28.1)	323	26.2 (23.2, 29.1)
	Never smoker	559	49.5 (45.8, 53.1)	546	47.7 (44.1, 51.3)	544	47.4 (43.7, 51.0)	567	47.1 (43.6, 50.6)	567	49.7 (46.1, 53.2)
NH-Other	Daily smoker	158	16.7 (13.6, 19.8)	164	15.0 (12.1, 17.8)	155	15.1 (12.0, 18.2)	140	12.4 (9.8, 15.1)	129	13.2 (10.5, 15.9)
	Someday smoker	65	6.8 (4.8, 8.9)	72	7.7 (5.5, 9.8)	67	7.2 (5.2, 9.2)	50	5.8 (3.9, 7.8)	47	5.4 (3.5, 7.3)
	Former smoker	301	18.7 (16.1, 21.4)	293	20.6 (17.8, 23.5)	253	22.0 (18.8, 25.1)	261	22.6 (19.3, 25.8)	262	23.4 (20.3, 26.6)
	Never smoker	611	57.7 (53.8, 61.7)	595	56.7 (52.9, 60.6)	529	55.7 (51.7, 59.8)	559	59.2 (55.2, 63.2)	560	58.0 (54.1, 61.9)

Race/Ethnicity	Smoking Status	2016-18		2017-19		2018-20		2019-21		2020-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Hispanic	Daily smoker	61	6.9 (4.9, 8.9)	62	6.3 (4.5, 8.0)	69	6.1 (4.5, 7.7)	75	6.6 (4.9, 8.4)	73	6.4 (4.7, 8.1)
	Someday smoker	68	9.4 (6.9, 12.0)	74	9.0 (6.5, 11.4)	56	5.9 (4.1, 7.7)	53	5.4 (3.6, 7.2)	42	4.1 (2.5, 5.6)
	Former smoker	164	16.3 (13.4, 19.2)	176	15.6 (12.9, 18.3)	164	13.9 (11.4, 16.4)	180	14.9 (12.3, 17.5)	184	15.9 (13.3, 18.6)
	Never smoker	561	67.3 (63.5, 71.2)	635	69.1 (65.5, 72.7)	673	74.1 (70.9, 77.3)	708	73.1 (69.8, 76.3)	716	73.5 (70.3, 76.7)

Race/Ethnicity	Smoking Status	2021-23	
		N	Weighted% (95%CI)
NH-White	Daily smoker	1282	12.2 (11.4, 12.9)
	Someday smoker	405	3.7 (3.3, 4.2)
	Former smoker	3620	28.8 (27.8, 29.8)
	Never smoker	6786	55.3 (54.2, 56.4)
NH-Black	Daily smoker	79	10.0 (7.6, 12.5)
	Someday smoker	46	6.0 (4.1, 7.9)
	Former smoker	183	19.5 (16.5, 22.6)
	Never smoker	521	64.5 (60.7, 68.3)
NH-AI/AN	Daily smoker	175	15.5 (13.0, 18.0)
	Someday smoker	88	7.4 (5.7, 9.2)
	Former smoker	353	27.0 (24.0, 29.9)
	Never smoker	618	50.1 (46.7, 53.4)
NH-Other	Daily smoker	132	11.6 (9.3, 13.9)

Race/Ethnicity	Smoking Status	2021-23	
		N	Weighted% (95%CI)
	Someday smoker	60	5.8 (4.1, 7.6)
	Former smoker	318	23.8 (21.0, 26.6)
	Never smoker	685	58.8 (55.3, 62.2)
	Daily smoker	77	6.2 (4.6, 7.8)
Hispanic	Someday smoker	62	5.2 (3.7, 6.7)
	Former smoker	221	16.8 (14.4, 19.2)
	Never smoker	822	71.8 (68.8, 74.8)

Table 5. Three-year rolling weighted percentages of everyday and someday smokers among Oklahoma adults by education, Oklahoma BRFSS 2011 – 2023

Education	Smoking Status	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than high school	Daily smoker	728	30.1 (27.7, 32.5)	614	27.3 (24.9, 29.7)	497	27.0 (24.3, 29.7)	385	26.7 (23.7, 29.6)	318	25.5 (22.4, 28.5)
	Someday smoker	221	9.9 (8.2, 11.6)	210	10.1 (8.4, 11.8)	174	10.3 (8.3, 12.3)	135	9.6 (7.5, 11.7)	105	10.3 (7.9, 12.7)
	Former smoker	800	23.1 (21.1, 25.0)	740	23.7 (21.7, 25.7)	661	23.7 (21.5, 25.9)	570	24.8 (22.4, 27.3)	489	24.9 (22.2, 27.5)
	Never smoker	1119	36.9 (34.4, 39.4)	1066	38.9 (36.4, 41.4)	891	39.0 (36.2, 41.8)	734	38.9 (35.9, 41.9)	579	39.3 (35.9, 42.7)
High school	Daily smoker	1482	21.4 (20.2, 22.7)	1349	19.9 (18.7, 21.1)	1185	19.4 (18.1, 20.6)	1016	17.8 (16.6, 19.1)	892	17.5 (16.1, 18.8)
	Someday smoker	450	6.2 (5.5, 7.0)	434	6.4 (5.6, 7.1)	386	6.3 (5.6, 7.1)	353	6.3 (5.5, 7.2)	300	6.3 (5.4, 7.1)
	Former smoker	2240	24.6 (23.4, 25.8)	2255	25.4 (24.2, 26.5)	2206	26.0 (24.7, 27.3)	2076	26.2 (24.8, 27.5)	1808	25.0 (23.6, 26.4)
	Never smoker	3588	47.7 (46.2, 49.3)	3590	48.4 (46.9, 49.8)	3440	48.3 (46.7, 49.9)	3273	49.7 (48.0, 51.3)	3011	51.3 (49.6, 53.0)
Some college	Daily smoker	1082	16.8 (15.6, 17.9)	996	15.3 (14.2, 16.4)	933	15.1 (14.0, 16.2)	854	13.8 (12.7, 14.9)	747	12.9 (11.8, 14.0)
	Someday smoker	415	6.5 (5.7, 7.3)	392	6.3 (5.5, 7.0)	384	6.2 (5.4, 7.0)	328	5.8 (5.0, 6.6)	339	6.6 (5.7, 7.5)
	Former smoker	2089	25.7 (24.4, 26.9)	2052	25.8 (24.6, 27.0)	1996	25.7 (24.4, 27.0)	1942	26.1 (24.7, 27.4)	1860	26.0 (24.6, 27.4)
	Never smoker	3431	51.0 (49.5, 52.6)	3489	52.6 (51.1, 54.1)	3378	52.9 (51.4, 54.5)	3337	54.4 (52.8, 56.0)	3238	54.4 (52.8, 56.1)
College graduate	Daily smoker	453	6.1 (5.4, 6.8)	423	5.4 (4.7, 6.0)	394	5.5 (4.8, 6.2)	343	5.2 (4.5, 6.0)	312	5.4 (4.6, 6.1)
	Someday smoker	225	3.4 (2.8, 4.0)	210	3.1 (2.6, 3.7)	186	3.0 (2.4, 3.6)	168	3.1 (2.5, 3.7)	149	3.0 (2.4, 3.6)
	Former smoker	1871	23.7 (22.5, 25.0)	1876	23.9 (22.7, 25.1)	1851	23.5 (22.3, 24.7)	1712	22.5 (21.3, 23.8)	1592	22.1 (20.8, 23.3)
	Never smoker	4364	66.7 (65.4, 68.1)	4548	67.6 (66.3, 68.9)	4483	68.0 (66.6, 69.4)	4371	69.1 (67.7, 70.6)	4032	69.6 (68.1, 71.0)

Education	Smoking Status	2016-18		2017-19		2018-20		2019-21		2020-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than high school	Daily smoker	294	24.5 (21.5, 27.4)	306	24.3 (21.4, 27.2)	298	25.5 (22.5, 28.4)	292	25.7 (22.7, 28.7)	258	23.7 (20.8, 26.7)
	Someday smoker	94	9.4 (7.2, 11.5)	91	9.2 (7.1, 11.4)	79	8.1 (6.0, 10.2)	70	7.5 (5.4, 9.6)	65	6.9 (4.8, 9.0)
	Former smoker	429	25.6 (22.8, 28.4)	401	24.9 (22.2, 27.7)	347	24.4 (21.6, 27.2)	321	23.7 (20.8, 26.5)	294	24.5 (21.6, 27.5)
	Never smoker	521	40.5 (37.1, 44.0)	530	41.5 (38.2, 44.9)	503	42.0 (38.6, 45.4)	496	43.1 (39.7, 46.6)	471	44.9 (41.3, 48.4)
High school	Daily smoker	776	16.4 (15.1, 17.7)	776	16.5 (15.2, 17.8)	714	16.3 (15.0, 17.6)	696	15.9 (14.6, 17.2)	616	14.9 (13.6, 16.2)
	Someday smoker	286	6.7 (5.7, 7.6)	288	6.8 (5.9, 7.7)	254	6.4 (5.5, 7.4)	224	5.8 (4.9, 6.6)	183	5.2 (4.3, 6.1)
	Former smoker	1573	24.7 (23.3, 26.1)	1513	25.7 (24.2, 27.1)	1364	26.6 (25.0, 28.1)	1306	26.6 (25.1, 28.2)	1229	27.2 (25.6, 28.8)
	Never smoker	2681	52.3 (50.5, 54.0)	2490	51.0 (49.2, 52.8)	2191	50.7 (48.9, 52.6)	2187	51.7 (49.9, 53.5)	2058	52.7 (50.9, 54.6)
Some college	Daily smoker	675	12.6 (11.5, 13.7)	623	12.1 (11.0, 13.2)	622	12.8 (11.7, 13.9)	615	12.0 (10.9, 13.0)	624	12.2 (11.1, 13.3)
	Someday smoker	298	6.3 (5.5, 7.2)	307	6.6 (5.8, 7.5)	253	5.8 (5.0, 6.6)	269	5.9 (5.0, 6.7)	242	5.1 (4.3, 5.8)
	Former smoker	1733	26.6 (25.1, 28.0)	1653	26.4 (25.0, 27.8)	1473	26.6 (25.1, 28.0)	1497	27.4 (25.9, 28.9)	1475	28.3 (26.7, 29.8)
	Never smoker	3013	54.5 (52.9, 56.2)	2841	54.9 (53.2, 56.5)	2546	54.8 (53.1, 56.5)	2556	54.8 (53.1, 56.5)	2500	54.5 (52.8, 56.2)
College graduate	Daily smoker	276	4.9 (4.2, 5.7)	271	4.6 (3.9, 5.3)	249	4.6 (3.9, 5.4)	245	4.4 (3.7, 5.0)	245	4.4 (3.7, 5.1)
	Someday smoker	153	3.2 (2.6, 3.8)	149	3.0 (2.4, 3.6)	133	2.8 (2.2, 3.4)	119	2.2 (1.7, 2.8)	108	2.1 (1.6, 2.6)
	Former smoker	1462	21.6 (20.4, 22.9)	1417	21.6 (20.4, 22.9)	1253	21.6 (20.3, 22.9)	1278	22.0 (20.7, 23.3)	1256	22.4 (21.0, 23.7)
	Never smoker	3818	70.2 (68.8, 71.7)	3775	70.8 (69.3, 72.2)	3502	70.9 (69.5, 72.4)	3628	71.4 (69.9, 72.9)	3494	71.1 (69.7, 72.6)

Education	Smoking Status	2021-23	
		N	Weighted% (95%CI)
Less than high school	Daily smoker	256	23.0 (20.2, 25.9)
	Someday smoker	71	6.4 (4.6, 8.1)
	Former smoker	301	24.9 (22.0, 27.9)
	Never smoker	470	45.7 (42.2, 49.2)
High school	Daily smoker	616	13.6 (12.4, 14.8)
	Someday smoker	209	5.3 (4.5, 6.1)
	Former smoker	1323	27.4 (25.8, 28.9)
	Never smoker	2235	53.7 (51.9, 55.5)
Some college	Daily smoker	636	11.2 (10.3, 12.2)
	Someday smoker	256	4.9 (4.2, 5.6)
	Former smoker	1649	28.7 (27.3, 30.1)
	Never smoker	2753	55.2 (53.6, 56.8)
College graduate	Daily smoker	268	3.9 (3.4, 4.5)
	Someday smoker	130	2.0 (1.6, 2.4)
	Former smoker	1496	22.6 (21.4, 23.8)
	Never smoker	4084	71.5 (70.2, 72.8)

Table 6. Three-year rolling weighted percentages of everyday and someday smokers among Oklahoma adults by income, Oklahoma BRFSS 2011 – 2023

Income	Smoking Status	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than 15,000	Daily smoker	961	32.5 (30.2, 34.9)	807	29.2 (27.0, 31.4)	664	28.4 (26.0, 30.9)	500	26.8 (24.0, 29.5)	404	24.7 (21.9, 27.6)
	Someday smoker	352	10.6 (9.1, 12.0)	309	10.4 (9.0, 11.9)	263	10.6 (8.9, 12.2)	198	9.8 (8.0, 11.5)	170	10.6 (8.6, 12.6)
	Former smoker	850	19.6 (17.9, 21.4)	798	20.7 (18.9, 22.5)	698	21.6 (19.5, 23.6)	572	22.1 (19.8, 24.3)	470	21.9 (19.4, 24.3)
	Never smoker	1256	37.3 (34.8, 39.7)	1184	39.7 (37.3, 42.1)	1014	39.4 (36.7, 42.2)	875	41.4 (38.3, 44.5)	752	42.8 (39.5, 46.1)
15,000 to Less than 25,000	Daily smoker	898	23.6 (21.8, 25.4)	819	22.9 (21.1, 24.7)	739	24.2 (22.2, 26.3)	621	23.0 (20.8, 25.2)	524	22.3 (20.1, 24.6)
	Someday smoker	301	8.7 (7.4, 10.1)	287	8.2 (6.9, 9.5)	239	7.3 (6.0, 8.5)	205	7.3 (6.0, 8.6)	169	7.7 (6.2, 9.2)
	Former smoker	1333	24.1 (22.5, 25.6)	1262	24.1 (22.6, 25.7)	1182	24.5 (22.7, 26.2)	1074	25.2 (23.3, 27.1)	931	24.0 (22.0, 25.9)
	Never smoker	2004	43.6 (41.6, 45.6)	1960	44.7 (42.7, 46.7)	1775	44.0 (41.8, 46.2)	1611	44.6 (42.2, 46.9)	1461	46.0 (43.4, 48.5)
25,000 to Less than 35,000	Daily smoker	448	19.7 (17.6, 21.8)	396	18.0 (16.0, 20.1)	324	16.5 (14.4, 18.7)	283	15.3 (13.1, 17.5)	241	14.9 (12.7, 17.2)
	Someday smoker	155	7.0 (5.6, 8.3)	160	7.9 (6.4, 9.4)	130	6.6 (5.2, 8.1)	120	7.2 (5.6, 8.8)	107	7.1 (5.3, 8.8)
	Former smoker	806	26.3 (24.1, 28.5)	813	26.3 (24.2, 28.4)	777	26.0 (23.8, 28.2)	715	26.1 (23.8, 28.5)	633	25.8 (23.4, 28.3)
	Never smoker	1306	47.0 (44.4, 49.6)	1317	47.7 (45.2, 50.3)	1257	50.8 (48.1, 53.6)	1145	51.4 (48.5, 54.3)	1002	52.1 (49.0, 55.2)
35,000 to Less than 50,000	Daily smoker	449	16.1 (14.4, 17.8)	403	14.6 (13.0, 16.2)	355	14.3 (12.6, 16.0)	306	14.2 (12.3, 16.1)	276	14.8 (12.7, 16.8)
	Someday smoker	143	5.4 (4.3, 6.4)	131	5.3 (4.3, 6.4)	129	5.4 (4.3, 6.5)	109	4.9 (3.7, 6.0)	107	5.2 (4.0, 6.5)
	Former smoker	1053	27.9 (26.0, 29.8)	997	27.8 (26.0, 29.7)	964	27.8 (25.8, 29.8)	893	27.2 (25.1, 29.3)	814	27.7 (25.5, 30.0)
	Never smoker	1674	50.6 (48.4, 52.9)	1686	52.2 (50.0, 54.4)	1613	52.5 (50.2, 54.9)	1511	53.7 (51.2, 56.3)	1343	52.3 (49.6, 55.0)
50,000 or more	Daily smoker	666	10.3 (9.4, 11.2)	634	9.3 (8.5, 10.2)	590	9.4 (8.5, 10.4)	556	9.3 (8.4, 10.3)	487	9.0 (8.0, 10.0)
	Someday smoker	256	4.0 (3.4, 4.6)	245	3.7 (3.2, 4.3)	242	4.4 (3.7, 5.2)	223	4.3 (3.5, 5.1)	204	4.7 (3.8, 5.5)

Income	Smoking Status	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
	Former smoker	2208	26.3 (25.1, 27.5)	2325	27.3 (26.1, 28.4)	2254	26.9 (25.6, 28.1)	2128	26.8 (25.5, 28.2)	1921	26.2 (24.9, 27.6)
	Never smoker	4580	59.4 (58.0, 60.7)	4785	59.7 (58.4, 61.0)	4635	59.3 (57.8, 60.7)	4488	59.5 (58.0, 61.0)	4084	60.1 (58.5, 61.7)

Income	Smoking Status	2016-18		2017-19		2018-20		2019-21		2020-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than 15,000	Daily smoker	336	24.6 (21.6, 27.5)	327	23.9 (21.0, 26.7)	292	25.2 (22.1, 28.3)	273	24.9 (21.7, 28.1)	230	25.5 (22.1, 28.9)
	Someday smoker	143	10.2 (8.2, 12.2)	136	10.7 (8.6, 12.7)	110	9.4 (7.3, 11.5)	90	8.1 (6.0, 10.2)	69	6.8 (4.7, 8.9)
	Former smoker	405	21.3 (18.8, 23.7)	376	21.7 (19.2, 24.3)	299	21.2 (18.4, 24.0)	285	23.3 (20.2, 26.4)	258	24.2 (20.9, 27.5)
	Never smoker	663	43.9 (40.6, 47.3)	573	43.7 (40.3, 47.2)	501	44.1 (40.5, 47.7)	460	43.7 (40.0, 47.4)	414	43.5 (39.6, 47.3)
15,000 to Less than 25,000	Daily smoker	453	20.8 (18.6, 23.0)	454	21.3 (19.2, 23.5)	433	21.0 (18.8, 23.2)	394	20.6 (18.3, 22.8)	328	19.4 (17.0, 21.7)
	Someday smoker	149	8.2 (6.5, 9.8)	154	8.4 (6.8, 10.1)	138	7.4 (5.8, 8.9)	130	7.1 (5.7, 8.6)	103	6.7 (5.2, 8.3)
	Former smoker	799	23.4 (21.4, 25.4)	731	22.6 (20.7, 24.6)	665	23.7 (21.6, 25.7)	585	23.7 (21.6, 25.9)	491	25.0 (22.6, 27.5)
	Never smoker	1339	47.7 (45.1, 50.2)	1281	47.6 (45.1, 50.2)	1144	48.0 (45.4, 50.6)	1031	48.6 (45.9, 51.3)	799	48.9 (45.9, 51.9)
25,000 to Less than 35,000	Daily smoker	240	16.4 (14.0, 18.7)	235	16.7 (14.3, 19.1)	236	18.3 (15.7, 20.9)	246	17.8 (15.4, 20.3)	271	17.7 (15.4, 20.0)
	Someday smoker	111	8.2 (6.3, 10.2)	89	7.0 (5.2, 8.8)	77	6.1 (4.5, 7.8)	70	5.7 (4.1, 7.3)	75	4.9 (3.6, 6.2)
	Former smoker	566	25.7 (23.1, 28.3)	533	26.1 (23.5, 28.8)	463	26.9 (24.1, 29.7)	471	27.2 (24.4, 29.9)	514	27.7 (25.1, 30.3)
	Never smoker	886	49.7 (46.5, 52.8)	848	50.2 (47.0, 53.3)	761	48.7 (45.4, 51.9)	796	49.3 (46.1, 52.4)	852	49.7 (46.7, 52.7)
35,000 to Less than 50,000	Daily smoker	263	14.6 (12.5, 16.6)	243	13.0 (11.1, 14.8)	220	11.9 (10.1, 13.8)	225	12.3 (10.5, 14.2)	215	11.9 (10.1, 13.7)
	Someday smoker	101	5.7	102	6.2	87	6.8	87	6.6	78	6.2

Income	Smoking Status	2016-18		2017-19		2018-20		2019-21		2020-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
			(4.3, 7.1)		(4.7, 7.6)		(5.0, 8.6)		(4.8, 8.3)		(4.4, 8.0)
	Former smoker	732	27.6 (25.3, 29.9)	694	28.2 (25.9, 30.5)	627	27.1 (24.7, 29.4)	573	25.9 (23.5, 28.2)	546	25.8 (23.4, 28.2)
	Never smoker	1221	52.1 (49.3, 54.8)	1148	52.6 (49.9, 55.4)	1107	54.2 (51.4, 57.0)	1110	55.2 (52.4, 58.1)	1099	56.1 (53.3, 58.9)
	Daily smoker	442	8.2 (7.3, 9.2)	438	7.9 (7.0, 8.8)	432	8.1 (7.2, 9.0)	431	7.8 (6.9, 8.6)	432	7.5 (6.6, 8.3)
	Someday smoker	197	4.0 (3.3, 4.7)	226	4.3 (3.7, 5.0)	209	3.9 (3.3, 4.5)	203	3.9 (3.2, 4.5)	170	3.2 (2.6, 3.7)
50,000 or more	Former smoker	1847	27.0 (25.6, 28.3)	1865	26.8 (25.5, 28.2)	1702	26.8 (25.5, 28.2)	1742	27.4 (26.0, 28.8)	1745	27.9 (26.6, 29.3)
	Never smoker	3933	60.8 (59.3, 62.4)	3977	60.9 (59.4, 62.4)	3691	61.2 (59.6, 62.7)	3756	60.9 (59.4, 62.5)	3745	61.5 (60.0, 63.0)

Income	Smoking Status	2021-23	
		N	Weighted% (95%CI)
Less than 15,000	Daily smoker	215	25.8 (22.2, 29.5)
	Someday smoker	70	7.4 (5.3, 9.5)
	Former smoker	242	24.3 (20.9, 27.8)
	Never smoker	365	42.4 (38.4, 46.5)
15,000 to Less than 25,000	Daily smoker	269	19.0 (16.5, 21.6)
	Someday smoker	88	6.6 (5.0, 8.2)
	Former smoker	429	26.4 (23.6, 29.2)
	Never smoker	647	48.0 (44.7, 51.3)
	Daily smoker	295	17.3 (15.2, 19.4)

Income	Smoking Status	2021-23	
		N	Weighted% (95%CI)
25,000 to Less than 35,000	Someday smoker	74	4.8 (3.5, 6.1)
	Former smoker	562	27.9 (25.4, 30.3)
	Never smoker	918	50.1 (47.2, 52.9)
35,000 to Less than 50,000	Daily smoker	257	11.8 (10.1, 13.4)
	Someday smoker	106	5.7 (4.4, 7.0)
	Former smoker	648	26.2 (24.0, 28.4)
	Never smoker	1244	56.3 (53.8, 58.9)
50,000 or more	Daily smoker	488	7.0 (6.3, 7.7)
	Someday smoker	217	3.4 (2.8, 3.9)
	Former smoker	2150	27.9 (26.7, 29.2)
	Never smoker	4645	61.7 (60.3, 63.0)

Table 7. Proportion of everyday and someday smokers among Oklahoma adults by area of residence, Oklahoma BRFSS 2011 – 2023

Area of Residence	Smoking Status	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Urban	Daily smoker	582	19.2 (17.3, 21.2)	350	13.3 (11.8, 14.9)	393	14.5 (12.9, 16.2)	305	12.1 (10.5, 13.7)	262	13.6 (11.5, 15.6)
	Someday smoker	225	6.5 (5.3, 7.7)	158	5.9 (4.8, 7.0)	188	6.4 (5.3, 7.5)	161	6.3 (5.2, 7.5)	116	6.7 (4.9, 8.5)
	Former smoker	1057	23.2 (21.4, 25)	748	24.2 (22.4, 26.1)	937	24.2 (22.4, 26)	870	25.1 (23.3, 27)	699	23.3 (21.1, 25.4)
	Never smoker	1865	51.1 (48.8, 53.4)	1537	56.5 (54.2, 58.8)	1810	54.9 (52.7, 57.1)	1657	56.4 (54.2, 58.7)	1351	56.5 (53.6, 59.3)
Rural	Daily smoker	812	20.4 (18.7, 22.1)	864	19.6 (18.0, 21.2)	747	19.5 (17.8, 21.1)	725	17.3 (15.8, 18.8)	580	17.8 (15.9, 19.6)
	Someday smoker	238	6.1 (5.0, 7.2)	264	6.6 (5.6, 7.7)	239	6.3 (5.3, 7.3)	236	5.7 (4.7, 6.6)	190	5.8 (4.7, 6.9)
	Former smoker	1425	25.8 (24.2, 27.4)	1453	24.5 (23.0, 25.9)	1390	24.8 (23.3, 26.3)	1534	26.5 (25, 28)	1293	25.8 (23.9, 27.6)
	Never smoker	2304	47.8 (45.7, 49.8)	2574	49.3 (47.4, 51.2)	2430	49.5 (47.6, 51.4)	2709	50.5 (48.7, 52.3)	2262	50.6 (48.4, 52.9)

Area of Residence	Smoking Status	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Urban	Daily smoker	202	11.0 (9.1, 12.8)	217	11.4 (9.7, 13.1)	165	10.3 (8.6, 12.1)	184	10.9 (9.2, 12.7)	169	12.2 (10.2, 14.2)
	Someday smoker	106	5.8 (4.4, 7.2)	105	7.1 (5.4, 8.7)	86	5.9 (4.5, 7.3)	88	5.7 (4.4, 7.0)	67	5.1 (3.7, 6.5)
	Former smoker	662	22.2 (20.2, 24.2)	642	24.2 (22.0, 26.4)	519	23.9 (21.6, 26.3)	551	24.1 (22.0, 26.2)	429	23.3 (20.9, 25.7)
	Never smoker	1441	61.1 (58.4, 63.7)	1261	57.3 (54.6, 60.0)	1084	59.8 (57.1, 62.6)	1208	59.3 (56.7, 61.9)	962	59.4 (56.5, 62.3)
Rural	Daily smoker	527	16.3 (14.6, 18.0)	485	14.7 (13.2, 16.2)	430	16.3 (14.5, 18.0)	499	14.9 (13.4, 16.4)	439	15.4 (13.8, 17.0)
	Someday smoker	175	5.4 (4.3, 6.4)	201	6.7 (5.6, 7.9)	158	6.1 (4.9, 7.2)	198	5.8 (4.8, 6.7)	123	5.0 (3.8, 6.2)
	Former smoker	1252	26.8 (25.1, 28.6)	1214	24.8 (23.2, 26.4)	920	25.5 (23.7, 27.4)	1148	26.2 (24.5, 27.9)	878	26.7 (24.8, 28.6)
	Never smoker	2325	51.5 (49.4, 53.5)	2246	53.8 (51.8, 55.8)	1698	52.1 (49.9, 54.4)	2160	53.2 (51.2, 55.2)	1655	52.9 (50.7, 55.1)

Area of Residence	Smoking Status	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Urban	Daily smoker	135	8.7 (7.0, 10.4)	175	9.7 (8.2, 11.3)	169	8.2 (6.9, 9.6)
	Someday smoker	56	4.2 (2.9, 5.5)	67	3.8 (2.7, 4.8)	93	4.8 (3.7, 5.9)
	Former smoker	417	26.9 (24.3, 29.6)	522	26.3 (24.1, 28.5)	644	24.8 (22.8, 26.7)
	Never smoker	934	60.1 (57.1, 63.1)	1099	60.2 (57.6, 62.8)	1405	62.2 (59.9, 64.5)
Rural	Daily smoker	424	14.8 (13.3, 16.4)	402	13.1 (11.6, 14.5)	473	13.4 (12.0, 14.7)
	Someday smoker	152	5.2 (4.2, 6.2)	134	4.2 (3.4, 5.0)	165	4.7 (3.9, 5.5)
	Former smoker	991	24.7 (23.0, 26.5)	1033	27.9 (26.1, 29.6)	1176	27.1 (25.5, 28.8)
	Never smoker	1984	55.3 (53.2, 57.4)	1938	54.9 (52.9, 56.9)	2232	54.8 (52.9, 56.8)

Table 8. Proportion of everyday and someday smokers among Oklahoma adults by poor physical health, Oklahoma BRFSS 2011 – 2023

Poor Physical Health	Smoking Status	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Daily smoker	342	28.0 (24.7, 31.4)	305	26.6 (23.2, 30.0)	283	25.2 (22.0, 28.4)	244	22.1 (19.0, 25.1)	197	20.7 (17.2, 24.2)
	Someday smoker	124	8.7 (6.6, 10.8)	98	7.9 (6.0, 9.7)	105	9.0 (7.0, 11.1)	106	9.1 (6.9, 11.2)	84	11.7 (8.0, 15.4)
	Former smoker	491	30.2 (27.0, 33.3)	411	29.7 (26.5, 32.9)	431	29.3 (26.3, 32.4)	494	33.7 (30.5, 36.8)	400	32.7 (28.8, 36.6)
	Never smoker	507	33.1 (29.8, 36.5)	519	35.8 (32.4, 39.3)	516	36.4 (33.1, 39.8)	494	35.2 (31.9, 38.4)	417	34.9 (30.7, 39.1)
No	Daily smoker	1052	18.6 (17.2, 20.0)	909	15.4 (14.2, 16.6)	857	16.1 (14.9, 17.4)	786	13.9 (12.8, 15.1)	645	15.2 (13.7, 16.7)
	Someday smoker	339	5.8 (5.0, 6.7)	324	6.1 (5.2, 6.9)	322	5.9 (5.1, 6.7)	291	5.4 (4.6, 6.2)	222	5.2 (4.3, 6.2)
	Former smoker	1991	23.9 (22.5, 25.2)	1790	23.5 (22.3, 24.8)	1896	23.8 (22.5, 25.0)	1910	24.7 (23.4, 25.9)	1592	23.3 (21.8, 24.8)
	Never smoker	3662	51.7 (50.0, 53.4)	3592	55.0 (53.4, 56.6)	3724	54.2 (52.7, 55.8)	3872	56.0 (54.4, 57.5)	3196	56.3 (54.4, 58.2)
Poor Physical Health	Smoking Status	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Daily smoker	200	24.2 (20.3, 28.0)	170	19.9 (16.5, 23.3)	151	22.6 (18.8, 26.5)	173	21.4 (17.8, 24.9)	149	25.3 (21.1, 29.6)
	Someday smoker	60	7.7 (5.4, 9.9)	80	9.9 (7.3, 12.4)	55	9.0 (6.1, 11.8)	69	9.4 (6.8, 12.0)	37	6.2 (3.8, 8.7)
	Former smoker	358	30.9 (27.3, 34.4)	348	30.8 (27.2, 34.4)	268	28.1 (24.2, 31.9)	295	29.0 (25.5, 32.5)	204	30.9 (26.4, 35.3)
	Never smoker	464	37.3 (33.5, 41.1)	413	39.4 (35.3, 43.6)	339	40.3 (35.9, 44.7)	378	40.2 (36.2, 44.2)	268	37.6 (33.0, 42.2)
No	Daily smoker	505	12.1 (10.8, 13.4)	506	11.9 (10.7, 13.1)	424	12.0 (10.6, 13.3)	478	11.6 (10.4, 12.9)	435	12.1 (10.8, 13.4)
	Someday smoker	202	4.8 (4.0, 5.7)	215	6.3 (5.3, 7.4)	181	5.3 (4.4, 6.2)	202	4.9 (4.1, 5.7)	145	4.9 (3.9, 5.9)
	Former smoker	1504	23.7 (22.3, 25.2)	1432	23.2 (21.7, 24.6)	1131	24.1 (22.5, 25.7)	1338	24.6 (23.1, 26.1)	1055	24.4 (22.7, 26.0)
	Never smoker	3205	59.3 (57.5, 61.1)	3016	58.7 (56.9, 60.4)	2389	58.6 (56.7, 60.5)	2862	58.9 (57.1, 60.6)	2261	58.6 (56.7, 60.6)

Poor Physical Health	Smoking Status	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Daily smoker	123	20.5 (16.7, 24.3)	157	21.5 (18.1, 24.9)	142	18.6 (15.3, 21.9)
	Someday smoker	44	8.5 (5.5, 11.5)	44	5.9 (3.9, 7.9)	60	8.1 (5.7, 10.4)
	Former smoker	214	28.8 (24.6, 33.0)	275	32.8 (29.0, 36.5)	304	32.6 (28.8, 36.4)
	Never smoker	297	42.2 (37.6, 46.9)	337	39.9 (35.9, 43.9)	370	40.8 (36.8, 44.8)
No	Daily smoker	408	10.9 (9.6, 12.1)	399	9.8 (8.7, 10.9)	469	9.6 (8.7, 10.6)
	Someday smoker	149	4.0 (3.2, 4.8)	149	3.7 (3.0, 4.4)	182	4.0 (3.4, 4.7)
	Former smoker	1131	25.2 (23.6, 26.9)	1222	26.1 (24.6, 27.6)	1423	24.7 (23.3, 26.1)
	Never smoker	2499	59.9 (57.9, 61.8)	2617	60.4 (58.7, 62.1)	3146	61.6 (60.0, 63.2)

Table 9. Proportion of everyday and someday smokers among Oklahoma adults by Frequent Mental Distress (FMD), Oklahoma BRFSS 2011 – 2023

Frequent Mental Distress	Smoking Status	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Daily smoker	379	33.8 (29.9, 37.6)	304	31.3 (27.5, 35.1)	303	34.1 (30.0, 38.2)	255	28.7 (25.1, 32.3)	186	28.4 (23.8, 32.9)
	Someday smoker	114	9.4 (6.8, 12.1)	110	11.6 (8.7, 14.5)	100	11.1 (8.6, 13.7)	93	10.2 (7.7, 12.7)	79	12.8 (9.0, 16.6)
	Former smoker	328	22.6 (19.5, 25.7)	248	20.8 (17.7, 23.9)	276	21.9 (18.8, 25)	301	24.7 (21.6, 27.9)	211	22.5 (18.6, 26.4)
	Never smoker	395	34.2 (30.2, 38.1)	369	36.3 (32.3, 40.3)	375	32.9 (29, 36.7)	381	36.3 (32.6, 40.1)	280	36.4 (31.4, 41.3)
No	Daily smoker	1015	17.5 (16.2, 18.9)	910	14.8 (13.6, 16.0)	837	14.6 (13.5, 15.8)	775	13.0 (11.9, 14.1)	656	14.2 (12.7, 15.6)
	Someday smoker	349	5.7 (4.8, 6.5)	312	5.5 (4.8, 6.3)	327	5.5 (4.8, 6.3)	304	5.3 (4.5, 6.1)	227	5.2 (4.2, 6.2)
	Former smoker	2154	25.1 (23.8, 26.4)	1953	24.9 (23.7, 26.2)	2051	25.0 (23.7, 26.2)	2103	26.1 (24.9, 27.4)	1781	25.0 (23.5, 26.5)
	Never smoker	3774	51.7 (50.1, 53.4)	3742	54.8 (53.2, 56.3)	3865	54.9 (53.3, 56.4)	3985	55.5 (54.0, 57.0)	3333	55.6 (53.7, 57.5)

Frequent Mental Distress	Smoking Status	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Daily smoker	180	27.4 (23.1, 31.7)	163	23.3 (19.5, 27.0)	154	24.3 (20.1, 28.4)	178	24.7 (20.8, 28.6)	175	26.7 (22.6, 30.7)
	Someday smoker	62	8.6 (5.7, 11.6)	71	10.6 (7.4, 13.7)	61	10.7 (7.6, 13.8)	65	9.1 (6.5, 11.6)	44	7.1 (4.7, 9.4)
	Former smoker	234	24.5 (20.9, 28.2)	233	22.9 (19.6, 26.2)	176	21.7 (17.9, 25.4)	220	24.4 (20.8, 27.9)	181	25.5 (21.6, 29.4)
	Never smoker	333	39.4 (34.9, 43.9)	338	43.3 (38.8, 47.8)	276	43.4 (38.5, 48.3)	324	41.9 (37.6, 46.2)	268	40.8 (36.2, 45.3)
No	Daily smoker	532	11.7 (10.4, 12.9)	515	11.1 (10.0, 12.2)	418	11.5 (10.2, 12.8)	466	10.7 (9.5, 11.8)	406	11.4 (10.1, 12.7)
	Someday smoker	209	4.9 (4.1, 5.8)	229	6.2 (5.2, 7.2)	173	5.0 (4.0, 5.9)	212	5.1 (4.3, 5.9)	137	4.5 (3.6, 5.5)
	Former smoker	1643	25.0 (23.5, 26.4)	1577	24.9 (23.4, 26.3)	1237	25.6 (23.9, 27.2)	1426	25.6 (24.1, 27.1)	1080	25.2 (23.6, 26.9)
	Never smoker	3362	58.4 (56.6, 60.1)	3105	57.8 (56.0, 59.5)	2449	58.0 (56.1, 59.9)	2930	58.6 (56.9, 60.3)	2261	58.8 (56.9, 60.8)

Frequent Mental Distress	Smoking Status	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Daily smoker	148	21.2 (17.5, 24.9)	142	18.4 (15.2, 21.7)	174	17.9 (14.9, 20.9)
	Someday smoker	47	6.8 (4.5, 9.0)	45	5.1 (3.4, 6.9)	71	8.1 (5.9, 10.2)
	Former smoker	213	26.4 (22.5, 30.3)	254	27.1 (23.7, 30.5)	293	29.3 (25.8, 32.7)
	Never smoker	333	45.6 (41.2, 50.1)	415	49.3 (45.4, 53.3)	404	44.8 (40.9, 48.7)
No	Daily smoker	382	9.9 (8.8, 11.1)	411	9.8 (8.7, 10.8)	443	9.6 (8.6, 10.7)
	Someday smoker	152	4.3 (3.5, 5.2)	148	3.7 (3.0, 4.5)	179	4.0 (3.3, 4.7)
	Former smoker	1148	25.6 (23.9, 27.3)	1252	27.2 (25.6, 28.8)	1467	25.5 (24.1, 26.9)
	Never smoker	2503	60.2 (58.2, 62.1)	2536	59.3 (57.5, 61.0)	3120	60.9 (59.2, 62.5)

Frequent Mental Distress = 14 or more days of poor mental health

Table 10. Proportion of everyday and someday smokers among Oklahoma adults by health insurance coverage, Oklahoma BRFSS 2011 – 2023

Health Insurance Status	Smoking Status	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Insured	Daily smoker	1039	17.0 (15.7, 18.3)	924	14.2 (13.1, 15.3)	853	14.6 (13.4, 15.8)	819	13.0 (12.0, 14.1)	696	14.0 (12.6, 15.4)
	Someday smoker	353	5.3 (4.5, 6.1)	322	5.7 (4.8, 6.5)	320	5.4 (4.7, 6.2)	335	5.6 (4.8, 6.3)	259	5.7 (4.7, 6.8)
	Former smoker	2227	26.9 (25.5, 28.2)	2023	26.6 (25.3, 27.9)	2132	26.2 (24.9, 27.4)	2256	27.3 (26.0, 28.5)	1897	26.4 (24.9, 27.9)
	Never smoker	3644	50.8 (49.2, 52.4)	3615	53.6 (52.0, 55.1)	3791	53.8 (52.3, 55.3)	4003	54.1 (52.6, 55.6)	3374	53.8 (52.0, 55.7)
Uninsured	Daily smoker	351	30.3 (26.7, 33.8)	287	29.8 (26.2, 33.4)	283	29.3 (25.9, 32.8)	209	27.9 (23.9, 31.9)	143	28.9 (23.8, 33.9)
	Someday smoker	110	9.6 (7.2, 12.0)	100	9.5 (7.3, 11.7)	106	10.5 (8.2, 12.8)	60	8.1 (5.7, 10.5)	46	9.4 (6.1, 12.6)
	Former smoker	248	17.0 (14.3, 19.6)	176	14.8 (12.2, 17.3)	189	17.9 (15, 20.7)	142	17.6 (14.3, 20.8)	93	14.5 (10.9, 18.0)
	Never smoker	516	43.1 (39.3, 47.0)	483	45.9 (42.1, 49.7)	437	42.3 (38.6, 46.1)	352	46.4 (42.1, 50.8)	224	47.3 (41.6, 53)
Health Insurance Status	Smoking Status	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Insured	Daily smoker	595	12.3 (11.1, 13.6)	567	11.7 (10.6, 12.8)	476	12.3 (11.0, 13.6)	540	12.0 (10.8, 13.2)	495	13.1 (11.8, 14.4)
	Someday smoker	238	4.9 (4.1, 5.7)	262	6.5 (5.5, 7.5)	202	5.2 (4.4, 6.1)	234	5.2 (4.4, 5.9)	164	5.2 (4.2, 6.2)
	Former smoker	1794	26.1 (24.7, 27.6)	1748	26.6 (25.1, 28.0)	1335	26.0 (24.4, 27.5)	1573	26.7 (25.2, 28.1)	1182	26.1 (24.5, 27.7)
	Never smoker	3486	56.6 (54.9, 58.3)	3209	55.2 (53.5, 56.9)	2528	56.5 (54.6, 58.4)	3019	56.2 (54.5, 57.9)	2333	55.6 (53.7, 57.5)
Uninsured	Daily smoker	132	24.2 (19.7, 28.8)	134	22.4 (18.3, 26.6)	118	22.4 (18.1, 26.7)	142	19.7 (16.2, 23.3)	111	19.6 (15.7, 23.5)
	Someday smoker	42	9.2 (5.9, 12.5)	44	9.1 (5.9, 12.3)	41	10.8 (7.1, 14.4)	52	9.0 (6.3, 11.7)	25	4.5 (2.4, 6.5)
	Former smoker	114	17.7 (14.1, 21.3)	105	13.4 (10.4, 16.5)	99	18.1 (14.1, 22.1)	123	17.9 (14.5, 21.3)	121	21.1 (17.0, 25.3)
	Never smoker	268	48.8 (43.6, 54.0)	282	55.0 (49.8, 60.1)	245	48.7 (43.5, 54.0)	340	53.4 (48.8, 57.9)	267	54.8 (49.7, 59.9)

Health Insurance Status	Smoking Status	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Insured	Daily smoker	448	10.8 (9.7, 12.0)	505	11.1 (10.0, 12.2)	572	11.0 (9.9, 12.0)
	Someday smoker	187	4.9 (4.0, 5.7)	181	4.0 (3.3, 4.6)	218	4.3 (3.7, 5.0)
	Former smoker	1310	27.0 (25.3, 28.6)	1449	28.0 (26.5, 29.5)	1709	27.5 (26.1, 28.8)
	Never smoker	2661	57.3 (55.5, 59.2)	2775	56.9 (55.3, 58.6)	3307	57.2 (55.7, 58.8)
Uninsured	Daily smoker	98	23.4 (18.6, 28.3)	63	16.2 (12.0, 20.3)	58	12.7 (9.2, 16.2)
	Someday smoker	15	4.2 (1.8, 6.7)	18	4.8 (2.3, 7.3)	35	9.1 (5.8, 12.3)
	Former smoker	75	18.8 (14.1, 23.6)	80	20.9 (16.3, 25.6)	75	15.8 (11.9, 19.7)
	Never smoker	176	53.5 (47.4, 59.5)	208	58.1 (52.3, 63.9)	238	62.4 (57.0, 67.7)

Table 11. Proportion of everyday and someday smokers among Oklahoma adults by disability status, Oklahoma BRFSS 2016 – 2023

Disability Status	Smoking Status	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
One or more disability	Daily smoker	368	20.3 (17.9, 22.7)	362	19.4 (17.2, 21.6)	311	20.3 (17.8, 22.8)	360	18.1 (16.0, 20.1)	331	21.1 (18.7, 23.6)
	Someday smoker	135	7.6 (6.0, 9.2)	148	8.5 (6.8, 10.2)	101	6.7 (5.1, 8.3)	133	7.1 (5.7, 8.5)	98	7.0 (5.3, 8.7)
	Former smoker	798	29.8 (27.4, 32.1)	861	29.2 (27.0, 31.5)	670	30.7 (28.1, 33.4)	804	31.8 (29.4, 34.1)	586	30.4 (27.7, 33.0)
	Never smoker	1193	42.3 (39.6, 44.9)	1136	42.9 (40.1, 45.6)	900	42.3 (39.4, 45.2)	1104	43.1 (40.5, 45.6)	795	41.5 (38.5, 44.4)
No disability	Daily smoker	345	10.8 (9.3, 12.2)	330	9.9 (8.7, 11.1)	282	10.3 (8.8, 11.7)	312	10.3 (9.0, 11.7)	270	10.3 (8.8, 11.7)
	Someday smoker	141	4.5 (3.5, 5.4)	156	6.1 (4.9, 7.3)	141	5.5 (4.4, 6.6)	149	5.0 (4.0, 5.9)	85	3.9 (2.8, 4.9)
	Former smoker	1087	22.4 (20.8, 24.0)	976	22.0 (20.4, 23.7)	762	21.8 (20.0, 23.5)	873	21.4 (19.8, 23.0)	696	22.3 (20.5, 24.1)
	Never smoker	2525	62.4 (60.4, 64.4)	2334	62.0 (60.0, 64.0)	1874	62.5 (60.3, 64.6)	2225	63.3 (61.3, 65.3)	1797	63.6 (61.4, 65.8)

Disability Status	Smoking Status	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
One or more disability	Daily smoker	283	17.4 (15.2, 19.6)	317	16.2 (14.2, 18.1)	359	16.5 (14.7, 18.4)
	Someday smoker	109	6.8 (5.3, 8.4)	105	5.2 (4.0, 6.3)	129	6.1 (4.9, 7.3)
	Former smoker	637	29.8 (27.2, 32.4)	778	33.0 (30.7, 35.3)	808	30.0 (27.8, 32.1)
	Never smoker	909	46.0 (43.1, 48.9)	1045	45.7 (43.1, 48.2)	1158	47.4 (44.9, 49.9)
No disability	Daily smoker	261	9.2 (7.9, 10.5)	254	8.7 (7.5, 9.9)	270	7.6 (6.5, 8.6)
	Someday smoker	93	3.6 (2.7, 4.5)	94	3.3 (2.5, 4.1)	120	3.7 (2.9, 4.4)
	Former smoker	750	23.8 (21.8, 25.7)	756	23.2 (21.5, 24.9)	969	23.4 (21.8, 25.0)
	Never smoker	1948	63.5 (61.3, 65.7)	1949	64.8 (62.8, 66.8)	2418	65.3 (63.5, 67.2)

Section III. Smokeless tobacco (ST) use

Table 1. Proportion of ST users among Oklahoma adults, Oklahoma BRFSS 2011 – 2023

Smokeless Tobacco Use	2011		2012		2013		2014		2015	
	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	381	7.0 (6.0, 7.9)	399	6.7 (5.9, 7.4)	360	6.3 (5.5, 7.1)	374	6.5 (5.7, 7.2)	307	6.6 (5.7, 7.6)
No	8141	93.0 (92.1, 94.0)	7556	93.3 (92.6, 94.1)	7788	93.7 (92.9, 94.5)	7839	93.6 (92.8, 94.3)	6459	93.4 (92.4, 94.3)

Smokeless Tobacco Use	2016		2017		2018		2019		2020	
	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	284	6.0 (5.2, 6.9)	332	7.4 (6.4, 8.4)	262	6.8 (5.8, 7.8)	366	7.3 (6.4, 8.2)	299	7.1 (6.1, 8.0)
No	6432	94.0 (93.1, 94.8)	6090	92.6 (91.6, 93.6)	4816	93.2 (92.2, 94.2)	5728	92.7 (91.8, 93.6)	4492	92.9 (92.0, 93.9)

Smokeless Tobacco Use	2021		2022		2023	
	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	288	6.4 (5.5, 7.2)	287	5.7 (5.0, 6.4)	349	6.3 (5.5, 7.0)
No	4875	93.6 (92.8, 94.5)	5134	94.3 (93.6, 95.0)	6070	93.7 (93.0, 94.5)

Table 2. Proportion of ST users among Oklahoma adults by sex, Oklahoma BRFSS 2011 – 2023

Sex	Smokeless Tobacco Use	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Male	Yes	345	13.4 (11.6, 15.1)	361	12.6 (11.1, 14.1)	320	11.8 (10.3, 13.3)	350	12.6 (11.1, 14.2)	277	12.5 (10.7, 14.3)
	No	2823	86.6 (84.9, 88.4)	2852	87.4 (85.9, 88.9)	2915	88.2 (86.7, 89.7)	2995	87.4 (85.8, 88.9)	2404	87.5 (85.7, 89.3)
Female	Yes	36	0.8 (0.5, 1.2)	38	1.0 (0.6, 1.4)	40	1.0 (0.6, 1.5)	24	0.5 (0.3, 0.8)	30	1.0 (0.4, 1.5)
	No	5318	99.2 (98.8, 99.5)	4704	99.0 (98.6, 99.4)	4873	99.0 (98.5, 99.4)	4844	99.5 (99.2, 99.7)	4055	99.0 (98.5, 99.6)

Sex	Smokeless Tobacco Use	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Male	Yes	256	11.4 (9.7, 13.1)	302	13.7 (11.9, 15.4)	227	12.2 (10.4, 14.0)	339	13.9 (12.2, 15.6)	260	12.6 (10.8, 14.3)
	No	2463	88.6 (86.9, 90.3)	2309	86.3 (84.6, 88.1)	1908	87.8 (86.0, 89.6)	2298	86.1 (84.4, 87.8)	1802	87.4 (85.7, 89.2)
Female	Yes	28	0.9 (0.5, 1.3)	29	1.3 (0.6, 1.9)	35	1.5 (0.9, 2.1)	27	1.1 (0.6, 1.5)	39	1.8 (1.1, 2.5)
	No	3969	99.1 (98.7, 99.5)	3780	98.7 (98.1, 99.4)	2907	98.5 (97.9, 99.1)	3430	98.9 (98.5, 99.4)	2690	98.2 (97.5, 98.9)

Sex	Smokeless Tobacco Use	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Male	Yes	257	11.9 (10.2, 13.6)	253	10.5 (9.1, 11.9)	297	10.8 (9.4, 12.1)
	No	2017	88.1 (86.4, 89.8)	2188	89.5 (88.1, 90.9)	2576	89.2 (87.9, 90.6)
Female	Yes	31	1.1 (0.7, 1.6)	34	1.1 (0.7, 1.5)	52	2.0 (1.4, 2.6)
	No	2858	98.9 (98.4, 99.3)	2946	98.9 (98.5, 99.3)	3494	98.0 (97.4, 98.6)

Table 3. Three-year rolling weighted percentages of ST users among Oklahoma male adults by age, Oklahoma BRFSS 2011 – 2023

Age Group	Smokeless Tobacco Use	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
18-24	Yes	85	14.1 (10.9, 17.4)	96	15.9 (12.6, 19.1)	86	16.6 (13.0, 20.2)	73	16.0 (12.1, 19.9)	64	15.6 (11.6, 19.5)
	No	446	85.9 (82.6, 89.1)	476	84.1 (80.9, 87.4)	414	83.4 (79.8, 87.0)	380	84.0 (80.1, 87.9)	351	84.4 (80.5, 88.4)
25-34	Yes	149	15.4 (12.8, 18.0)	154	15.0 (12.6, 17.4)	127	13.9 (11.4, 16.4)	126	14.2 (11.6, 16.8)	115	13.9 (11.3, 16.6)
	No	903	84.6 (82.0, 87.2)	911	85.0 (82.6, 87.4)	812	86.1 (83.6, 88.6)	771	85.8 (83.2, 88.4)	746	86.1 (83.4, 88.7)
35-44	Yes	183	16.5 (14.0, 19.0)	182	16.1 (13.6, 18.5)	146	14.6 (12.1, 17.1)	129	13.9 (11.3, 16.5)	136	15.9 (13.0, 18.8)
	No	990	83.5 (81.0, 86.0)	1041	83.9 (81.5, 86.4)	924	85.4 (82.9, 87.9)	830	86.1 (83.5, 88.7)	703	84.1 (81.2, 87.0)
45-54	Yes	228	13.6 (11.7, 15.6)	223	13.0 (11.2, 14.8)	219	15.0 (12.8, 17.2)	202	15.2 (12.9, 17.4)	172	14.9 (12.5, 17.3)
	No	1425	86.4 (84.4, 88.3)	1422	87.0 (85.2, 88.8)	1273	85.0 (82.8, 87.2)	1136	84.8 (82.6, 87.1)	954	85.1 (82.7, 87.5)
55-64	Yes	175	8.2 (6.8, 9.6)	167	7.3 (6.0, 8.5)	154	7.5 (6.2, 8.9)	151	8.1 (6.7, 9.6)	169	9.8 (8.1, 11.4)
	No	1985	91.8 (90.4, 93.2)	2026	92.7 (91.5, 94.0)	1949	92.5 (91.1, 93.8)	1826	91.9 (90.4, 93.3)	1622	90.2 (88.6, 91.9)
65 or more	Yes	206	7.2 (6.1, 8.4)	209	7.2 (6.1, 8.3)	215	7.0 (5.9, 8.1)	202	6.5 (5.5, 7.6)	179	6.1 (5.1, 7.2)
	No	2841	92.8 (91.6, 93.9)	2884	92.8 (91.7, 93.9)	2940	93.0 (91.9, 94.1)	2917	93.5 (92.4, 94.5)	2800	93.9 (92.8, 94.9)

Age Group	Smokeless Tobacco Use	2016-18		2017-19		2018-20		2019-21		2020-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
18-24	Yes	65	15.6 (11.6, 19.5)	74	17.5 (13.5, 21.6)	63	15.6 (11.7, 19.5)	57	12.8 (9.3, 16.3)	40	7.6 (5.1, 10.2)
	No	351	84.4 (80.5, 88.4)	364	82.5 (78.4, 86.5)	358	84.4 (80.5, 88.3)	409	87.2 (83.7, 90.7)	446	92.4 (89.8, 94.9)
25-34	Yes	110	13.7 (10.9, 16.4)	111	13.6 (10.9, 16.3)	101	12.7 (10.0, 15.4)	109	13.6 (10.9, 16.4)	108	13.9 (11.1, 16.7)
	No	752	86.3 (83.6, 89.1)	767	86.4 (83.7, 89.1)	711	87.3 (84.6, 90.0)	720	86.4 (83.6, 89.1)	694	86.1 (83.3, 88.9)
35-44	Yes	133	15.5 (12.7, 18.3)	175	17.0 (14.3, 19.7)	150	14.8 (12.3, 17.3)	158	14.7 (12.2, 17.1)	123	12.7 (10.3, 15.1)
	No	693	84.5 (81.7, 87.3)	770	83.0 (80.3, 85.7)	777	85.2 (82.7, 87.7)	834	85.3 (82.9, 87.8)	829	87.3 (84.9, 89.7)
45-54	Yes	155	14.3 (11.9, 16.7)	151	13.9 (11.5, 16.2)	160	14.9 (12.5, 17.4)	149	14.6 (12.0, 17.1)	148	15.0 (12.4, 17.6)
	No	885	85.7 (83.3, 88.1)	877	86.1 (83.8, 88.5)	854	85.1 (82.6, 87.5)	871	85.4 (82.9, 88.0)	882	85.0 (82.4, 87.6)
55-64	Yes	166	10.4 (8.6, 12.1)	181	11.7 (9.8, 13.5)	166	12.1 (10.1, 14.1)	178	12.8 (10.8, 14.9)	168	12.6 (10.5, 14.7)
	No	1430	89.6 (87.9, 91.4)	1308	88.3 (86.5, 90.2)	1170	87.9 (85.9, 89.9)	1118	87.2 (85.1, 89.2)	1069	87.4 (85.3, 89.5)
65 or more	Yes	156	6.3 (5.2, 7.4)	176	7.3 (6.0, 8.5)	186	8.5 (7.1, 9.9)	205	9.1 (7.7, 10.5)	183	8.3 (6.9, 9.7)
	No	2569	93.7 (92.6, 94.8)	2429	92.7 (91.5, 94.0)	2138	91.5 (90.1, 92.9)	2165	90.9 (89.5, 92.3)	2087	91.7 (90.3, 93.1)

Age Group	Smokeless Tobacco Use	2021-23	
		N	Weighted% (95%CI)
18-24	Yes	44	6.2 (4.3, 8.2)
	No	501	93.8 (91.8, 95.7)
25-34	Yes	118	14.1 (11.4, 16.8)
	No	794	85.9 (83.2, 88.6)
35-44	Yes	129	11.4 (9.3, 13.5)
	No	980	88.6 (86.5, 90.7)
45-54	Yes	173	15.7 (13.2, 18.2)
	No	968	84.3 (81.8, 86.8)
55-64	Yes	174	12.6 (10.5, 14.6)
	No	1175	87.4 (85.4, 89.5)
65 or more	Yes	169	7.0 (5.7, 8.2)
	No	2363	93.0 (91.8, 94.3)

Table 4. Three-year rolling weighted percentages of ST users among Oklahoma male adults by race/ethnicity, Oklahoma BRFSS 2011 – 2023

Race/Ethnicity	Smokeless Tobacco Use	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
NH-White	Yes	769	13.8 (12.7, 14.9)	774	13.3 (12.2, 14.3)	719	12.9 (11.8, 14.0)	681	12.8 (11.7, 13.9)	631	12.8 (11.6, 14.0)
	No	6342	86.2 (85.1, 87.3)	6475	86.7 (85.7, 87.8)	6182	87.1 (86.0, 88.2)	5914	87.2 (86.1, 88.3)	5417	87.2 (86.0, 88.4)
NH-Black	Yes	25	4.8 (2.5, 7.1)	24	4.8 (2.5, 7.1)	27	7.0 (3.9, 10.2)	24	6.5 (3.4, 9.6)	21	6.0 (2.9, 9.1)
	No	481	95.2 (92.9, 97.5)	508	95.2 (92.9, 97.5)	510	93.0 (89.8, 96.1)	467	93.5 (90.4, 96.6)	408	94.0 (90.9, 97.1)
NH-AI/AN	Yes	81	16.3 (12.5, 20.2)	86	16.7 (13.0, 20.4)	79	17.3 (13.2, 21.5)	77	16.0 (11.9, 20.1)	80	19.2 (14.6, 23.8)
	No	530	83.7 (79.8, 87.5)	536	83.3 (79.6, 87.0)	507	82.7 (78.5, 86.8)	503	84.0 (79.9, 88.1)	439	80.8 (76.2, 85.4)
NH-Other	Yes	121	13.0 (10.1, 15.9)	121	16.0 (12.0, 20.0)	97	15.1 (10.7, 19.4)	62	12.2 (8.0, 16.4)	56	11.6 (7.5, 15.6)
	No	703	87.0 (84.1, 89.9)	692	84.0 (80.0, 88.0)	597	84.9 (80.6, 89.3)	533	87.8 (83.6, 92.0)	477	88.4 (84.4, 92.5)
Hispanic	Yes	26	5.7 (3.2, 8.2)	24	4.9 (2.6, 7.2)	19	4.6 (2.1, 7.1)	26	6.9 (3.8, 10.0)	33	9.8 (5.9, 13.7)
	No	481	94.3 (91.8, 96.8)	491	95.1 (92.8, 97.4)	447	95.4 (92.9, 97.9)	377	93.1 (90.0, 96.2)	348	90.2 (86.3, 94.1)

Race/Ethnicity	Smokeless Tobacco Use	2016-18		2017-19		2018-20		2019-21		2020-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
NH-White	Yes	589	12.7 (11.5, 13.9)	632	13.4 (12.2, 14.6)	612	13.4 (12.2, 14.6)	642	14.1 (12.8, 15.4)	587	13.0 (11.8, 14.2)
	No	4994	87.3 (86.1, 88.5)	4799	86.6 (85.4, 87.8)	4365	86.6 (85.4, 87.8)	4382	85.9 (84.6, 87.2)	4280	87.0 (85.8, 88.2)
NH-Black	Yes	14	3.6 (1.2, 6.0)	19	4.4 (1.9, 6.9)	18	5.9 (2.8, 9.1)	16	5.2 (2.3, 8.1)	11	4.4 (1.6, 7.3)
	No	358	96.4 (94.0, 98.8)	348	95.6 (93.1, 98.1)	317	94.1 (90.9, 97.2)	320	94.8 (91.9, 97.7)	298	95.6 (92.7, 98.4)
NH-AI/AN	Yes	72	18.6 (13.8, 23.3)	86	21.4 (16.5, 26.2)	85	20.0 (15.2, 24.7)	92	19.2 (14.9, 23.5)	77	16.6 (12.5, 20.7)
	No	405	81.4 (76.7, 86.2)	399	78.6 (73.8, 83.5)	408	80.0 (75.3, 84.8)	430	80.8 (76.5, 85.1)	416	83.4 (79.3, 87.5)
NH-Other	Yes	58	12.3 (8.2, 16.4)	75	14.0 (9.9, 18.1)	60	11.2 (7.6, 14.7)	58	9.1 (6.4, 11.9)	52	10.0 (7.0, 13.0)
	No	485	87.7 (83.6, 91.8)	461	86.0 (81.9, 90.1)	431	88.8 (85.3, 92.4)	432	90.9 (88.1, 93.6)	445	90.0 (87.0, 93.0)
Hispanic	Yes	38	10.9 (7.0, 14.7)	44	10.9 (7.2, 14.6)	36	8.8 (5.6, 11.9)	32	6.9 (4.1, 9.6)	30	6.0 (3.5, 8.5)
	No	350	89.1 (85.3, 93.0)	395	89.1 (85.4, 92.8)	385	91.2 (88.1, 94.4)	417	93.1 (90.4, 95.9)	429	94.0 (91.5, 96.5)

Race/Ethnicity	Smokeless Tobacco Use	2021-23	
		N	Weighted% (95%CI)
NH-White	Yes	601	12.4 (11.3, 13.6)
	No	4798	87.6 (86.4, 88.7)
NH-Black	Yes	14	3.7 (1.6, 5.9)
	No	324	96.3 (94.1, 98.4)
NH-AI/AN	Yes	78	14.7 (11.3, 18.2)
	No	445	85.3 (81.8, 88.7)
NH-Other	Yes	64	10.9 (8.0, 13.8)
	No	529	89.1 (86.2, 92.0)
Hispanic	Yes	37	5.4 (3.5, 7.4)
	No	521	94.6 (92.6, 96.5)

Table 5. Three-year rolling weighted percentages of ST users among Oklahoma male adults by education, Oklahoma BRFSS 2011 – 2023

Education	Smokeless Tobacco Use	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than high school	Yes	138	13.1 (10.2, 15.9)	110	12.0 (9.3, 14.8)	91	12.0 (8.9, 15.1)	76	12.8 (9.4, 16.2)	83	14.7 (11.0, 18.3)
	No	964	86.9 (84.1, 89.8)	879	88.0 (85.2, 90.7)	721	88.0 (84.9, 91.1)	591	87.2 (83.8, 90.6)	504	85.3 (81.7, 89.0)
High school	Yes	402	15.5 (13.8, 17.3)	399	15.1 (13.5, 16.7)	360	14.8 (13.0, 16.5)	326	15.1 (13.2, 16.9)	317	15.9 (13.9, 17.9)
	No	2529	84.5 (82.7, 86.2)	2657	84.9 (83.3, 86.5)	2494	85.2 (83.5, 87.0)	2310	84.9 (83.1, 86.8)	1989	84.1 (82.1, 86.1)
Some college	Yes	281	12.6 (11.0, 14.3)	305	13.1 (11.5, 14.7)	294	13.4 (11.6, 15.1)	289	12.6 (11.0, 14.3)	258	11.6 (10.0, 13.3)
	No	2319	87.4 (85.7, 89.0)	2347	86.9 (85.3, 88.5)	2302	86.6 (84.9, 88.4)	2262	87.4 (85.7, 89.0)	2208	88.4 (86.7, 90.0)
College graduate	Yes	205	7.6 (6.4, 8.8)	215	7.2 (6.1, 8.3)	200	7.2 (6.0, 8.4)	190	6.9 (5.7, 8.1)	177	7.7 (6.3, 9.0)
	No	2768	92.4 (91.2, 93.6)	2866	92.8 (91.7, 93.9)	2782	92.8 (91.6, 94.0)	2685	93.1 (91.9, 94.3)	2461	92.3 (91.0, 93.7)
Education	Smokeless Tobacco Use	2016-18		2017-19		2018-20		2019-21		202-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than high school	Yes	85	15.6 (11.8, 19.4)	87	15.4 (11.7, 19.2)	78	14.3 (10.7, 18.0)	70	12.3 (9.0, 15.6)	59	10.0 (7.1, 12.9)
	No	474	84.4 (80.6, 88.2)	491	84.6 (80.8, 88.3)	461	85.7 (82.0, 89.3)	452	87.7 (84.4, 91.0)	423	90.0 (87.1, 92.9)
High school	Yes	306	16.6 (14.5, 18.6)	341	16.7 (14.7, 18.6)	322	16.0 (14.0, 17.9)	330	16.2 (14.2, 18.2)	301	15.8 (13.9, 17.8)
	No	1782	83.4 (81.4, 85.5)	1787	83.3 (81.4, 85.3)	1674	84.0 (82.1, 86.0)	1664	83.8 (81.8, 85.8)	1552	84.2 (82.2, 86.1)
Some college	Yes	226	10.5 (8.9, 12.0)	254	12.5 (10.8, 14.2)	240	12.6 (10.9, 14.4)	254	13.0 (11.2, 14.8)	218	11.0 (9.4, 12.7)
	No	2048	89.5 (88.0, 91.1)	1900	87.5 (85.8, 89.2)	1716	87.4 (85.6, 89.1)	1770	87.0 (85.2, 88.8)	1823	89.0 (87.3, 90.6)
College graduate	Yes	168	7.6 (6.2, 8.9)	185	8.2 (6.9, 9.6)	183	8.2 (6.8, 9.6)	196	8.0 (6.7, 9.3)	187	7.6 (6.3, 8.8)
	No	2359	92.4 (91.1, 93.8)	2318	91.8 (90.4, 93.1)	2135	91.8 (90.4, 93.2)	2203	92.0 (90.7, 93.3)	2170	92.4 (91.2, 93.7)

Education	Smokeless Tobacco Use	2021-23	
		N	Weighted% (95%CI)
Less than high school	Yes	55	8.4 (5.9, 10.9)
	No	451	91.6 (89.1, 94.1)
High school	Yes	304	15.0 (13.2, 16.8)
	No	1728	85.0 (83.2, 86.8)
Some college	Yes	244	11.1 (9.6, 12.7)
	No	2013	88.9 (87.3, 90.4)
College graduate	Yes	201	6.8 (5.7, 7.9)
	No	2550	93.2 (92.1, 94.3)

Table 6. Three-year rolling weighted percentages of ST users among Oklahoma male adults by income, Oklahoma BRFSS 2011 – 2023

Income	Smokeless Tobacco Use	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than 15,000	Yes	123	12.8 (9.9, 15.8)	114	13.1 (10.2, 15.9)	99	13.3 (10.1, 16.5)	73	12.5 (9.1, 16.0)	54	10.4 (7.2, 13.7)
	No	974	87.2 (84.2, 90.1)	877	86.9 (84.1, 89.8)	732	86.7 (83.5, 89.9)	591	87.5 (84.0, 90.9)	524	89.6 (86.3, 92.8)
15,000 to Less than 25,000	Yes	162	11.3 (9.1, 13.6)	155	11.2 (8.9, 13.4)	126	10.8 (8.4, 13.2)	122	11.6 (8.9, 14.2)	119	14.4 (11.2, 17.5)
	No	1391	88.7 (86.4, 90.9)	1373	88.8 (86.6, 91.1)	1249	89.2 (86.8, 91.6)	1099	88.4 (85.8, 91.1)	913	85.6 (82.5, 88.8)
25,000 to Less than 35,000	Yes	97	10.2 (7.8, 12.6)	107	12.6 (9.9, 15.2)	93	11.3 (8.6, 14.1)	95	13.3 (10.0, 16.5)	78	12.7 (9.3, 16.1)
	No	967	89.8 (87.4, 92.2)	975	87.4 (84.8, 90.1)	887	88.7 (85.9, 91.4)	803	86.7 (83.5, 90.0)	710	87.3 (83.9, 90.7)
35,000 to Less than 50,000	Yes	151	13.2 (10.8, 15.7)	159	13.9 (11.5, 16.4)	146	14.5 (11.8, 17.1)	131	14.3 (11.5, 17.1)	122	14.3 (11.3, 17.2)
	No	1236	86.8 (84.3, 89.2)	1231	86.1 (83.6, 88.5)	1154	85.5 (82.9, 88.2)	1090	85.7 (82.9, 88.5)	968	85.7 (82.8, 88.7)
50,000 or more	Yes	410	14.1 (12.6, 15.6)	413	12.7 (11.3, 14.0)	393	13.2 (11.7, 14.7)	361	12.5 (11.0, 14.0)	349	13.2 (11.5, 14.8)
	No	3174	85.9 (84.4, 87.4)	3395	87.3 (86.0, 88.7)	3270	86.8 (85.3, 88.3)	3142	87.5 (86.0, 89.0)	2818	86.8 (85.2, 88.5)

Income	Smokeless Tobacco Use	2016-18		2017-19		2018-20		2019-21		2020-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than 15,000	Yes	52	12.1 (8.3, 15.8)	58	12.6 (8.9, 16.4)	57	13.8 (9.7, 17.9)	47	10.9 (7.3, 14.4)	34	8.7 (5.4, 12.1)
	No	437	87.9 (84.2, 91.7)	411	87.4 (83.6, 91.1)	364	86.2 (82.1, 90.3)	356	89.1 (85.6, 92.7)	302	91.3 (87.9, 94.6)
15,000 to Less than 25,000	Yes	112	13.6 (10.6, 16.5)	126	15.5 (12.4, 18.6)	106	12.6 (9.8, 15.5)	94	12.8 (9.7, 15.9)	66	10.0 (7.1, 12.8)
	No	844	86.4 (83.5, 89.4)	805	84.5 (81.4, 87.6)	755	87.4 (84.5, 90.2)	664	87.2 (84.1, 90.3)	559	90.0 (87.2, 92.9)
25,000 to Less than 35,000	Yes	72	13.4 (9.8, 16.9)	65	11.8 (8.5, 15.1)	64	11.7 (8.5, 15.0)	66	10.2 (7.5, 12.9)	72	9.2 (6.8, 11.5)
	No	637	86.6 (83.1, 90.2)	615	88.2 (84.9, 91.5)	555	88.3 (85.0, 91.5)	610	89.8 (87.1, 92.5)	647	90.8 (88.5, 93.2)
35,000 to Less than 50,000	Yes	102	12.3 (9.6, 15.1)	111	11.9 (9.4, 14.5)	108	12.4 (9.6, 15.1)	129	15.7 (12.6, 18.9)	112	15.1 (11.9, 18.3)
	No	882	87.7 (84.9, 90.4)	851	88.1 (85.5, 90.6)	797	87.6 (84.9, 90.4)	777	84.3 (81.1, 87.4)	737	84.9 (81.7, 88.1)
50,000 or more	Yes	331	12.9 (11.3, 14.5)	392	14.5 (12.8, 16.2)	377	14.0 (12.4, 15.7)	397	13.7 (12.1, 15.2)	374	12.5 (11.1, 13.9)
	No	2764	87.1 (85.5, 88.7)	2828	85.5 (83.8, 87.2)	2675	86.0 (84.3, 87.6)	2722	86.3 (84.8, 87.9)	2784	87.5 (86.1, 88.9)

Income	Smokeless Tobacco Use	2021-23	
		N	Weighted% (95%CI)
Less than 15,000	Yes	30	7.8 (4.7, 10.8)
	No	273	92.2 (89.2, 95.3)
15,000 to Less than 25,000	Yes	52	8.5 (5.8, 11.2)
	No	489	91.5 (88.8, 94.2)
25,000 to Less than 35,000	Yes	73	8.3 (6.2, 10.4)
	No	676	91.7 (89.6, 93.8)
35,000 to Less than 50,000	Yes	127	13.5 (10.8, 16.1)
	No	884	86.5 (83.9, 89.2)
50,000 or more	Yes	426	12.2 (10.9, 13.5)
	No	3417	87.8 (86.5, 89.1)

Table 7. Proportion of ST users among Oklahoma male adults by area of residence, Oklahoma BRFSS 2011 – 2023

Area of Residence	Smokeless Tobacco Use	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Urban	Yes	86	8.0 (6.0, 10.1)	84	6.8 (5.1, 8.5)	66	6.1 (4.3, 7.8)	78	7.7 (5.7, 9.6)	65	8.6 (6.1, 11.1)
	No	1269	92.0 (89.9, 94.0)	1059	93.2 (91.5, 94.9)	1232	93.9 (92.2, 95.7)	1144	92.3 (90.4, 94.3)	905	91.4 (88.9, 93.9)
Rural	Yes	259	17.2 (14.6, 19.8)	277	16.8 (14.5, 19.0)	254	16.0 (13.8, 18.1)	272	16.2 (14.0, 18.4)	212	15.4 (12.9, 17.9)
	No	1554	82.8 (80.2, 85.4)	1793	83.2 (81.0, 85.5)	1683	84.0 (81.9, 86.2)	1851	83.8 (81.6, 86.0)	1499	84.6 (82.1, 87.1)

Area of Residence	Smokeless Tobacco Use	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Urban	Yes	51	6.6 (4.4, 8.7)	61	8.4 (5.9, 10.8)	44	5.7 (3.9, 7.6)	64	8.7 (6.3, 11.1)	55	7.9 (5.5, 10.3)
	No	903	93.4 (91.3, 95.6)	880	91.6 (89.2, 94.1)	744	94.3 (92.4, 96.1)	812	91.3 (88.9, 93.7)	673	92.1 (89.7, 94.5)
Rural	Yes	205	14.9 (12.5, 17.3)	241	17.6 (15.1, 20.1)	183	17.0 (14.3, 19.8)	275	17.8 (15.4, 20.2)	205	16.0 (13.6, 18.5)
	No	1560	85.1 (82.7, 87.5)	1429	82.4 (79.9, 84.9)	1164	83.0 (80.2, 85.7)	1486	82.2 (79.8, 84.6)	1129	84.0 (81.5, 86.4)

Area of Residence	Smokeless Tobacco Use	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Urban	Yes	45	7.3 (4.9, 9.8)	59	7.3 (5.3, 9.2)	72	6.9 (5.1, 8.6)
	No	611	92.7 (90.2, 95.1)	795	92.7 (90.8, 94.7)	992	93.1 (91.4, 94.9)
Rural	Yes	212	15.4 (13.1, 17.7)	194	13.0 (11.0, 15.0)	225	13.8 (11.8, 15.8)
	No	1406	84.6 (82.3, 86.9)	1393	87.0 (85.0, 89.0)	1584	86.2 (84.2, 88.2)

Table 8. Proportion of ST users among Oklahoma male adults by physical health status, Oklahoma BRFSS 2011 – 2023

Poor Physical Health	Smokeless Tobacco Use	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Yes	56	12.0 (8.1, 16)	48	12.2 (8.1, 16.3)	51	10.8 (7.1, 14.4)	50	10.6 (7.2, 14.0)	48	12.4 (8.1, 16.8)
	No	467	88.0 (84, 91.9)	425	87.8 (83.7, 91.9)	426	89.2 (85.6, 92.9)	454	89.4 (86.0, 92.8)	364	87.6 (83.2, 91.9)
No	Yes	289	13.6 (11.6, 15.5)	313	12.7 (11.0, 14.3)	269	11.9 (10.3, 13.5)	300	12.9 (11.2, 14.6)	229	12.5 (10.6, 14.5)
	No	2356	86.4 (84.5, 88.4)	2427	87.3 (85.7, 89.0)	2489	88.1 (86.5, 89.7)	2541	87.1 (85.4, 88.8)	2040	87.5 (85.5, 89.4)

Poor Physical Health	Smokeless Tobacco Use	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Yes	41	11.6 (7.6, 15.5)	42	14.1 (8.8, 19.3)	34	11.6 (6.4, 16.9)	43	12.3 (8.0, 16.5)	36	13.6 (8.6, 18.6)
	No	358	88.4 (84.5, 92.4)	336	85.9 (80.7, 91.2)	263	88.4 (83.1, 93.6)	328	87.7 (83.5, 92.0)	241	86.4 (81.4, 91.4)
No	Yes	212	11.4 (9.5, 13.2)	254	13.8 (11.8, 15.7)	189	12.1 (10.2, 14.0)	276	14.1 (12.2, 16.0)	214	12.3 (10.4, 14.2)
	No	2040	88.6 (86.8, 90.5)	1906	86.2 (84.3, 88.2)	1598	87.9 (86.0, 89.8)	1872	85.9 (84.0, 87.8)	1509	87.7 (85.8, 89.6)

Poor Physical Health	Smokeless Tobacco Use	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Yes	29	10.1 (5.9, 14.3)	47	14.0 (9.7, 18.3)	37	11.4 (7.0, 15.7)
	No	243	89.9 (85.7, 94.1)	303	86.0 (81.7, 90.3)	309	88.6 (84.3, 93.0)
No	Yes	220	12.3 (10.4, 14.1)	196	9.8 (8.3, 11.3)	247	10.5 (9.0, 11.9)
	No	1684	87.7 (85.9, 89.6)	1813	90.2 (88.7, 91.7)	2170	89.5 (88.1, 91.0)

Table 9. Proportion of ST users among Oklahoma male adults by mental health status, Oklahoma BRFSS 2011 – 2023

Frequent Mental Distress	Smokeless Tobacco Use	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Yes	49	14.5 (8.9, 20.1)	50	19.2 (13.4, 25.1)	40	14.7 (9.6, 19.9)	51	16.9 (11.8, 22.0)	33	13.9 (8.2, 19.6)
	No	346	85.5 (79.9, 91.1)	283	80.8 (74.9, 86.6)	301	85.3 (80.1, 90.4)	312	83.1 (78.0, 88.2)	209	86.1 (80.4, 91.8)
No	Yes	296	13.2 (11.4, 15.0)	311	11.8 (10.3, 13.3)	280	11.4 (9.9, 12.9)	299	12.1 (10.5, 13.7)	244	12.4 (10.4, 14.3)
	No	2477	86.8 (85.0, 88.6)	2569	88.2 (86.7, 89.7)	2614	88.6 (87.1, 90.1)	2683	87.9 (86.3, 89.5)	2195	87.6 (85.7, 89.6)

Frequent Mental Distress	Smokeless Tobacco Use	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Yes	29	12.7 (7.6, 17.8)	32	14.4 (8.6, 20.3)	23	9.7 (4.5, 14.9)	35	13.5 (8.6, 18.5)	28	13.0 (7.6, 18.4)
	No	245	87.3 (82.2, 92.4)	224	85.6 (79.7, 91.4)	200	90.3 (85.1, 95.5)	248	86.5 (81.5, 91.4)	211	87.0 (81.6, 92.4)
No	Yes	224	11.1 (9.3, 12.8)	262	13.5 (11.6, 15.4)	199	12.4 (10.5, 14.4)	286	13.6 (11.8, 15.5)	220	12.4 (10.5, 14.3)
	No	2170	88.9 (87.2, 90.7)	2041	86.5 (84.6, 88.4)	1672	87.6 (85.6, 89.5)	1977	86.4 (84.5, 88.2)	1530	87.6 (85.7, 89.5)

Frequent Mental Distress	Smokeless Tobacco Use	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Yes	25	10.4 (6.0, 14.8)	36	9.9 (6.4, 13.4)	44	12.5 (8.4, 16.6)
	No	224	89.6 (85.2, 94.0)	301	90.1 (86.6, 93.6)	300	87.5 (83.4, 91.6)
No	Yes	222	12.0 (10.1, 13.8)	210	10.6 (9.0, 12.2)	243	10.4 (9.0, 11.9)
	No	1734	88.0 (86.2, 89.9)	1823	89.4 (87.8, 91.0)	2191	89.6 (88.1, 91.0)

Table 10. Proportion of ST users among Oklahoma male adults by health insurance coverage, Oklahoma BRFSS 2011 – 2022

Health Insurance Status	Smokeless Tobacco Use	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Insured	Yes	282	13.3 (11.4, 15.2)	302	12.4 (10.8, 14.0)	280	12.0 (10.4, 13.6)	306	12.4 (10.8, 14.1)	249	12.7 (10.7, 14.7)
	No	2385	86.7 (84.8, 88.6)	2488	87.6 (86.0, 89.2)	2518	88.0 (86.4, 89.6)	2703	87.6 (85.9, 89.2)	2198	87.3 (85.3, 89.3)
Uninsured	Yes	61	13.6 (9.5, 17.7)	58	13.9 (9.9, 17.9)	38	10.3 (6.6, 14)	43	13.9 (9.4, 18.4)	28	12.1 (7.3, 16.9)
	No	431	86.4 (82.3, 90.5)	356	86.1 (82.1, 90.1)	391	89.7 (86.0, 93.4)	281	86.1 (81.6, 90.6)	194	87.9 (83.1, 92.7)

Health Insurance Status	Smokeless Tobacco Use	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Insured	Yes	219	11.2 (9.5, 13.0)	259	13.1 (11.3, 15.0)	192	12.2 (10.2, 14.2)	300	14.2 (12.3, 16.1)	225	12.7 (10.8, 14.6)
	No	2218	88.8 (87.0, 90.5)	2062	86.9 (85.0, 88.7)	1699	87.8 (85.8, 89.8)	2025	85.8 (83.9, 87.7)	1561	87.3 (85.4, 89.2)
Uninsured	Yes	37	13.1 (7.9, 18.2)	42	16.5 (10.9, 22.0)	34	12.3 (7.6, 17.0)	39	12.6 (8.3, 16.9)	34	12.3 (7.5, 17.2)
	No	231	86.9 (81.8, 92.1)	241	83.5 (78.0, 89.1)	200	87.7 (83.0, 92.4)	265	87.4 (83.1, 91.7)	228	87.7 (82.8, 92.5)

Health Insurance Status	Smokeless Tobacco Use	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Insured	Yes	232	12.5 (10.6, 14.3)	230	10.7 (9.2, 12.2)	268	11.0 (9.5, 12.4)
	No	1810	87.5 (85.7, 89.4)	1993	89.3 (87.8, 90.8)	2338	89.0 (87.6, 90.5)
Uninsured	Yes	22	9.6 (5.2, 14.1)	17	8.0 (3.8, 12.1)	24	10.3 (5.9, 14.8)
	No	156	90.4 (85.9, 94.8)	160	92.0 (87.9, 96.2)	177	89.7 (85.2, 94.1)

Table 11. Proportion of ST users among Oklahoma male adults by disability status, Oklahoma BRFSS 2016 – 2023

Disability Status	Smokeless Tobacco Use	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
One or more disability	Yes	84	11.3 (8.5, 14.2)	111	13.0 (10.0, 16.0)	92	13.1 (9.7, 16.4)	131	13.8 (11.1, 16.5)	103	14.2 (11.1, 17.3)
	No	879	88.7 (85.8, 91.5)	861	87.0 (84.0, 90.0)	723	86.9 (83.6, 90.3)	883	86.2 (83.5, 88.9)	646	85.8 (82.7, 88.9)
No disability	Yes	167	11.2 (9.2, 13.3)	189	14.1 (11.9, 16.4)	135	11.8 (9.6, 14.0)	206	14.0 (11.8, 16.3)	151	11.6 (9.4, 13.7)
	No	1543	88.8 (86.7, 90.8)	1417	85.9 (83.6, 88.1)	1182	88.2 (86.0, 90.4)	1384	86.0 (83.7, 88.2)	1134	88.4 (86.3, 90.6)

Disability Status	Smokeless Tobacco Use	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
One or more disability	Yes	104	13.2 (10.4, 16.1)	115	11.3 (9.0, 13.5)	117	10.8 (8.6, 12.9)
	No	746	86.8 (83.9, 89.6)	877	88.7 (86.5, 91.0)	985	89.2 (87.1, 91.4)
No disability	Yes	148	11.2 (9.1, 13.4)	132	9.8 (8.0, 11.6)	173	10.6 (8.8, 12.3)
	No	1230	88.8 (86.6, 90.9)	1292	90.2 (88.4, 92.0)	1538	89.4 (87.7, 91.2)

Section IV. Use of Electronic Vaping Products (EVP)

Table 1. Proportion of EVP users among Oklahoma adults, Oklahoma BRFSS 2017 – 2022

EVP Use	2017		2018		2019		2020		2021		2022	
	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	237	6.8 (5.8, 7.9)	280	8.7 (7.5, 9.9)	387	10.8 (9.6, 12.0)	268	9.1 (7.8, 10.3)	278	9.3 (8.0, 10.5)	344	9.3 (8.3, 10.4)
No	5326	93.2 (92.1, 94.2)	4253	91.3 (90.1, 92.5)	4977	89.2 (88.0, 90.4)	3817	90.9 (89.7, 92.2)	4094	90.7 (89.5, 92.0)	4898	90.7 (89.6, 91.7)

EVP Use	2023	
	N	Weighted% (95%CI)
Yes	350	8.5 (7.5, 9.5)
No	5585	91.5 (90.5, 92.5)

Table 2. Prevalence of current use of EVP among Oklahoma adults by sex, Oklahoma BRFSS 2017 – 2023

Sex	EVP Use	2017		2018		2019		2020		2021		2022	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Male	Yes	110	7.4 (5.8, 9.0)	131	10.4 (8.3, 12.4)	192	12.7 (10.8, 14.6)	122	10.2 (8.2, 12.2)	133	9.8 (7.9, 11.7)	159	10.4 (8.6, 12.1)
	No	2112	92.6 (91.0, 94.2)	1750	89.6 (87.6, 91.7)	2084	87.3 (85.4, 89.2)	1603	89.8 (87.8, 91.8)	1771	90.2 (88.3, 92.1)	2199	89.6 (87.9, 91.4)
Female	Yes	126	6.2 (4.8, 7.5)	149	7.1 (5.8, 8.5)	195	9.0 (7.6, 10.5)	146	8.1 (6.6, 9.5)	145	8.8 (7.1, 10.4)	185	8.4 (7.0, 9.7)
	No	3213	93.8 (92.5, 95.2)	2502	92.9 (91.5, 94.2)	2893	91.0 (89.5, 92.4)	2214	91.9 (90.5, 93.4)	2323	91.2 (89.6, 92.9)	2699	91.6 (90.3, 93.0)

Sex	EVP Use	N	2023 Weighted% (95%CI)
Male	Yes	178	10.1 (8.5, 11.8)
	No	2458	89.9 (88.2, 91.5)
Female	Yes	172	6.9 (5.7, 8.1)
	No	3127	93.1 (91.9, 94.3)

Table 3. Three-year rolling weighted percentages of EVP users among Oklahoma adults by age, Oklahoma BRFSS 2017 – 2023

Age Group	EVP Use	2017-19		2018-20		2019-21		2020-22		2021-23	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
18-24	Yes	135	20.5 (17.0, 24.0)	168	23.3 (19.8, 26.8)	190	25.3 (21.8, 28.8)	214	26.6 (23.1, 30.1)	226	26.7 (23.4, 30.1)
	No	541	79.5 (76.0, 83.0)	523	76.7 (73.2, 80.2)	547	74.7 (71.2, 78.2)	588	73.4 (69.9, 76.9)	637	73.3 (69.9, 76.6)
25-34	Yes	181	12.1 (10.1, 14.0)	210	14.3 (12.2, 16.4)	213	14.8 (12.7, 17.0)	221	15.5 (13.3, 17.6)	241	15.3 (13.3, 17.4)
	No	1348	87.9 (86.0, 89.9)	1264	85.7 (83.6, 87.8)	1255	85.2 (83.0, 87.3)	1290	84.5 (82.4, 86.7)	1465	84.7 (82.6, 86.7)
35-44	Yes	178	10.8 (9.1, 12.4)	191	11.2 (9.5, 12.8)	198	11.4 (9.7, 13.1)	177	9.8 (8.2, 11.3)	203	9.8 (8.3, 11.3)
	No	1649	89.2 (87.6, 90.9)	1565	88.8 (87.2, 90.5)	1669	88.6 (86.9, 90.3)	1729	90.2 (88.7, 91.8)	2034	90.2 (88.7, 91.7)
45-54	Yes	145	7.0 (5.7, 8.3)	133	6.8 (5.5, 8.2)	138	7.1 (5.8, 8.4)	121	6.1 (4.9, 7.3)	134	5.8 (4.7, 6.9)
	No	1945	93.0 (91.7, 94.3)	1841	93.2 (91.8, 94.5)	1844	92.9 (91.6, 94.2)	1889	93.9 (92.7, 95.1)	2169	94.2 (93.1, 95.3)
55-64	Yes	155	5.8 (4.7, 6.8)	130	5.6 (4.5, 6.8)	111	5.0 (3.9, 6.1)	88	3.6 (2.7, 4.5)	99	3.4 (2.6, 4.2)
	No	2907	94.2 (93.2, 95.3)	2519	94.4 (93.2, 95.5)	2386	95.0 (93.9, 96.1)	2367	96.4 (95.5, 97.3)	2692	96.6 (95.8, 97.4)
65 or more	Yes	110	2.3 (1.8, 2.8)	103	2.2 (1.7, 2.6)	83	1.6 (1.2, 2.0)	69	1.2 (0.9, 1.6)	69	1.2 (0.8, 1.6)
	No	6166	97.7 (97.2, 98.2)	5335	97.8 (97.4, 98.3)	5187	98.4 (98.0, 98.8)	4946	98.8 (98.4, 99.1)	5580	98.8 (98.4, 99.2)

Table 4. Three-year rolling weighted percentages of EVP users among Oklahoma adults by race/ethnicity, Oklahoma BRFSS 2017 – 2023

Race/Ethnicity	EVP Use	2017-19		2018-20		2019-21		2020-22		2021-23	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
NH-White	Yes	650	8.7 (8.0, 9.5)	652	9.5 (8.7, 10.4)	633	9.5 (8.7, 10.4)	592	8.9 (8.1, 9.7)	649	8.8 (8.0, 9.5)
	No	11043	91.3 (90.5, 92.0)	9717	90.5 (89.6, 91.3)	9445	90.5 (89.6, 91.3)	9285	91.1 (90.3, 91.9)	10539	91.2 (90.5, 92.0)
NH-Black	Yes	45	7.1 (4.6, 9.6)	45	7.1 (4.7, 9.5)	39	6.7 (4.3, 9.1)	29	5.4 (3.3, 7.5)	34	5.4 (3.4, 7.4)
	No	747	92.9 (90.4, 95.4)	666	92.9 (90.5, 95.3)	666	93.3 (90.9, 95.7)	664	94.6 (92.5, 96.7)	723	94.6 (92.6, 96.6)
NH-AI/AN	Yes	76	10.6 (7.9, 13.3)	84	11.1 (8.5, 13.6)	89	11.7 (9.2, 14.2)	96	12.0 (9.4, 14.6)	104	11.8 (9.3, 14.3)
	No	945	89.4 (86.7, 92.1)	941	88.9 (86.4, 91.5)	957	88.3 (85.8, 90.8)	965	88.0 (85.4, 90.6)	1037	88.2 (85.7, 90.7)
NH-Other	Yes	79	10.7 (8.0, 13.3)	95	13.2 (10.1, 16.3)	101	14.9 (11.5, 18.4)	98	14.4 (11.2, 17.7)	88	11.6 (8.9, 14.3)
	No	904	89.3 (86.7, 92.0)	790	86.8 (83.7, 89.9)	800	85.1 (81.6, 88.5)	825	85.6 (82.3, 88.8)	1025	88.4 (85.7, 91.1)
Hispanic	Yes	43	7.4 (4.8, 10.0)	50	7.4 (5.0, 9.7)	61	8.1 (5.9, 10.4)	62	8.1 (5.9, 10.3)	74	8.2 (6.1, 10.3)
	No	775	92.6 (90.0, 95.2)	802	92.6 (90.3, 95.0)	838	91.9 (89.6, 94.1)	864	91.9 (89.7, 94.1)	1006	91.8 (89.7, 93.9)

Table 5. Three-year rolling weighted percentages of EVP users among Oklahoma adults by education, Oklahoma BRFSS 2017 – 2023

Education	EVP Use	2017-19		2018-20		2019-21		2020-22		2021-23	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than high school	Yes	94	11.5 (8.9, 14.1)	80	10.1 (7.6, 12.6)	76	9.6 (7.3, 12.0)	71	9.5 (7.1, 11.8)	78	10.2 (7.9, 12.6)
	No	1051	88.5 (85.9, 91.1)	988	89.9 (87.4, 92.4)	942	90.4 (88.0, 92.7)	900	90.5 (88.2, 92.9)	930	89.8 (87.4, 92.1)
High school	Yes	297	10.0 (8.7, 11.3)	312	11.2 (9.9, 12.6)	314	12.1 (10.7, 13.6)	294	11.8 (10.3, 13.2)	326	11.6 (10.2, 12.9)
	No	4149	90.0 (88.7, 91.3)	3662	88.8 (87.4, 90.1)	3515	87.9 (86.4, 89.3)	3354	88.2 (86.8, 89.7)	3666	88.4 (87.1, 89.8)
Some college	Yes	339	9.6 (8.4, 10.7)	351	10.5 (9.3, 11.7)	358	10.8 (9.5, 12.0)	358	10.3 (9.1, 11.5)	391	10.2 (9.1, 11.4)
	No	4478	90.4 (89.3, 91.6)	4003	89.5 (88.3, 90.7)	3959	89.2 (88.0, 90.5)	4025	89.7 (88.5, 90.9)	4443	89.8 (88.6, 90.9)
College graduate	Yes	173	4.6 (3.9, 5.4)	192	5.6 (4.7, 6.5)	184	5.2 (4.3, 6.0)	166	4.5 (3.7, 5.3)	175	3.7 (3.1, 4.4)
	No	4854	95.4 (94.6, 96.1)	4368	94.4 (93.5, 95.3)	4436	94.8 (94.0, 95.7)	4473	95.5 (94.7, 96.3)	5478	96.3 (95.6, 96.9)

Table 6. Three-year rolling weighted percentages of EVP users among Oklahoma adults by income, Oklahoma BRFSS 2017 – 2023

Income	EVP Use	2017-19		2018-20		2019-21		2020-22		2021-23	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than 15,000	Yes	107	9.9 (7.7, 12.1)	97	11.1 (8.6, 13.6)	97	13.0 (10.1, 15.8)	82	11.5 (8.7, 14.3)	77	12.0 (8.9, 15.1)
	No	1123	90.1 (87.9, 92.3)	953	88.9 (86.4, 91.4)	872	87.0 (84.2, 89.9)	786	88.5 (85.7, 91.3)	740	88.0 (84.9, 91.1)
15,000 to Less than 25,000	Yes	168	11.4 (9.4, 13.4)	166	10.7 (8.9, 12.5)	150	10.3 (8.4, 12.1)	128	10.4 (8.4, 12.4)	111	11.0 (8.8, 13.3)
	No	2130	88.6 (86.6, 90.6)	1922	89.3 (87.5, 91.1)	1712	89.7 (87.9, 91.6)	1420	89.6 (87.6, 91.6)	1218	89.0 (86.7, 91.2)
25,000 to Less than 35,000	Yes	105	11.7 (9.1, 14.3)	111	12.8 (10.1, 15.6)	105	11.6 (9.2, 14.1)	117	10.6 (8.5, 12.7)	114	9.0 (7.2, 10.8)
	No	1412	88.3 (85.7, 90.9)	1272	87.2 (84.4, 89.9)	1281	88.4 (85.9, 90.8)	1412	89.4 (87.3, 91.5)	1547	91.0 (89.2, 92.8)
35,000 to Less than 50,000	Yes	111	8.8 (6.9, 10.6)	124	10.4 (8.4, 12.4)	128	10.5 (8.5, 12.5)	124	10.2 (8.3, 12.2)	160	10.4 (8.6, 12.1)
	No	1827	91.2 (89.4, 93.1)	1673	89.6 (87.6, 91.6)	1596	89.5 (87.5, 91.5)	1619	89.8 (87.8, 91.7)	1928	89.6 (87.9, 91.4)
50,000 or more	Yes	300	7.3 (6.3, 8.2)	311	7.8 (6.8, 8.8)	307	7.9 (6.9, 8.9)	283	7.2 (6.3, 8.2)	339	7.0 (6.2, 7.9)
	No	5546	92.7 (91.8, 93.7)	5102	92.2 (91.2, 93.2)	5124	92.1 (91.1, 93.1)	5295	92.8 (91.8, 93.7)	6691	93.0 (92.1, 93.8)

Table 7. Proportion of EVP users among Oklahoma adults by area of residence, Oklahoma BRFSS 2017 – 2023

Area of Residence	EVP Use	2017		2018		2019		2020		2021		2022	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Urban	Yes	90	7.2 (5.4, 8.9)	109	8.6 (6.7, 10.5)	162	12.2 (10.2, 14.1)	90	8.6 (6.6, 10.6)	90	9.0 (6.9, 11.1)	132	9.2 (7.5, 10.9)
	No	1843	92.8 (91.1, 94.6)	1540	91.4 (89.5, 93.3)	1642	87.8 (85.9, 89.8)	1293	91.4 (89.4, 93.4)	1200	91.0 (88.9, 93.1)	1678	90.8 (89.1, 92.5)
Rural	Yes	147	6.6 (5.3, 7.9)	171	8.8 (7.2, 10.4)	225	9.8 (8.3, 11.2)	178	9.4 (7.9, 11.0)	188	9.4 (7.9, 11.0)	212	9.5 (8.1, 10.9)
	No	3483	93.4 (92.1, 94.7)	2713	91.2 (89.6, 92.8)	3335	90.2 (88.8, 91.7)	2524	90.6 (89.0, 92.1)	2894	90.6 (89.0, 92.1)	3220	90.5 (89.1, 91.9)

Area of Residence	EVP Use	2023	
		N	Weighted% (95%CI)
Urban	Yes	133	8.4 (6.9, 9.9)
	No	2025	91.6 (90.1, 93.1)
Rural	Yes	217	8.6 (7.3, 9.9)
	No	3560	91.4 (90.1, 92.7)

Table 8. Proportion of EVP users among Oklahoma adults by physical health status, Oklahoma BRFSS 2017 – 2023

Poor Physical Health	EVP Use	2017		2018		2019		2020		2021		2022	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Yes	60	9.3 (6.4, 12.1)	65	12.7 (9.1, 16.2)	61	10.0 (7.2, 12.7)	48	11.5 (7.8, 15.2)	50	10.0 (6.8, 13.3)	63	10.3 (7.5, 13.1)
	No	834	90.7 (87.9, 93.6)	665	87.3 (83.8, 90.9)	764	90.0 (87.3, 92.8)	517	88.5 (84.8, 92.2)	563	90.0 (86.7, 93.2)	725	89.7 (86.9, 92.5)
No	Yes	171	6.4 (5.2, 7.5)	206	7.9 (6.6, 9.2)	314	11.1 (9.7, 12.4)	211	8.8 (7.4, 10.1)	218	9.2 (7.8, 10.6)	269	9.1 (7.9, 10.3)
	No	4320	93.6 (92.5, 94.8)	3482	92.1 (90.8, 93.4)	4015	88.9 (87.6, 90.3)	3159	91.2 (89.9, 92.6)	3336	90.8 (89.4, 92.2)	4014	90.9 (89.7, 92.1)

Poor Physical Health	EVP Use	2023	
		N	Weighted% (95%CI)
Yes	Yes	66	11.2 (8.2, 14.3)
	No	742	88.8 (85.7, 91.8)
No	Yes	269	7.9 (6.9, 9.0)
	No	4611	92.1 (91.0, 93.1)

Table 9. Proportion of EVP users among Oklahoma adults by mental health status, Oklahoma BRFSS 2017 – 2023

Frequent Mental Distress	EVP Use	2017		2018		2019		2020		2021		2022	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Yes	74	14.6 (10.7, 18.4)	87	16.9 (12.9, 20.8)	104	19.8 (15.9, 23.7)	86	17.9 (13.9, 21.8)	83	17.3 (13.3, 21.3)	119	19.3 (15.8, 22.8)
	No	619	85.4 (81.6, 89.3)	517	83.1 (79.2, 87.1)	587	80.2 (76.3, 84.1)	485	82.1 (78.2, 86.1)	549	82.7 (78.7, 86.7)	693	80.7 (77.2, 84.2)
No	Yes	159	5.5 (4.5, 6.6)	182	6.9 (5.7, 8.2)	265	9.0 (7.8, 10.3)	161	7.1 (5.8, 8.3)	184	7.5 (6.2, 8.8)	203	6.9 (5.8, 7.9)
	No	4587	94.5 (93.4, 95.5)	3646	93.1 (91.8, 94.3)	4210	91.0 (89.7, 92.2)	3193	92.9 (91.7, 94.2)	3412	92.5 (91.2, 93.8)	4064	93.1 (92.1, 94.2)

Frequent Mental Distress	EVP Use	2023	
		N	Weighted% (95%CI)
Yes	Yes	107	17.1 (13.5, 20.7)
	No	719	82.9 (79.3, 86.5)
No	Yes	233	6.9 (5.9, 7.8)
	No	4685	93.1 (92.2, 94.1)

Frequent Mental Distress = 14 or more days of poor mental health

Table 10. Proportion of EVP users among Oklahoma adults by health insurance coverage, Oklahoma BRFSS 2017 – 2023

Health Insurance Status	EVP Use	2017		2018		2019		2020		2021		2022	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Insured	Yes	192	6.1 (5.1, 7.1)	246	8.5 (7.2, 9.8)	300	9.6 (8.4, 10.8)	221	8.5 (7.2, 9.8)	220	8.2 (7.0, 9.5)	290	8.8 (7.7, 9.9)
	No	4898	93.9 (92.9, 94.9)	3840	91.5 (90.2, 92.8)	4479	90.4 (89.2, 91.6)	3397	91.5 (90.2, 92.8)	3754	91.8 (90.5, 93.0)	4523	91.2 (90.1, 92.3)
Uninsured	Yes	45	11.6 (7.6, 15.6)	33	10.0 (6.4, 13.7)	85	17.1 (13.4, 20.9)	47	12.8 (8.8, 16.8)	49	18.3 (12.9, 23.7)	48	14.3 (10.1, 18.4)
	No	411	88.4 (84.4, 92.4)	401	90.0 (86.3, 93.6)	488	82.9 (79.1, 86.6)	400	87.2 (83.2, 91.2)	245	81.7 (76.3, 87.1)	294	85.7 (81.6, 89.9)

Health Insurance Status	EVP Use	2023	
		N	Weighted% (95%CI)
Insured	Yes	298	8.1 (7.0, 9.1)
	No	5142	91.9 (90.9, 93.0)
Uninsured	Yes	43	12.2 (8.3, 16.2)
	No	317	87.8 (83.8, 91.7)

Table 11. Proportion of EVP users among Oklahoma adults by disability status, Oklahoma BRFSS 2017 – 2023

Disability Status	EVP Use	2017		2018		2019		2020		2021		2022	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
One or more disability	Yes	120	9.8 (7.7, 11.9)	123	10.5 (8.3, 12.7)	163	12.0 (10.0, 14.1)	118	10.2 (8.1, 12.3)	122	10.1 (8.0, 12.1)	160	10.7 (8.8, 12.5)
	No	2101	90.2 (88.1, 92.3)	1685	89.5 (87.3, 91.7)	1987	88.0 (85.9, 90.0)	1449	89.8 (87.7, 91.9)	1586	89.9 (87.9, 92.0)	2028	89.3 (87.5, 91.2)
No disability	Yes	113	5.1 (4.0, 6.3)	157	7.8 (6.3, 9.2)	219	10.1 (8.6, 11.6)	147	8.6 (7.0, 10.1)	146	8.6 (7.0, 10.2)	178	8.5 (7.2, 9.9)
	No	3165	94.9 (93.7, 96.0)	2553	92.2 (90.8, 93.7)	2925	89.9 (88.4, 91.4)	2316	91.4 (89.9, 93.0)	2424	91.4 (89.8, 93.0)	2804	91.5 (90.1, 92.8)

Disability Status	EVP Use	2023	
		N	Weighted% (95%CI)
One or more disability	Yes	170	10.9 (9.1, 12.8)
	No	2108	89.1 (87.2, 90.9)
No disability	Yes	172	6.8 (5.7, 8.0)
	No	3366	93.2 (92.0, 94.3)

Section V. Past quit attempts (in the past 12 months)

Table 1. Past quit attempts among Oklahoma adult current smokers, Oklahoma BRFSS 2011 – 2023

Quit attempts in the last 12 months	2011		2012		2013		2014		2015	
	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	1075	58.5 (55.4, 61.7)	950	61.5 (58.4, 64.6)	892	58.2 (55.0, 61.4)	833	61.3 (58.1, 64.65)	622	58.2 (54.1, 62.2)
No	779	41.5 (38.3, 44.6)	680	38.5 (35.4, 41.6)	669	41.8 (38.6, 45.0)	585	38.7 (35.4, 41.9)	523	41.9 (37.8, 45.9)

Quit attempts in the last 12 months	2016		2017		2018		2019		2020	
	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	530	57.5 (53.4, 61.5)	565	59.4 (55.6, 63.2)	430	54.3 (50.1, 58.5)	526	58.0 (54.1, 61.8)	397	55.2 (50.9, 59.4)
No	474	42.5 (38.5, 46.6)	441	40.6 (36.8, 44.4)	403	45.7 (41.5, 50.0)	434	42.0 (38.2, 45.9)	394	44.8 (40.6, 49.1)

Quit attempts in the last 12 months	2021		2022		2023	
	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	311	51.8 (47.1, 56.5)	324	47.0 (42.6, 51.5)	310	51.5 (46.8, 56.2)
No	343	48.2 (43.5, 52.9)	353	53.0 (48.5, 57.4)	335	48.5 (43.8, 53.2)

Table 2. Past quit attempts among Oklahoma adult current smokers by gender, Oklahoma BRFSS 2011 – 2023

Gender	Quit attempts in the last 12 months	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Male	Yes	397	54.1 (49.2, 59.0)	374	58.8 (54.0, 63.5)	358	54.5 (49.9, 59.1)	355	58.5 (53.7, 63.4)	246	55.1 (49.0, 61.2)
	No	354	45.9 (41.0, 50.8)	306	41.2 (36.5, 46.0)	329	45.5 (40.9, 50.1)	286	41.5 (36.6, 46.3)	258	44.9 (38.8, 51)
Female	Yes	678	63.4 (59.6, 67.3)	576	64.4 (60.4, 68.3)	534	62.4 (58.1, 66.7)	478	64.6 (60.1, 69.0)	376	61.6 (56.4, 66.8)
	No	425	36.6 (32.7, 40.4)	374	35.6 (31.7, 39.6)	340	37.6 (33.3, 41.9)	299	35.4 (31.0, 39.9)	265	38.4 (33.2, 43.6)

Gender	Quit attempts in the last 12 months	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Male	Yes	223	56.1 (50.1, 62.1)	251	57.9 (52.2, 63.6)	203	52.0 (46.0, 58.0)	248	58.4 (53.0, 63.8)	178	53.6 (47.2, 60.0)
	No	218	43.9 (37.9, 49.9)	194	42.1 (36.4, 47.8)	201	48.0 (42.0, 54.0)	214	41.6 (36.2, 47.0)	187	46.4 (40.0, 52.8)
Female	Yes	307	59.0 (53.7, 64.3)	314	61.0 (56.0, 66.0)	227	57.2 (51.3, 63.0)	278	57.4 (51.9, 62.9)	219	56.9 (51.4, 62.4)
	No	256	41.0 (35.7, 46.3)	247	39.0 (34.0, 44.0)	202	42.8 (37.0, 48.7)	220	42.6 (37.1, 48.1)	207	43.1 (37.6, 48.6)

Gender	Quit attempts in the last 12 months	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Male	Yes	125	49.9 (42.6, 57.1)	146	43.0 (36.6, 49.3)	123	48.3 (41.3, 55.3)
	No	162	50.1 (42.9, 57.4)	181	57.0 (50.7, 63.4)	158	51.7 (44.7, 58.7)
Female	Yes	186	53.5 (47.4, 59.7)	178	51.7 (45.5, 57.8)	187	54.5 (48.3, 60.8)
	No	181	46.5 (40.3, 52.6)	172	48.3 (42.2, 54.5)	177	45.5 (39.2, 51.7)

Table 3. Past quit attempts among Oklahoma adult current smokers by age (three-year rolling weighted percentages), Oklahoma BRFSS 2011 – 2023

Age Group	Quit attempts in the last 12 months	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
18-24	Yes	204	68.1 (61.6, 74.5)	211	71.7 (65.6, 77.8)	172	74.1 (67.6, 80.5)	147	77.3 (70.7, 83.9)	117	74.7 (67.3, 82.0)
	No	92	31.9 (25.5, 38.4)	83	28.3 (22.2, 34.4)	64	25.9 (19.5, 32.4)	48	22.7 (16.1, 29.3)	45	25.3 (18.0, 32.7)
25-34	Yes	470	65.8 (61.7, 69.8)	432	65.3 (61.0, 69.6)	363	62.7 (57.9, 67.6)	309	62.2 (57.0, 67.3)	288	62.6 (57.3, 67.8)
	No	259	34.2 (30.2, 38.3)	230	34.7 (30.4, 39.0)	203	37.3 (32.4, 42.1)	186	37.8 (32.7, 43.0)	176	37.4 (32.2, 42.7)
35-44	Yes	454	58.2 (53.9, 62.5)	418	57.4 (53.0, 61.8)	344	55.3 (50.4, 60.1)	271	54.4 (49.0, 59.9)	235	55.1 (49.4, 60.7)
	No	304	41.8 (37.5, 46.1)	270	42.6 (38.2, 47.0)	249	44.7 (39.9, 49.6)	205	45.6 (40.1, 51.0)	188	44.9 (39.3, 50.6)
45-54	Yes	643	54.4 (50.9, 57.8)	561	56.8 (53.3, 60.4)	480	56.0 (51.7, 60.2)	383	54.8 (50.2, 59.5)	314	54.2 (49.2, 59.2)
	No	510	45.6 (42.2, 49.1)	415	43.2 (39.6, 46.7)	341	44.0 (39.8, 48.3)	290	45.2 (40.5, 49.8)	257	45.8 (40.8, 50.8)
55-64	Yes	664	54.8 (51.2, 58.4)	629	55.7 (52.1, 59.2)	579	56.0 (52.3, 59.7)	496	55.3 (51.3, 59.4)	431	54.6 (50.3, 58.9)
	No	512	45.2 (41.6, 48.8)	494	44.3 (40.8, 47.9)	461	44.0 (40.3, 47.7)	407	44.7 (40.6, 48.7)	353	45.4 (41.1, 49.7)
65 or more	Yes	482	50.2 (46.1, 54.2)	424	49.7 (45.6, 53.8)	409	47.8 (43.6, 52.1)	379	48.2 (43.8, 52.6)	332	46.5 (41.9, 51.0)
	No	451	49.8 (45.8, 53.9)	442	50.3 (46.2, 54.4)	459	52.2 (47.9, 56.4)	446	51.8 (47.4, 56.2)	419	53.5 (49.0, 58.1)

Age Group	Quit attempts in the last 12 months	2016-18		2017-19		2018-20		2019-21		2020-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
18-24	Yes	103	68.8 (60.6, 77.0)	102	67.1 (58.6, 75.6)	76	64.8 (55.1, 74.4)	67	67.5 (57.6, 77.4)	44	64.1 (52.5, 75.7)
	No	51	31.2 (23.0, 39.4)	48	32.9 (24.4, 41.4)	44	35.2 (25.6, 44.9)	35	32.5 (22.6, 42.4)	32	35.9 (24.3, 47.5)
25-34	Yes	266	65.3 (60.2, 70.4)	257	66.8 (61.6, 72.0)	219	67.0 (61.5, 72.5)	187	66.2 (60.2, 72.2)	149	61.2 (54.4, 68.0)
	No	161	34.7 (29.6, 39.8)	145	33.2 (28.0, 38.4)	126	33.0 (27.5, 38.5)	111	33.8 (27.8, 39.8)	104	38.8 (32.0, 45.6)
35-44	Yes	222	54.3 (48.6, 59.9)	246	54.5 (49.2, 59.7)	238	55.9 (50.5, 61.3)	236	55.8 (50.4, 61.2)	197	52.8 (46.9, 58.6)
	No	181	45.7 (40.1, 51.4)	203	45.5 (40.3, 50.8)	191	44.1 (38.7, 49.5)	193	44.2 (38.8, 49.6)	178	47.2 (41.4, 53.1)
45-54	Yes	287	52.3 (47.3, 57.2)	281	53.9 (48.9, 58.9)	247	49.4 (44.3, 54.6)	211	49.4 (43.9, 54.9)	173	45.3 (39.5, 51.0)
	No	247	47.7 (42.8, 52.7)	234	46.1 (41.1, 51.1)	248	50.6 (45.4, 55.7)	228	50.6 (45.1, 56.1)	214	54.7 (49.0, 60.5)
55-64	Yes	355	52.2 (47.6, 56.8)	342	52.5 (47.8, 57.2)	298	52.1 (47.1, 57.0)	286	52.3 (47.3, 57.3)	245	48.9 (43.7, 54.0)
	No	312	47.8 (43.2, 52.4)	290	47.5 (42.8, 52.2)	279	47.9 (43.0, 52.9)	264	47.7 (42.7, 52.7)	260	51.1 (46.0, 56.3)
65 or more	Yes	292	46.2 (41.4, 50.9)	293	46.0 (41.4, 50.6)	275	45.5 (40.8, 50.3)	247	44.0 (39.0, 49.0)	224	44.4 (39.1, 49.7)
	No	366	53.8 (49.1, 58.6)	358	54.0 (49.4, 58.6)	343	54.5 (49.7, 59.2)	340	56.0 (51.0, 61.0)	302	55.6 (50.3, 60.9)

Age Group	Quit attempts in the last 12 months	2021-23	
		N	Weighted% (95%CI)
18-24	Yes	38	60.6 (48.2, 73.0)
	No	31	39.4 (27.0, 51.8)
25-34	Yes	124	57.8 (50.3, 65.3)
	No	93	42.2 (34.7, 49.7)
35-44	Yes	181	50.4 (44.4, 56.3)
	No	174	49.6 (43.7, 55.6)
45-54	Yes	165	48.4 (42.3, 54.5)
	No	184	51.6 (45.5, 57.7)
55-64	Yes	225	47.3 (42.1, 52.5)
	No	255	52.7 (47.5, 57.9)
65 or more	Yes	212	43.0 (37.5, 48.4)
	No	294	57.0 (51.6, 62.5)

Table 4. Past quit attempts among Oklahoma adult current smokers by race/ethnicity (three-year rolling weighted percentages), Oklahoma BRFSS 2011 – 2023

Race/Ethnicity	Quit attempts in the last 12 months	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
NH-White	Yes	1973	58.6 (56.4, 60.8)	1822	58.3 (56.1, 60.5)	1615	58.0 (55.6, 60.4)	1385	57.5 (54.9, 60.1)	1187	57.1 (54.4, 59.8)
	No	1538	41.4 (39.2, 43.6)	1407	41.7 (39.5, 43.9)	1283	42.0 (39.6, 44.4)	1191	42.5 (39.9, 45.1)	1085	42.9 (40.2, 45.6)
NH-Black	Yes	264	67.2 (60.8, 73.7)	220	70.6 (64.4, 76.8)	197	69.3 (62.6, 76.0)	150	66.5 (58.4, 74.6)	127	67.2 (58.5, 75.9)
	No	123	32.8 (26.3, 39.2)	97	29.4 (23.2, 35.6)	96	30.7 (24.0, 37.4)	75	33.5 (25.4, 41.6)	61	32.8 (24.1, 41.5)
NH-AI/AN	Yes	256	57.7 (51.6, 63.8)	231	57.3 (51.1, 63.6)	212	54.6 (48.1, 61.1)	191	56.5 (49.7, 63.3)	176	57.8 (50.8, 64.9)
	No	194	42.3 (36.2, 48.4)	182	42.7 (36.4, 48.9)	189	45.4 (38.9, 51.9)	154	43.5 (36.7, 50.3)	144	42.2 (35.1, 49.2)
NH-Other	Yes	279	65.2 (58.8, 71.6)	269	71.5 (65.5, 77.5)	207	65.6 (57.9, 73.2)	161	67.0 (58.8, 75.1)	121	57.6 (48.3, 66.9)
	No	173	34.8 (28.4, 41.2)	166	28.5 (22.5, 34.5)	141	34.4 (26.8, 42.1)	104	33.0 (24.9, 41.2)	88	42.4 (33.1, 51.7)
Hispanic	Yes	137	56.6 (48.4, 64.8)	125	60.7 (52.5, 68.9)	103	58.0 (47.5, 68.4)	81	60.0 (48.8, 71.2)	82	61.0 (50.2, 71.8)
	No	91	43.4 (35.2, 51.6)	74	39.3 (31.1, 47.5)	59	42.0 (31.6, 52.5)	47	40.0 (28.8, 51.2)	49	39.0 (28.2, 49.8)

Race/Ethnicity	Quit attempts in the last 12 months	2016-18		2017-19		2018-20		2019-21		2020-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
NH-White	Yes	1049	55.9 (53.1, 58.6)	1013	55.0 (52.2, 57.8)	912	54.9 (52.1, 57.8)	809	52.5 (49.5, 55.4)	686	50.2 (47.1, 53.3)
	No	991	44.1 (41.4, 46.9)	940	45.0 (42.2, 47.8)	875	45.1 (42.2, 47.9)	835	47.5 (44.6, 50.5)	777	49.8 (46.7, 52.9)
NH-Black	Yes	106	60.6 (51.4, 69.9)	112	68.1 (59.8, 76.5)	99	67.1 (58.5, 75.7)	95	74.9 (66.8, 82.9)	76	68.4 (58.9, 78.0)
	No	61	39.4 (30.1, 48.6)	53	31.9 (23.5, 40.2)	52	32.9 (24.3, 41.5)	37	25.1 (17.1, 33.2)	37	31.6 (22.0, 41.1)
NH-AI/AN	Yes	139	56.0 (48.3, 63.7)	152	56.9 (49.7, 64.1)	139	53.6 (46.0, 61.2)	142	52.9 (45.6, 60.2)	118	50.9 (43.0, 58.9)
	No	116	44.0 (36.3, 51.7)	127	43.1 (35.9, 50.3)	136	46.4 (38.8, 54.0)	146	47.1 (39.8, 54.4)	127	49.1 (41.1, 57.0)
NH-Other	Yes	130	56.2 (47.6, 64.9)	140	57.5 (48.9, 66.1)	123	55.0 (46.2, 63.8)	100	60.9 (51.9, 69.9)	78	47.1 (37.5, 56.7)
	No	92	43.8 (35.1, 52.4)	95	42.5 (33.9, 51.1)	97	45.0 (36.2, 53.8)	84	39.1 (30.1, 48.1)	90	52.9 (43.3, 62.5)
Hispanic	Yes	80	65.0 (55.1, 74.8)	85	66.5 (57.0, 76.0)	67	56.8 (46.6, 67.0)	69	59.7 (49.4, 70.0)	58	55.9 (44.2, 67.6)
	No	49	35.0 (25.2, 44.9)	51	33.5 (24.0, 43.0)	58	43.2 (33.0, 53.4)	55	40.3 (30.0, 50.6)	45	44.1 (32.4, 55.8)

Race/Ethnicity	Quit attempts in the last 12 months	2021-23	
		N	Weighted% (95%CI)
NH-White	Yes	629	47.2 (44.1, 50.4)
	No	746	52.8 (49.6, 55.9)
NH-Black	Yes	63	68.3 (58.0, 78.7)
	No	32	31.7 (21.3, 42.0)
NH-AI/AN	Yes	101	49.2 (41.2, 57.2)
	No	113	50.8 (42.8, 58.8)
NH-Other	Yes	71	45.8 (36.5, 55.2)
	No	91	54.2 (44.8, 63.5)
Hispanic	Yes	63	65.1 (54.1, 76.1)
	No	36	34.9 (23.9, 45.9)

Table 5. Past quit attempts among Oklahoma adult current smokers by education (three-year rolling weighted percentages), Oklahoma BRFSS 2011 – 2023

Education	Quit attempts in the last 12 months	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than high school	Yes	543	60.2 (55.9, 64.5)	469	59.5 (55.0, 64.0)	359	54.6 (49.3, 59.8)	279	56.4 (50.7, 62.2)	226	57.1 (51.1, 63.2)
	No	404	39.8 (35.5, 44.1)	351	40.5 (36.0, 45.0)	307	45.4 (40.2, 50.7)	236	43.6 (37.8, 49.3)	195	42.9 (36.8, 48.9)
High school	Yes	1097	57.3 (54.3, 60.2)	1030	59.6 (56.7, 62.5)	915	61.0 (57.9, 64.1)	769	59.4 (56.0, 62.7)	630	57.2 (53.7, 60.8)
	No	832	42.7 (39.8, 45.7)	748	40.4 (37.5, 43.3)	652	39.0 (35.9, 42.1)	596	40.6 (37.3, 44.0)	559	42.8 (39.2, 46.3)
Some college	Yes	886	61.2 (58.1, 64.3)	831	62.4 (59.3, 65.5)	767	61.6 (58.3, 64.9)	669	60.8 (57.2, 64.3)	610	60.5 (56.8, 64.2)
	No	606	38.8 (35.7, 41.9)	550	37.6 (34.5, 40.7)	544	38.4 (35.1, 41.7)	506	39.2 (35.7, 42.8)	470	39.5 (35.8, 43.2)
College graduate	Yes	390	60.1 (55.3, 65.0)	345	58.4 (53.4, 63.3)	306	55.8 (50.3, 61.4)	267	58.2 (52.5, 64.0)	250	58.8 (53.0, 64.7)
	No	283	39.9 (35.0, 44.7)	283	41.6 (36.7, 46.6)	271	44.2 (38.6, 49.7)	242	41.8 (36.0, 47.5)	211	41.2 (35.3, 47.0)

Education	Quit attempts in the last 12 months	2016-18		2017-19		2018-20		2019-21		2020-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than high school	Yes	206	57.2 (51.3, 63.1)	199	54.4 (48.4, 60.3)	179	54.1 (48.0, 60.1)	162	54.3 (48.0, 60.5)	131	49.6 (42.8, 56.4)
	No	181	42.8 (36.9, 48.7)	196	45.6 (39.7, 51.6)	196	45.9 (39.9, 52.0)	179	45.7 (39.5, 52.0)	160	50.4 (43.6, 57.2)
High school	Yes	551	55.4 (51.6, 59.1)	555	55.5 (51.9, 59.2)	488	53.4 (49.5, 57.2)	431	53.1 (49.2, 57.0)	335	49.7 (45.4, 54.0)
	No	508	44.6 (40.9, 48.4)	502	44.5 (40.8, 48.1)	472	46.6 (42.8, 50.5)	444	46.9 (43.0, 50.8)	389	50.3 (46.0, 54.6)
Some college	Yes	538	59.3 (55.5, 63.1)	542	62.4 (58.7, 66.1)	483	60.5 (56.7, 64.3)	460	58.8 (55.0, 62.7)	411	55.3 (51.2, 59.3)
	No	427	40.7 (36.9, 44.5)	382	37.6 (33.9, 41.3)	384	39.5 (35.7, 43.3)	379	41.2 (37.3, 45.0)	377	44.7 (40.7, 48.8)
College graduate	Yes	229	55.8 (49.7, 61.8)	225	54.0 (48.0, 60.1)	203	54.1 (47.7, 60.5)	181	54.7 (48.1, 61.2)	155	52.5 (45.7, 59.2)
	No	199	44.2 (38.2, 50.3)	194	46.0 (39.9, 52.0)	176	45.9 (39.5, 52.3)	165	45.3 (38.8, 51.9)	162	47.5 (40.8, 54.3)

Education	Quit attempts in the last 12 months	2021-23	
		N	Weighted% (95%CI)
Less than high school	Yes	128	48.8 (41.8, 55.7)
	No	138	51.2 (44.3, 58.2)
High school	Yes	305	50.0 (45.5, 54.5)
	No	354	50.0 (45.5, 54.5)
Some college	Yes	366	52.4 (48.2, 56.6)
	No	355	47.6 (43.4, 51.8)
College graduate	Yes	146	45.1 (38.9, 51.3)
	No	182	54.9 (48.7, 61.1)

Table 6. Past quit attempts among Oklahoma adult current smokers by income (three-year rolling weighted percentages), Oklahoma BRFSS
2011 – 2023

Income	Quit attempts in the last 12 months	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than 15,000	Yes	843	66.3 (62.8, 69.8)	718	65.7 (62.0, 69.4)	589	64.1 (59.9, 68.3)	427	63.4 (58.5, 68.3)	346	64.2 (59.0, 69.3)
	No	467	33.7 (30.2, 37.2)	396	34.3 (30.6, 38.0)	336	35.9 (31.7, 40.1)	269	36.6 (31.7, 41.5)	227	35.8 (30.7, 41.0)
15,000 to Less than 25,000	Yes	698	60.7 (56.9, 64.5)	640	60.9 (57.0, 64.8)	544	56.3 (51.9, 60.7)	456	57.2 (52.5, 61.9)	381	58.3 (53.3, 63.3)
	No	500	39.3 (35.5, 43.1)	463	39.1 (35.2, 43.0)	431	43.7 (39.3, 48.1)	364	42.8 (38.1, 47.5)	307	41.7 (36.7, 46.7)
25,000 to Less than 35,000	Yes	340	59.5 (54.4, 64.5)	323	62.1 (56.9, 67.2)	259	63.3 (57.6, 69.0)	224	62.6 (56.4, 68.8)	183	57.5 (50.7, 64.3)
	No	262	40.5 (35.5, 45.6)	231	37.9 (32.8, 43.1)	194	36.7 (31.0, 42.4)	176	37.4 (31.2, 43.6)	163	42.5 (35.7, 49.3)
35,000 to Less than 50,000	Yes	322	59.0 (54.1, 63.9)	302	61.6 (56.6, 66.5)	276	60.0 (54.6, 65.5)	247	62.5 (56.4, 68.6)	212	57.2 (50.8, 63.6)
	No	268	41.0 (36.1, 45.9)	230	38.4 (33.5, 43.4)	206	40.0 (34.5, 45.4)	168	37.5 (31.4, 43.6)	171	42.8 (36.4, 49.2)
50,000 or more	Yes	484	53.5 (49.5, 57.6)	465	55.5 (51.5, 59.6)	440	56.8 (52.3, 61.2)	412	57.3 (52.6, 62.1)	361	55.7 (50.7, 60.7)
	No	433	46.5 (42.4, 50.5)	409	44.5 (40.4, 48.5)	390	43.2 (38.8, 47.7)	366	42.7 (37.9, 47.4)	330	44.3 (39.3, 49.3)

Income	Quit attempts in the last 12 months	2016-18		2017-19		2018-20		2019-21		2020-22	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than 15,000	Yes	284	62.1 (56.5, 67.6)	274	60.6 (55.0, 66.1)	221	55.9 (49.8, 62.0)	185	53.8 (47.3, 60.4)	143	50.9 (43.7, 58.2)
	No	193	37.9 (32.4, 43.5)	186	39.4 (33.9, 45.0)	178	44.1 (38.0, 50.2)	166	46.2 (39.6, 52.7)	135	49.1 (41.8, 56.3)
15,000 to Less than 25,000	Yes	331	60.6 (55.5, 65.6)	330	58.9 (53.9, 63.9)	288	55.2 (50.0, 60.4)	251	54.6 (49.2, 59.9)	193	53.2 (47.2, 59.3)
	No	266	39.4 (34.4, 44.5)	273	41.1 (36.1, 46.1)	279	44.8 (39.6, 50.0)	249	45.4 (40.1, 50.8)	201	46.8 (40.7, 52.8)
25,000 to Less than 35,000	Yes	188	55.8 (49.2, 62.4)	172	54.9 (48.1, 61.7)	161	54.4 (47.6, 61.2)	147	52.4 (45.4, 59.4)	151	50.7 (43.9, 57.4)
	No	161	44.2 (37.6, 50.8)	152	45.1 (38.3, 51.9)	151	45.6 (38.8, 52.4)	147	47.6 (40.6, 54.6)	155	49.3 (42.6, 56.1)
35,000 to Less than 50,000	Yes	191	56.9 (50.4, 63.3)	171	55.8 (49.4, 62.2)	160	60.7 (53.9, 67.5)	170	63.3 (56.6, 70.1)	146	59.7 (52.3, 67.2)
	No	170	43.1 (36.7, 49.6)	170	44.2 (37.8, 50.6)	143	39.3 (32.5, 46.1)	127	36.7 (29.9, 43.4)	121	40.3 (32.8, 47.7)
50,000 or more	Yes	335	54.8 (49.8, 59.7)	362	56.1 (51.3, 60.8)	340	56.1 (51.4, 60.8)	303	54.2 (49.3, 59.1)	244	46.8 (41.7, 51.9)
	No	304	45.2 (40.3, 50.2)	301	43.9 (39.2, 48.7)	297	43.9 (39.2, 48.6)	301	45.8 (40.9, 50.7)	306	53.2 (48.1, 58.3)

Income	Quit attempts in the last 12 months	2021-23	
		N	Weighted% (95%CI)
Less than 15,000	Yes	130	54.0 (46.1, 61.9)
	No	107	46.0 (38.1, 53.9)
15,000 to Less than 25,000	Yes	146	51.7 (44.8, 58.7)
	No	149	48.3 (41.3, 55.2)
25,000 to Less than 35,000	Yes	144	51.8 (45.0, 58.6)
	No	145	48.2 (41.4, 55.0)
35,000 to Less than 50,000	Yes	161	57.2 (50.4, 63.9)
	No	139	42.8 (36.1, 49.6)
50,000 or more	Yes	236	43.3 (38.3, 48.3)
	No	328	56.7 (51.7, 61.7)

Table 7. Past quit attempts among Oklahoma adult current smokers by area of residence, Oklahoma BRFSS 2011 – 2023

Area of Residence	Quit attempts in the last 12 months	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Urban	Yes	493	59.2 (54.4, 64.1)	303	63.3 (58.3, 68.3)	349	60.4 (55.3, 65.5)	283	62.7 (57.2, 68.2)	222	61.6 (54.6, 68.5)
	No	313	40.8 (35.9, 45.6)	202	36.7 (31.7, 41.7)	229	39.6 (34.5, 44.7)	182	37.3 (31.8, 42.8)	155	38.4 (31.5, 45.4)
Rural	Yes	582	58.1 (53.9, 62.2)	647	60.5 (56.6, 64.5)	543	56.9 (52.8, 61.0)	550	60.5 (56.4, 64.6)	400	55.9 (51.0, 60.9)
	No	466	41.9 (37.8, 46.1)	478	39.5 (35.5, 43.4)	440	43.1 (39.0, 47.2)	403	39.5 (35.4, 43.6)	368	44.1 (39.1, 49.0)

Area of Residence	Quit attempts in the last 12 months	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Urban	Yes	169	57.5 (50.3, 64.7)	186	61.0 (54.5, 67.6)	129	54.6 (74.2, 61.9)	163	61.3 (54.5, 68.0)	138	60.5 (53.2, 67.8)
	No	138	42.5 (35.3, 49.7)	136	39.0 (32.4, 45.5)	119	45.4 (38.1, 52.8)	108	38.7 (32.0, 45.5)	96	39.5 (32.2, 46.8)
Rural	Yes	361	57.4 (52.6, 62.3)	379	58.3 (53.7, 63.0)	301	54.2 (49.0, 59.3)	363	55.9 (51.3, 60.5)	259	51.7 (46.5, 56.9)
	No	336	42.6 (37.7, 47.4)	305	41.7 (37.0, 46.3)	284	45.8 (40.7, 51.0)	326	44.1 (39.5, 48.7)	298	48.3 (43.1, 53.5)

Area of Residence	Quit attempts in the last 12 months	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Urban	Yes	79	58.7 (49.5, 67.8)	117	55.2 (47.6, 62.8)	96	52.9 (45.0, 60.8)
	No	75	41.3 (32.2, 50.5)	95	44.8 (37.2, 52.4)	102	47.1 (39.2, 55.0)
Rural	Yes	232	48.6 (43.2, 53.9)	207	41.9 (36.6, 47.2)	214	50.6 (44.9, 56.4)
	No	268	51.4 (46.1, 56.8)	258	58.1 (52.8, 63.4)	233	49.4 (43.6, 55.1)

Table 8. Past quit attempts among Oklahoma adult current smokers by poor physical health, Oklahoma BRFSS 2011 – 2023

Poor Physical Health	Quit attempts in the last 12 months	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Yes	294	62.6 (56.6, 68.5)	255	68.4 (62.6, 74.3)	244	65.2 (59.1, 71.3)	218	63.2 (56.6, 69.7)	172	62.0 (53.4, 70.6)
	No	172	37.4 (31.5, 43.4)	146	31.6 (25.7, 37.4)	144	34.8 (28.7, 40.9)	131	36.8 (30.3, 43.4)	108	38.0 (29.4, 46.6)
No	Yes	781	57.6 (53.9, 61.2)	695	59.7 (56.2, 63.3)	648	56.5 (52.8, 60.1)	615	60.9 (57.1, 64.7)	450	57.1 (52.5, 61.7)
	No	607	42.4 (38.8, 46.1)	534	40.3 (36.7, 43.8)	525	43.5 (39.9, 47.2)	454	39.1 (35.3, 42.9)	415	42.9 (38.3, 47.5)

Poor Physical Health	Quit attempts in the last 12 months	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Yes	132	59.2 (51.4, 67.1)	153	65.0 (57.6, 72.3)	134	71.0 (63.5, 78.4)	138	62.0 (54.5, 69.5)	94	57.2 (48.6, 65.7)
	No	126	40.8 (32.9, 48.6)	97	35.0 (27.7, 42.4)	71	29.0 (21.6, 36.5)	104	38.0 (30.5, 45.5)	92	42.8 (34.3, 51.4)
No	Yes	378	57.8 (53.0, 62.6)	395	58.7 (54.2, 63.2)	284	49.2 (44.2, 54.2)	363	56.9 (52.3, 61.5)	283	54.2 (49.2, 59.3)
	No	325	42.2 (37.4, 47.0)	324	41.3 (36.8, 45.8)	316	50.8 (45.8, 55.8)	311	43.1 (38.5, 47.7)	290	45.8 (40.7, 50.8)

Poor Physical Health	Quit attempts in the last 12 months	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Yes	75	52.5 (43.0, 62.0)	94	53.4 (44.8, 61.9)	84	65.0 (55.8, 74.1)
	No	79	47.5 (38.0, 57.0)	80	46.6 (38.1, 55.2)	59	35.0 (25.9, 44.2)
No	Yes	218	52.1 (46.4, 57.8)	216	43.9 (38.6, 49.1)	213	48.4 (42.9, 53.8)
	No	243	47.9 (42.2, 53.6)	262	56.1 (50.9, 61.4)	256	51.6 (46.2, 57.1)

Table 9. Past quit attempts among Oklahoma adult current smokers by Frequent Mental Distress (FMD), Oklahoma BRFSS 2011 – 2023

Frequent Mental Distress	Quit attempts in the last 12 months	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Yes	290	58.8 (52.6, 65.0)	260	65.0 (58.7, 71.3)	255	64.5 (58.1, 71.0)	224	66.3 (60.1, 72.4)	163	63.4 (55.3, 71.6)
	No	203	41.2 (35.0, 47.4)	153	35.0 (28.7, 41.3)	148	35.5 (29.0, 41.9)	122	33.7 (27.6, 39.9)	102	36.6 (28.4, 44.7)
No	Yes	785	58.5 (54.8, 62.2)	690	60.4 (56.8, 64.0)	637	55.9 (52.2, 59.5)	609	59.8 (55.9, 63.7)	459	56.5 (51.8, 61.2)
	No	576	41.5 (37.8, 45.2)	527	39.6 (36.0, 43.2)	521	44.1 (40.5, 47.8)	463	40.2 (36.3, 44.1)	421	43.5 (38.8, 48.2)

Frequent Mental Distress	Quit attempts in the last 12 months	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Yes	127	57.7 (49.4, 65.9)	143	62.7 (54.9, 70.4)	127	63.1 (55.2, 71.0)	144	63.9 (56.7, 71.2)	116	58.2 (50.6, 65.8)
	No	111	42.3 (34.1, 50.6)	90	37.3 (29.6, 45.1)	86	36.9 (29.0, 44.8)	99	36.1 (28.8, 43.3)	102	41.8 (34.2, 49.4)
No	Yes	390	58.3 (53.6, 63.0)	404	58.8 (54.4, 63.3)	283	50.2 (45.1, 55.4)	361	56.7 (52.1, 61.3)	259	52.7 (47.4, 58.1)
	No	350	41.7 (37.0, 46.4)	339	41.2 (36.7, 45.6)	305	49.8 (44.6, 54.9)	310	43.3 (38.7, 47.9)	279	47.3 (41.9, 52.6)

Frequent Mental Distress	Quit attempts in the last 12 months	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Yes	88	55.6 (46.5, 64.6)	85	49.5 (40.2, 58.8)	101	61.8 (52.8, 70.9)
	No	73	44.4 (35.4, 53.5)	77	50.5 (41.2, 59.8)	74	38.2 (29.1, 47.2)
No	Yes	210	51.3 (45.5, 57.0)	229	47.2 (42.0, 52.5)	196	47.2 (41.7, 52.8)
	No	250	48.7 (43.0, 54.5)	256	52.8 (47.5, 58.0)	248	52.8 (47.2, 58.3)

Frequent Mental Distress = 14 or more days of poor mental health

Table 10. Past quit attempts among Oklahoma adult current smokers by health insurance coverage, Oklahoma BRFSS 2011 – 2023

Health Insurance Status	Quit attempts in the last 12 months	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Insured	Yes	800	58.1 (54.4, 61.7)	716	61.3 (57.7, 64.8)	658	58.5 (54.8, 62.2)	671	62.5 (58.9, 66.1)	508	58.9 (54.3, 63.4)
	No	589	41.9 (38.3, 45.6)	526	38.7 (35.2, 42.3)	510	41.5 (37.8, 45.2)	475	37.5 (33.9, 41.1)	445	41.1 (36.6, 45.7)
Uninsured	Yes	271	59.3 (53.2, 65.5)	232	62.1 (55.8, 68.4)	229	56.6 (50.5, 62.6)	160	57.5 (49.8, 65.1)	110	54.6 (45.6, 63.6)
	No	190	40.7 (34.5, 46.8)	153	37.9 (31.6, 44.2)	159	43.4 (37.4, 49.5)	108	42.5 (34.9, 50.2)	78	45.4 (36.4, 54.4)

Health Insurance Status	Quit attempts in the last 12 months	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Insured	Yes	434	57.1 (52.7, 61.6)	465	59.6 (55.3, 63.8)	348	54.3 (49.6, 59.1)	419	58.2 (53.9, 62.5)	327	55.1 (50.3, 59.8)
	No	393	42.9 (38.4, 47.3)	362	40.4 (36.2, 44.7)	324	45.7 (40.9, 50.4)	347	41.8 (37.5, 46.1)	327	44.9 (40.2, 49.7)
Uninsured	Yes	94	58.0 (48.8, 67.1)	100	59.1 (50.4, 67.8)	81	54.5 (45.3, 63.7)	107	57.4 (49.0, 65.7)	68	55.4 (45.5, 65.3)
	No	80	42.0 (32.9, 51.2)	78	40.9 (32.2, 49.6)	78	45.5 (36.3, 54.7)	86	42.6 (34.3, 51.0)	66	44.6 (34.7, 54.5)

Health Insurance Status	Quit attempts in the last 12 months	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Insured	Yes	262	52.0 (46.8, 57.2)	290	48.2 (43.4, 52.9)	266	49.3 (44.3, 54.4)
	No	285	48.0 (42.8, 53.2)	310	51.8 (47.1, 56.6)	296	50.7 (45.6, 55.7)
Uninsured	Yes	46	56.1 (44.6, 67.7)	29	40.3 (26.9, 53.6)	37	66.5 (54.8, 78.1)
	No	46	43.9 (32.3, 55.4)	37	59.7 (46.4, 73.1)	31	33.5 (21.9, 45.2)

Table 11. Past quit attempts among Oklahoma adult current smokers by disability status, Oklahoma BRFSS 2016 – 2023

Disability Status	Quit attempts in the last 12 months	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
One or more disability	Yes	277	60.7 (55.2, 66.1)	297	60.2 (54.7, 65.6)	218	53.9 (47.8, 60.0)	270	57.9 (52.5, 63.2)	207	52.1 (46.3, 57.9)
	No	224	39.3 (33.9, 44.8)	212	39.8 (34.4, 45.3)	192	46.1 (40.0, 52.2)	218	42.1 (36.8, 47.5)	220	47.9 (42.1, 53.7)
No disability	Yes	245	55.4 (49.5, 61.4)	262	58.7 (53.2, 64.1)	210	54.5 (48.6, 60.4)	248	58.3 (52.7, 63.9)	182	58.7 (52.4, 64.9)
	No	238	44.6 (38.6, 50.5)	223	41.3 (35.9, 46.8)	209	45.5 (39.6, 51.4)	211	41.7 (36.1, 47.3)	169	41.3 (35.1, 47.6)

Disability Status	Quit attempts in the last 12 months	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
One or more disability	Yes	277	60.7 (55.2, 66.1)	297	60.2 (54.7, 65.6)	218	53.9 (47.8, 60.0)	270	57.9 (52.5, 63.2)	207	52.1 (46.3, 57.9)
	No	224	39.3 (33.9, 44.8)	212	39.8 (34.4, 45.3)	192	46.1 (40.0, 52.2)	218	42.1 (36.8, 47.5)	220	47.9 (42.1, 53.7)
No disability	Yes	245	55.4 (49.5, 61.4)	262	58.7 (53.2, 64.1)	210	54.5 (48.6, 60.4)	248	58.3 (52.7, 63.9)	182	58.7 (52.4, 64.9)
	No	238	44.6 (38.6, 50.5)	223	41.3 (35.9, 46.8)	209	45.5 (39.6, 51.4)	211	41.7 (36.1, 47.3)	169	41.3 (35.1, 47.6)

Disability Status	Quit attempts in the last 12 months	2021		2022		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
One or more disability	Yes	175	53.7 (47.3, 60.1)	188	50.8 (44.7, 57.0)	183	55.6 (49.2, 62.0)
	No	168	46.3 (39.9, 52.7)	182	49.2 (43.0, 55.3)	165	44.4 (38.0, 50.8)
No disability	Yes	125	50.4 (43.3, 57.6)	133	43.0 (36.5, 49.4)	119	45.8 (38.8, 52.8)
	No	165	49.6 (42.4, 56.7)	166	57.0 (50.6, 63.5)	160	54.2 (47.2, 61.2)

Section VI. Smoking policies at home

Table 1. Proportion of Oklahoma adults reporting smoking ban at home, Oklahoma BRFSS 2011 - 2021, 2023

Home Smoking Policy	2011		2012		2013		2014		2015	
	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Smoking Ban	2712	75.3 (72.9, 77.6)	2832	77.6 (75.7, 79.5)	2963	80.4 (78.5, 82.2)	3033	83.7 (81.7, 85.5)	2581	82.7 (80.5, 84.9)
No Ban	831	24.7 (22.4, 27.1)	806	22.4 (20.5, 24.3)	679	19.6 (17.8, 21.5)	586	16.3 (14.5, 18.1)	453	17.3 (15.1, 19.5)

Home Smoking Policy	2016		2017		2018		2019		2020	
	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Smoking Ban	2777	84.0 (82.1, 85.9)	2341	83.3 (81.4, 85.3)	1938	82.8 (80.7, 85.0)	2399	86.6 (84.9, 88.2)	1709	87.6 (85.7, 89.4)
No Ban	452	16.0 (14.1, 17.9)	421	16.7 (14.7, 18.6)	329	17.2 (15.0, 19.3)	360	13.4 (11.8, 15.1)	257	12.4 (10.6, 14.3)

Home Smoking Policy	2021		2023	
	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Smoking Ban	1914	87.0 (85.0, 88.9)	2073	87.2 (85.5, 88.9)
No Ban	247	13.0 (11.1, 15.0)	267	12.8 (11.1, 14.5)

Table 2. Proportion of Oklahoma adults reporting smoking ban at home by sex, Oklahoma BRFSS 2011 – 2021, 2023

Sex	Home Smoking Policy	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Male	Smoking Ban	926	73.2 (69.1, 77.3)	1105	75.2 (72.2, 78.2)	1142	79.7 (77.0, 82.3)	1208	82.2 (79.3, 85.1)	970	78.3 (74.4, 82.1)
	No Ban	328	26.8 (22.7, 30.9)	368	24.8 (21.8, 27.8)	294	20.3 (17.7, 23.0)	259	17.8 (14.9, 20.7)	212	21.7 (17.9, 25.6)
Female	Smoking Ban	1786	77.3 (74.8, 79.7)	1727	79.9 (77.5, 82.2)	1821	81.0 (78.6, 83.5)	1825	85.1 (83.0, 87.2)	1611	86.9 (84.7, 89.2)
	No Ban	503	22.7 (20.3, 25.2)	438	20.1 (17.8, 22.5)	385	19.0 (16.5, 21.4)	327	14.9 (12.8, 17.0)	241	13.1 (10.8, 15.3)

Sex	Home Smoking Policy	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Male	Smoking Ban	1059	82.4 (79.4, 85.4)	939	82.0 (78.8, 85.2)	789	81.0 (77.5, 84.5)	1044	86.5 (84.0, 89.0)	743	87.8 (85.1, 90.5)
	No Ban	211	17.6 (14.6, 20.6)	181	18.0 (14.8, 21.2)	146	19.0 (15.5, 22.5)	156	13.5 (11.0, 16.0)	113	12.2 (9.5, 14.9)
Female	Smoking Ban	1718	85.5 (83.2, 87.9)	1402	84.6 (82.2, 87.0)	1149	84.7 (82.1, 87.3)	1355	86.7 (84.4, 88.9)	966	87.3 (84.9, 89.7)
	No Ban	241	14.5 (12.1, 16.8)	240	15.4 (13.0, 17.8)	183	15.3 (12.7, 17.9)	204	13.3 (11.1, 15.6)	144	12.7 (10.3, 15.1)

Sex	Home Smoking Policy	2021		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)
Male	Smoking Ban	813	86.5 (83.8, 89.3)	896	86.2 (83.6, 88.7)
	No Ban	127	13.5 (10.7, 16.2)	127	13.8 (11.3, 16.4)
Female	Smoking Ban	1101	87.4 (84.6, 90.2)	1177	88.3 (86.1, 90.5)
	No Ban	120	12.6 (9.8, 15.4)	140	11.7 (9.5, 13.9)

Table 3. Three-year rolling weighted percentages of Oklahoma adults reporting smoking ban at home by age, Oklahoma BRFSS 2011 – 2021, 2023

Age Group	Home Smoking Policy	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
18-24	Smoking Ban	329	74.8 (69.5, 80.2)	414	76.4 (71.8, 81.0)	379	78.3 (73.5, 83.2)	370	81.0 (76.1, 85.9)	290	77.7 (72.4, 83.1)
	No Ban	104	25.2 (19.8, 30.5)	113	23.6 (19.0, 28.2)	86	21.7 (16.8, 26.5)	66	19.0 (14.1, 23.9)	64	22.3 (16.9, 27.6)
25-34	Smoking Ban	871	80.7 (77.6, 83.8)	992	83.6 (81.0, 86.3)	844	85.0 (82.1, 87.8)	804	84.7 (81.7, 87.7)	714	84.7 (81.7, 87.6)
	No Ban	182	19.3 (16.2, 22.4)	172	16.4 (13.7, 19.0)	134	15.0 (12.2, 17.9)	125	15.3 (12.3, 18.3)	120	15.3 (12.4, 18.3)
35-44	Smoking Ban	995	80.3 (77.5, 83.0)	1065	82.6 (80.0, 85.2)	954	82.8 (79.8, 85.7)	913	85.1 (82.1, 88.1)	812	86.4 (83.5, 89.3)
	No Ban	226	19.7 (17.0, 22.5)	211	17.4 (14.8, 20.0)	169	17.2 (14.3, 20.2)	128	14.9 (11.9, 17.9)	99	13.6 (10.7, 16.5)
45-54	Smoking Ban	1347	74.4 (72.0, 76.8)	1379	79.1 (76.8, 81.3)	1281	80.9 (78.4, 83.4)	1164	81.2 (78.6, 83.8)	1023	81.8 (79.0, 84.5)
	No Ban	459	25.6 (23.2, 28.0)	374	20.9 (18.7, 23.2)	298	19.1 (16.6, 21.6)	255	18.8 (16.2, 21.4)	211	18.2 (15.5, 21.0)
55-64	Smoking Ban	1799	74.2 (72.1, 76.3)	1854	77.3 (75.2, 79.4)	1826	79.9 (77.9, 82.0)	1802	81.8 (79.7, 83.8)	1611	80.8 (78.5, 83.0)
	No Ban	605	25.8 (23.7, 27.9)	543	22.7 (20.6, 24.8)	445	20.1 (18.0, 22.1)	390	18.2 (16.2, 20.3)	347	19.2 (17.0, 21.5)
65 or more	Smoking Ban	3166	81.0 (79.5, 82.5)	3124	82.6 (81.1, 84.1)	3293	84.9 (83.5, 86.4)	3338	85.9 (84.5, 87.3)	3249	86.7 (85.3, 88.0)
	No Ban	740	19.0 (17.5, 20.5)	658	17.4 (15.9, 18.9)	586	15.1 (13.6, 16.5)	527	14.1 (12.7, 15.5)	485	13.3 (12.0, 14.7)

Age Group	Home Smoking Policy	2016-18		2017-19		2018-20		2019-21		2020-2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
18-24	Smoking Ban	285	79.4 (74.2, 84.6)	275	80.1 (74.9, 85.2)	288	84.4 (79.9, 89.0)	303	85.1 (80.8, 89.5)	291	85.1 (80.9, 89.3)
	No Ban	59	20.6 (15.4, 25.8)	58	19.9 (14.8, 25.1)	48	15.6 (11.0, 20.1)	47	14.9 (10.5, 19.2)	52	14.9 (10.7, 19.1)
25-34	Smoking Ban	703	84.8 (81.9, 87.7)	682	85.5 (82.7, 88.4)	643	88.2 (85.4, 91.0)	654	89.6 (86.9, 92.2)	648	90.2 (87.5, 92.9)
	No Ban	115	15.2 (12.3, 18.1)	106	14.5 (11.6, 17.3)	80	11.8 (9.0, 14.6)	72	10.4 (7.8, 13.1)	67	9.8 (7.1, 12.5)
35-44	Smoking Ban	790	86.6 (83.9, 89.3)	848	87.3 (84.8, 89.8)	796	87.2 (84.6, 89.7)	837	89.6 (87.3, 91.9)	810	89.2 (86.7, 91.6)
	No Ban	106	13.4 (10.7, 16.1)	113	12.7 (10.2, 15.2)	109	12.8 (10.3, 15.4)	95	10.4 (8.1, 12.7)	89	10.8 (8.4, 13.3)
45-54	Smoking Ban	941	80.5 (77.7, 83.4)	893	82.5 (79.7, 85.3)	843	83.1 (80.3, 85.9)	881	85.7 (83.1, 88.4)	847	86.4 (83.7, 89.1)
	No Ban	198	19.5 (16.6, 22.3)	166	17.5 (14.7, 20.3)	150	16.9 (14.1, 19.7)	131	14.3 (11.6, 16.9)	112	13.6 (10.9, 16.3)
55-64	Smoking Ban	1424	80.5 (78.1, 82.8)	1280	81.5 (79.2, 83.9)	1118	83.3 (80.9, 85.6)	1064	84.9 (82.6, 87.2)	966	84.3 (81.8, 86.8)
	No Ban	305	19.5 (17.2, 21.9)	267	18.5 (16.1, 20.8)	216	16.7 (14.4, 19.1)	194	15.1 (12.8, 17.4)	169	15.7 (13.2, 18.2)
65 or more	Smoking Ban	2913	86.6 (85.2, 88.1)	2700	86.6 (85.1, 88.1)	2358	86.6 (85.0, 88.2)	2283	86.6 (85.0, 88.3)	2134	87.5 (85.8, 89.2)
	No Ban	419	13.4 (11.9, 14.8)	400	13.4 (11.9, 14.9)	343	13.4 (11.8, 15.0)	325	13.4 (11.7, 15.0)	282	12.5 (10.8, 14.2)

Home smoking ban data were not available for 2022, therefore, three-year rolling percentage for 2020-2023 was based on data from 2020, 2021, and 2023.

Table 4. Three-year rolling weighted percentages of Oklahoma adults reporting smoking ban at home by race/ethnicity, Oklahoma BRFSS 2011 – 2021, 2023

Race/Ethnicity	Home Smoking Policy	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
NH-White	Smoking Ban	6485	78.6 (77.2, 79.9)	6698	80.9 (79.7, 82.2)	6559	82.8 (81.5, 84.1)	6525	83.5 (82.2, 84.9)	6042	84.2 (82.9, 85.5)
	No Ban	1622	21.4 (20.1, 22.8)	1469	19.1 (17.8, 20.3)	1214	17.2 (15.9, 18.5)	1098	16.5 (15.1, 17.8)	968	15.8 (14.5, 17.1)
NH-Black	Smoking Ban	454	70.4 (65.1, 75.7)	456	76.1 (71.6, 80.6)	454	74.8 (69.6, 80.0)	389	75.8 (70.3, 81.3)	356	74.4 (68.5, 80.4)
	No Ban	208	29.6 (24.3, 34.9)	158	23.9 (19.4, 28.4)	149	25.2 (20.0, 30.4)	113	24.2 (18.7, 29.7)	101	25.6 (19.6, 31.5)
NH-AI/AN	Smoking Ban	445	67.5 (62.5, 72.5)	496	71.2 (66.4, 75.9)	505	75.3 (70.3, 80.2)	531	79.7 (75.3, 84.1)	469	77.0 (72.0, 82.0)
	No Ban	195	32.5 (27.5, 37.5)	189	28.8 (24.1, 33.6)	151	24.7 (19.8, 29.7)	137	20.3 (15.9, 24.7)	116	23.0 (18.0, 28.0)
NH-Other	Smoking Ban	626	74.4 (69.2, 79.5)	640	78.7 (73.9, 83.4)	564	77.5 (71.9, 83.2)	495	81.8 (76.3, 87.4)	441	82.8 (77.8, 87.8)
	No Ban	222	25.6 (20.5, 30.8)	191	21.3 (16.6, 26.1)	150	22.5 (16.8, 28.1)	101	18.2 (12.6, 23.7)	92	17.2 (12.2, 22.2)
Hispanic	Smoking Ban	462	88.8 (85.2, 92.4)	498	89.7 (86.3, 93.2)	446	92.6 (90.0, 95.3)	396	92.6 (89.8, 95.5)	324	88.3 (84.0, 92.6)
	No Ban	62	11.2 (7.6, 14.8)	57	10.3 (6.8, 13.7)	46	7.4 (4.7, 10.0)	37	7.4 (4.5, 10.2)	40	11.7 (7.4, 16.0)

Race/Ethnicity	Home Smoking Policy	2016-18		2017-19		2018-20		2019-21		2020-23	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
NH-White	Smoking Ban	5486	83.8 (82.5, 85.1)	5071	85.1 (83.8, 86.3)	4480	85.9 (84.6, 87.1)	4423	88.2 (87.0, 89.3)	4191	88.5 (87.3, 89.7)
	No Ban	888	16.2 (14.9, 17.5)	791	14.9 (13.7, 16.2)	672	14.1 (12.9, 15.4)	576	11.8 (10.7, 13.0)	510	11.5 (10.3, 12.7)
NH-Black	Smoking Ban	334	77.4 (71.8, 83.0)	338	80.0 (74.7, 85.3)	319	82.2 (77.0, 87.3)	299	82.8 (77.6, 88.0)	265	83.3 (78.1, 88.6)
	No Ban	83	22.6 (17.0, 28.2)	75	20.0 (14.7, 25.3)	67	17.8 (12.7, 23.0)	65	17.2 (12.0, 22.4)	54	16.7 (11.4, 21.9)
NH-AI/AN	Smoking Ban	396	77.7 (72.7, 82.8)	392	76.9 (71.7, 82.1)	399	81.8 (77.2, 86.4)	421	81.1 (76.8, 85.4)	404	84.0 (79.9, 88.1)
	No Ban	101	22.3 (17.2, 27.3)	99	23.1 (17.9, 28.3)	91	18.2 (13.6, 22.8)	97	18.9 (14.6, 23.2)	77	16.0 (11.9, 20.1)
NH-Other	Smoking Ban	453	85.5 (81.2, 89.8)	448	84.9 (80.6, 89.2)	398	85.5 (81.2, 89.8)	383	85.0 (80.8, 89.2)	359	82.7 (78.5, 86.9)
	No Ban	79	14.5 (10.2, 18.8)	82	15.1 (10.8, 19.4)	64	14.5 (10.2, 18.8)	68	15.0 (10.8, 19.2)	74	17.3 (13.1, 21.5)
Hispanic	Smoking Ban	324	87.0 (82.5, 91.5)	354	85.7 (81.2, 90.2)	375	88.8 (84.8, 92.8)	397	88.0 (84.1, 92.0)	390	89.1 (85.2, 93.0)
	No Ban	43	13.0 (8.5, 17.5)	51	14.3 (9.8, 18.8)	43	11.2 (7.2, 15.2)	46	12.0 (8.0, 15.9)	38	10.9 (7.0, 14.8)

Home smoking ban data were not available for 2022, therefore, three-year rolling percentage for 2020-2023 was based on data from 2020, 2021, and 2023.

Table 5. Three-year rolling weighted percentages of Oklahoma adults reporting smoking ban at home by level of education, Oklahoma BRFSS 2011 – 2021, 2023

Education	Home Smoking Policy	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than high school	Smoking Ban	819	65.9 (61.7, 70.1)	812	69.2 (65.3, 73.1)	704	70.0 (65.6, 74.5)	606	70.4 (65.8, 75.0)	490	69.9 (65.0, 74.7)
	No Ban	407	34.1 (29.9, 38.3)	338	30.8 (26.9, 34.7)	252	30.0 (25.5, 34.4)	202	29.6 (25.0, 34.2)	170	30.1 (25.3, 35.0)
High school	Smoking Ban	2526	73.6 (71.4, 75.8)	2641	76.5 (74.4, 78.5)	2540	79.1 (77.1, 81.1)	2479	80.3 (78.2, 82.3)	2227	79.9 (77.7, 82.1)
	No Ban	864	26.4 (24.2, 28.6)	792	23.5 (21.5, 25.6)	657	20.9 (18.9, 22.9)	596	19.7 (17.7, 21.8)	508	20.1 (17.9, 22.3)
Some college	Smoking Ban	2396	79.0 (77.1, 81.0)	2527	82.1 (80.4, 83.8)	2491	83.8 (82.0, 85.6)	2498	85.7 (83.9, 87.4)	2405	85.7 (83.9, 87.4)
	No Ban	681	21.0 (19.0, 22.9)	617	17.9 (16.2, 19.6)	528	16.2 (14.4, 18.0)	453	14.3 (12.6, 16.1)	418	14.3 (12.6, 16.1)
College graduate	Smoking Ban	2756	90.4 (89.2, 91.5)	2841	91.8 (90.7, 92.9)	2835	92.5 (91.3, 93.7)	2797	93.1 (91.8, 94.3)	2564	92.7 (91.4, 93.9)
	No Ban	363	9.6 (8.5, 10.8)	323	8.2 (7.1, 9.3)	280	7.5 (6.3, 8.7)	239	6.9 (5.7, 8.2)	230	7.3 (6.1, 8.6)

Education	Home Smoking Policy	2016-18		2017-19		2018-20		2019-21		2020-23	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than high school	Smoking Ban	434	70.7 (66.0, 75.5)	419	72.1 (67.4, 76.9)	410	77.0 (72.6, 81.4)	393	80.1 (76.0, 84.1)	337	82.2 (78.1, 86.2)
	No Ban	159	29.3 (24.5, 34.0)	156	27.9 (23.1, 32.6)	134	23.0 (18.6, 27.4)	118	19.9 (15.9, 24.0)	86	17.8 (13.8, 21.9)
High school	Smoking Ban	1969	78.5 (76.2, 80.9)	1835	79.9 (77.6, 82.2)	1613	81.9 (79.7, 84.1)	1573	84.6 (82.5, 86.7)	1438	85.3 (83.3, 87.4)
	No Ban	468	21.5 (19.1, 23.8)	403	20.1 (17.8, 22.4)	343	18.1 (15.9, 20.3)	291	15.4 (13.3, 17.5)	261	14.7 (12.6, 16.7)
Some college	Smoking Ban	2258	86.4 (84.8, 88.1)	2126	86.6 (85.0, 88.3)	1912	87.5 (85.8, 89.1)	1871	87.6 (85.9, 89.3)	1746	86.7 (84.9, 88.5)
	No Ban	368	13.6 (11.9, 15.2)	341	13.4 (11.7, 15.0)	294	12.5 (10.9, 14.2)	284	12.4 (10.7, 14.1)	272	13.3 (11.5, 15.1)
College graduate	Smoking Ban	2382	92.8 (91.4, 94.1)	2288	93.0 (91.7, 94.2)	2100	92.5 (90.9, 94.0)	2168	93.0 (91.5, 94.5)	2157	93.0 (91.4, 94.5)
	No Ban	206	7.2 (5.9, 8.6)	209	7.0 (5.8, 8.3)	174	7.5 (6.0, 9.1)	169	7.0 (5.5, 8.5)	150	7.0 (5.5, 8.6)

Home smoking ban data were not available for 2022, therefore, three-year rolling percentage for 2020-2023 was based on data from 2020, 2021, and 2023.

Table 6. Three-year rolling weighted percentages of Oklahoma adults reporting smoking ban at home by income level, Oklahoma BRFSS 2011 – 2021, 2023

Income	Home Smoking Policy	2011-13		2012-14		2013-15		2014-16		2015-17	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than 15,000	Smoking Ban	912	59.7 (56.1, 63.3)	892	63.5 (60.0, 67.0)	760	65.4 (61.6, 69.3)	661	65.2 (60.8, 69.6)	599	68.2 (63.7, 72.6)
	No Ban	576	40.3 (36.7, 43.9)	502	36.5 (33.0, 40.0)	389	34.6 (30.7, 38.4)	308	34.8 (30.4, 39.2)	243	31.8 (27.4, 36.3)
15,000 to Less than 25,000	Smoking Ban	1475	70.2 (66.9, 73.5)	1474	73.5 (70.5, 76.4)	1320	74.1 (70.8, 77.4)	1208	73.2 (69.7, 76.7)	1078	74.1 (70.5, 77.6)
	No Ban	557	29.8 (26.5, 33.1)	481	26.5 (23.6, 29.5)	398	25.9 (22.6, 29.2)	355	26.8 (23.3, 30.3)	312	25.9 (22.4, 29.5)
25,000 to Less than 35,000	Smoking Ban	938	76.9 (73.6, 80.2)	958	78.4 (75.2, 81.6)	901	80.3 (76.9, 83.8)	856	82.4 (78.7, 86.0)	737	80.6 (76.6, 84.6)
	No Ban	285	23.1 (19.8, 26.4)	263	21.6 (18.4, 24.8)	216	19.7 (16.2, 23.1)	168	17.6 (14.0, 21.3)	152	19.4 (15.4, 23.4)
35,000 to Less than 50,000	Smoking Ban	1168	79.3 (76.3, 82.4)	1200	82.3 (79.6, 85.0)	1193	85.9 (83.5, 88.3)	1155	88.5 (86.2, 90.9)	1023	88.0 (85.5, 90.5)
	No Ban	289	20.7 (17.6, 23.7)	248	17.7 (15.0, 20.4)	205	14.1 (11.7, 16.5)	160	11.5 (9.1, 13.8)	150	12.0 (9.5, 14.5)
50,000 or more	Smoking Ban	3037	89.4 (88.1, 90.7)	3270	90.9 (89.7, 92.1)	3216	91.3 (89.9, 92.7)	3159	92.1 (90.8, 93.4)	2819	91.3 (89.8, 92.8)
	No Ban	356	10.6 (9.3, 11.9)	337	9.1 (7.9, 10.3)	292	8.7 (7.3, 10.1)	270	7.9 (6.6, 9.2)	237	8.7 (7.2, 10.2)

Income	Home Smoking Policy	2016-18		2017-19		2018-20		2019-21		2020-23	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Less than 15,000	Smoking Ban	517	66.8 (62.0, 71.6)	436	67.7 (62.9, 72.5)	361	70.0 (65.1, 74.9)	347	73.5 (68.5, 78.4)	285	76.0 (70.8, 81.2)
	No Ban	209	33.2 (28.4, 38.0)	185	32.3 (27.5, 37.1)	160	30.0 (25.1, 34.9)	135	26.5 (21.6, 31.5)	98	24.0 (18.8, 29.2)
15,000 to Less than 25,000	Smoking Ban	982	76.5 (73.1, 80.0)	931	80.1 (76.9, 83.3)	850	81.8 (78.6, 85.0)	764	83.2 (80.1, 86.3)	531	82.6 (78.8, 86.3)
	No Ban	263	23.5 (20.0, 26.9)	227	19.9 (16.7, 23.1)	181	18.2 (15.0, 21.4)	157	16.8 (13.7, 19.9)	114	17.4 (13.7, 21.2)
25,000 to Less than 35,000	Smoking Ban	672	80.0 (75.9, 84.0)	630	78.9 (74.7, 83.0)	557	79.9 (75.7, 84.0)	552	80.5 (76.6, 84.5)	494	81.3 (77.3, 85.4)
	No Ban	142	20.0 (16.0, 24.1)	142	21.1 (17.0, 25.3)	126	20.1 (16.0, 24.3)	124	19.5 (15.5, 23.4)	109	18.7 (14.6, 22.7)
35,000 to Less than 50,000	Smoking Ban	907	85.1 (82.1, 88.1)	824	84.7 (81.7, 87.8)	777	86.3 (83.3, 89.4)	766	88.6 (85.8, 91.4)	792	87.5 (84.8, 90.3)
	No Ban	148	14.9 (11.9, 17.9)	144	15.3 (12.2, 18.3)	122	13.7 (10.6, 16.7)	104	11.4 (8.6, 14.2)	111	12.5 (9.7, 15.2)
50,000 or more	Smoking Ban	2688	91.1 (89.6, 92.6)	2713	91.3 (89.8, 92.8)	2542	92.4 (91.0, 93.7)	2560	93.1 (91.8, 94.4)	2601	92.7 (91.4, 93.9)
	No Ban	234	8.9 (7.4, 10.4)	233	8.7 (7.2, 10.2)	195	7.6 (6.3, 9.0)	181	6.9 (5.6, 8.2)	183	7.3 (6.1, 8.6)

Home smoking ban data were not available for 2022, therefore, three-year rolling percentage for 2020-2023 was based on data from 2020, 2021, and 2023.

Table 7. Proportion of Oklahoma adults reporting smoking ban at home by area of residence, Oklahoma BRFSS 2011 – 2021, 2023

Area of Residence	Home Smoking Policy	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Urban	Smoking Ban	1217	76.8 (73.4, 80.2)	1055	82.7 (80.1, 85.2)	1267	83.8 (81.3, 86.3)	1132	85.8 (83.2, 88.4)	951	84.9 (81.6, 88.1)
	No Ban	353	23.2 (19.8, 26.6)	242	17.3 (14.8, 19.9)	243	16.2 (13.7, 18.7)	181	14.2 (11.6, 16.8)	156	15.1 (11.9, 18.4)
Rural	Smoking Ban	1495	74.1 (70.9, 77.3)	1777	73.9 (71.2, 76.5)	1696	77.9 (75.3, 80.4)	1901	82.1 (79.7, 84.5)	1630	81.1 (78.1, 84.1)
	No Ban	478	25.9 (22.7, 29.1)	564	26.1 (23.5, 28.8)	436	22.1 (19.6, 24.7)	405	17.9 (15.5, 20.3)	297	18.9 (15.9, 21.9)

Area of Residence	Home Smoking Policy	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Urban	Smoking Ban	1032	87.3 (84.4, 90.2)	829	86.7 (83.6, 89.7)	726	84.7 (81.4, 88.0)	823	87.7 (84.9, 90.5)	645	89.8 (86.9, 92.7)
	No Ban	128	12.7 (9.8, 15.6)	118	13.3 (10.3, 16.4)	108	15.3 (12.0, 18.6)	109	12.3 (9.5, 15.1)	71	10.2 (7.3, 13.1)
Rural	Smoking Ban	1745	81.7 (79.2, 84.2)	1512	80.9 (78.3, 83.5)	1212	81.4 (78.6, 84.3)	1576	85.7 (83.7, 87.8)	1064	85.8 (83.5, 88.2)
	No Ban	324	18.3 (15.8, 20.8)	303	19.1 (16.5, 21.7)	221	18.6 (15.7, 21.4)	251	14.3 (12.2, 16.3)	186	14.2 (11.8, 16.5)

Area of Residence	Home Smoking Policy	2021		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)
Urban	Smoking Ban	589	89.2 (85.9, 92.4)	774	88.0 (85.4, 90.6)
	No Ban	61	10.8 (7.6, 14.1)	90	12.0 (9.4, 14.6)
Rural	Smoking Ban	1325	85.3 (82.9, 87.7)	1299	86.6 (84.4, 88.8)
	No Ban	186	14.7 (12.3, 17.1)	177	13.4 (11.2, 15.6)

Table 8. Proportion of Oklahoma adults reporting smoking ban at home by Physical health (14 or more physically unhealthy days), Oklahoma BRFSS 2011 – 2021, 2023

Poor Physical Health	Home Smoking Policy	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Smoking Ban	416	63.4 (58.4, 68.5)	435	70.6 (65.7, 75.4)	410	69.0 (63.8, 74.1)	452	74.4 (69.0, 79.7)	374	72.8 (67.1, 78.5)
	No Ban	228	36.6 (31.5, 41.6)	191	29.4 (24.6, 34.3)	175	31.0 (25.9, 36.2)	139	25.6 (20.3, 31.0)	126	27.2 (21.5, 32.9)
No	Smoking Ban	2238	77.3 (74.6, 79.9)	2354	79.2 (77.1, 81.3)	2490	82.4 (80.5, 84.3)	2526	85.4 (83.5, 87.3)	2167	84.9 (82.5, 87.3)
	No Ban	587	22.7 (20.1, 25.4)	595	20.8 (18.7, 22.9)	486	17.6 (15.7, 19.5)	431	14.6 (12.7, 16.5)	313	15.1 (12.7, 17.5)

Poor Physical Health	Home Smoking Policy	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Smoking Ban	406	73.6 (68.2, 79.1)	362	74.6 (69.3, 79.9)	280	71.9 (65.7, 78.0)	345	76.5 (70.9, 82.0)	217	81.4 (75.7, 87.1)
	No Ban	122	26.4 (20.9, 31.8)	106	25.4 (20.1, 30.7)	81	28.1 (22.0, 34.3)	90	23.5 (18.0, 29.1)	51	18.6 (12.9, 24.3)
No	Smoking Ban	2298	86.5 (84.5, 88.4)	1910	85.3 (83.1, 87.4)	1611	84.7 (82.3, 87.0)	1960	88.6 (86.9, 90.3)	1416	88.3 (86.3, 90.3)
	No Ban	312	13.5 (11.6, 15.5)	294	14.7 (12.6, 16.9)	241	15.3 (13.0, 17.7)	253	11.4 (9.7, 13.1)	198	11.7 (9.7, 13.7)

Poor Physical Health	Home Smoking Policy	2021		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Smoking Ban	246	79.5 (74.1, 84.8)	254	79.2 (73.8, 84.7)
	No Ban	66	20.5 (15.2, 25.9)	59	20.8 (15.3, 26.2)
No	Smoking Ban	1575	88.2 (86.0, 90.4)	1737	88.6 (86.8, 90.4)
	No Ban	166	11.8 (9.6, 14.0)	197	11.4 (9.6, 13.2)

Table 9. Proportion of Oklahoma adults reporting smoking ban at home by Frequent Mental Distress (FMD), Oklahoma BRFSS 2011 – 2021, 2023

Frequent Mental Distress	Home Smoking Policy	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Smoking Ban	262	50.7 (43.1, 58.3)	282	61.0 (55.1, 67.0)	309	65.5 (59.2, 71.8)	335	76.1 (71.0, 81.2)	241	71.1 (64.0, 78.2)
	No Ban	213	49.3 (41.7, 56.9)	181	39.0 (33.0, 44.9)	157	34.5 (28.2, 40.8)	118	23.9 (18.8, 29.0)	92	28.9 (21.8, 36.0)
No	Smoking Ban	2416	79.5 (77.3, 81.8)	2505	80.1 (78.1, 82.1)	2606	82.9 (81.1, 84.7)	2660	84.9 (83.0, 86.8)	2306	84.7 (82.4, 87.1)
	No Ban	603	20.5 (18.2, 22.7)	616	19.9 (17.9, 21.9)	510	17.1 (15.3, 18.9)	459	15.1 (13.2, 17.0)	348	15.3 (12.9, 17.6)

Frequent Mental Distress	Home Smoking Policy	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Smoking Ban	282	69.7 (63.1, 76.4)	271	70.4 (64.1, 76.8)	233	69.5 (62.6, 76.5)	295	79.6 (74.3, 84.8)	212	77.5 (71.4, 83.7)
	No Ban	95	30.3 (23.6, 36.9)	88	29.6 (23.2, 35.9)	79	30.5 (23.5, 37.4)	82	20.4 (15.2, 25.7)	60	22.5 (16.3, 28.6)
No	Smoking Ban	2453	86.3 (84.4, 88.2)	2022	86.2 (84.3, 88.2)	1660	85.6 (83.4, 87.8)	2026	88.8 (87.1, 90.4)	1432	89.7 (87.9, 91.5)
	No Ban	349	13.7 (11.8, 15.6)	321	13.8 (11.8, 15.7)	242	14.4 (12.2, 16.6)	256	11.2 (9.6, 12.9)	185	10.3 (8.5, 12.1)

Frequent Mental Distress	Home Smoking Policy	2021		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)
Yes	Smoking Ban	256	80.3 (74.4, 86.1)	268	77.1 (71.7, 82.5)
	No Ban	52	19.7 (13.9, 25.6)	71	22.9 (17.5, 28.3)
No	Smoking Ban	1600	88.5 (86.5, 90.6)	1743	89.7 (88.0, 91.4)
	No Ban	185	11.5 (9.4, 13.5)	181	10.3 (8.6, 12.0)

Frequent Mental Distress = 14 or more days of poor mental health

Table 10. Proportion of Oklahoma adults reporting smoking ban at home by health insurance coverage, Oklahoma BRFSS 2011 – 2021, 2023

Health Insurance Status	Home Smoking Policy	2011		2012		2013		2014		2015	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Insured	Smoking Ban	2436	78.1 (75.9, 80.3)	2491	78.9 (77.0, 80.9)	2654	82.1 (80.2, 84.0)	2783	84.5 (82.6, 86.4)	2426	84.3 (82.1, 86.6)
	No Ban	681	21.9 (19.7, 24.1)	663	21.1 (19.1, 23.0)	559	17.9 (16.0, 19.8)	495	15.5 (13.6, 17.4)	390	15.7 (13.4, 17.9)
Uninsured	Smoking Ban	270	64.2 (56.7, 71.6)	335	70.5 (65.0, 76.1)	298	73.1 (68.0, 78.3)	240	78.1 (72.6, 83.6)	149	71.8 (63.6, 79.9)
	No Ban	146	35.8 (28.4, 43.3)	142	29.5 (23.9, 35.0)	117	26.9 (21.7, 32.0)	91	21.9 (16.4, 27.4)	61	28.2 (20.1, 36.4)

Health Insurance Status	Home Smoking Policy	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
Insured	Smoking Ban	2569	86.0 (84.2, 87.9)	2160	85.3 (83.4, 87.2)	1782	85.1 (82.9, 87.3)	2159	87.8 (86.1, 89.4)	1524	88.3 (86.4, 90.2)
	No Ban	379	14.0 (12.1, 15.8)	362	14.7 (12.8, 16.6)	262	14.9 (12.7, 17.1)	302	12.2 (10.6, 13.9)	212	11.7 (9.8, 13.6)
Uninsured	Smoking Ban	200	73.3 (66.6, 80.0)	175	71.8 (64.0, 79.5)	150	69.3 (62.0, 76.6)	232	79.1 (73.4, 84.9)	175	82.8 (76.9, 88.7)
	No Ban	70	26.7 (20.0, 33.4)	58	28.2 (20.5, 36.0)	65	30.7 (23.4, 38.0)	57	20.9 (15.1, 26.6)	44	17.2 (11.3, 23.1)

Health Insurance Status	Home Smoking Policy	2021		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)
Insured	Smoking Ban	1761	88.2 (86.2, 90.2)	1919	87.6 (85.9, 89.3)
	No Ban	207	11.8 (9.8, 13.8)	238	12.4 (10.7, 14.1)
Uninsured	Smoking Ban	116	79.9 (72.3, 87.5)	117	84.3 (77.3, 91.2)
	No Ban	31	20.1 (12.5, 27.7)	20	15.7 (8.8, 22.7)

Table 11. Proportion of Oklahoma adults reporting smoking ban at home by disability status, Oklahoma BRFSS 2016 – 2021, 2023

Disability Status	Home Smoking Policy	2016		2017		2018		2019		2020	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)	N	Weighted% (95%CI)
One or more disability	Smoking Ban	979	75.8 (72.3, 79.2)	892	75.6 (72.0, 79.3)	726	75.8 (71.8, 79.7)	902	80.0 (76.9, 83.1)	613	82.4 (79.2, 85.7)
	No Ban	246	24.2 (20.8, 27.7)	221	24.4 (20.7, 28.0)	173	24.2 (20.3, 28.2)	199	20.0 (16.9, 23.1)	140	17.6 (14.3, 20.8)
No disability	Smoking Ban	1765	88.2 (86.0, 90.4)	1419	87.4 (85.1, 89.7)	1205	86.8 (84.3, 89.3)	1470	90.4 (88.5, 92.3)	1076	90.2 (87.9, 92.4)
	No Ban	197	11.8 (9.6, 14.0)	194	12.6 (10.3, 14.9)	154	13.2 (10.7, 15.7)	156	9.6 (7.7, 11.5)	113	9.8 (7.6, 12.1)

Disability Status	Home Smoking Policy	2021		2023	
		N	Weighted% (95%CI)	N	Weighted% (95%CI)
One or more disability	Smoking Ban	705	80.8 (77.2, 84.4)	766	82.0 (78.9, 85.1)
	No Ban	136	19.2 (15.6, 22.8)	143	18.0 (14.9, 21.1)
No disability	Smoking Ban	1171	90.9 (88.6, 93.2)	1267	90.8 (89.0, 92.7)
	No Ban	101	9.1 (6.8, 11.4)	119	9.2 (7.3, 11.0)