

ANALYSIS OF Diabetes in Oklahoma's SoonerCare Program



SoonerCare Members

From SFY2019 to SFY2020, the number of SoonerCare members diagnosed with diabetes decreased by 1.8%.


Type II continues to be the most prevalent type of diabetes among SoonerCare members (4.6% vs. 0.3% for Type I), although unspecified diabetes continued to increase in prevalence.

Total SoonerCare Enrollment SFY2011 - SFY2020



Diabetes Prevalence Rates SFY2011 - SFY2020

Diabetes Type: Total | Type I | Type II | Unspecified




In SFY2020 there were

56,586

members with diabetes, or 5.6% of all members

Total Prevalence Rate of Diabetes Type Among SoonerCare Members SFY2020



Members with Diabetes by Eligibility Type SFY2020




54.2%

were Dual Eligible*
30,667 members

**Dual Eligible: members enrolled in both Medicaid and Medicare*

Members with Diabetes by Delivery System SFY2020




65.2%

were enrolled in
SoonerCare Traditional
36,899 members


ANALYSIS OF Diabetes in Oklahoma's SoonerCare Program



Prevalence Rates



In SFY2020, the prevalence of diabetes was higher in eastern counties than in western counties. Prevalence of diabetes was highest in Okfuske (10.4%), Choctaw (9.5%), Tillman (8.3%), McIntosh, Johnston, and Seminole counties (all 8.6%).




SoonerCare Members with Diabetes

In SFY2020, diabetes was more prevalent among adults, females, Black/African Americans, and Caucasians than would be expected based on the demographic composition of all SoonerCare members. See details below. The black line represents the percent of total SoonerCare members.


95.6%

were adults
54,115 members




66.0%

were female
37,369 members



67.6%


were Caucasian
38,256 members



Prevalence Rates Among SoonerCare Members


13.2%

of all adult members had diabetes




6.6%

of all female members had diabetes



7.0%

of all Black or African American members had diabetes



Demographics

ANALYSIS OF Diabetes in Oklahoma's SoonerCare Program




Reimbursement Cost


While only 5.6% of SoonerCare members had diabetes in SFY2020, the total reimbursement associated with their services made up 16.7% of all SoonerCare reimbursements paid in SFY2020. The total reimbursement for members with diabetes increased by 7.6% from \$878,888,942 in SFY2019 to \$945,894,267 in SFY2020.

Members with Type I Diabetes had significantly higher reimbursement costs, on average, than other members with diabetes.


Annual Reimbursement for Members with Diabetes
SFY2011 - SFY2020



SoonerCare Members with Diabetes: Percent of Total Reimbursement Compared to Percent of Total SoonerCare Enrollment



Average Reimbursement per Member with Diabetes
Diabetes Type: Total | Type I | Type II | Unspecified



\$945,894,267

Total reimbursement for members with diabetes

\$16,716

Average reimbursement for members with diabetes

	Total Reimbursement	% of Total Reimbursement	Reimbursement per Member
Dual Eligible	\$419,805,722	44.4%	\$13,689
Non-Dual Eligible	\$526,088,545	55.6%	\$20,297
Adult	\$908,861,569	96.1%	\$16,795
Child	\$37,032,698	3.9%	\$14,987
Female	\$587,078,001	62.1%	\$15,710
Male	\$358,816,266	37.9%	\$18,672